

CP violation in D decays to two pseudoscalars: A SM-based calculation

Eleftheria Solomonidi

In collaboration with Antonio Pich & Luiz Vale Silva
Based on hep-ph/2305.11951 and upcoming publication

May 30

FPCP 2023, Lyon

Introduction


Charm Physics in the limelight

- Complementary to K and B Physics (CKM parameters) but different (*masses*)
- Experimental programme is growing (LHCb, Belle II, BESIII)

Rare decays

Mixing

CP violation in decays



- CPV in hadronic D modes: only discovery of CPV in the charm sector
- Plus new result of KK has puzzling implications

A new Flavour Physics 'anomaly' or an incomplete theory prediction?

$$\Delta A_{CP}^{exp} \equiv A_{CP}(D^0 \rightarrow K^+ K^-) - A_{CP}(D^0 \rightarrow \pi^+ \pi^-) = [-1.54 \pm 0.29] \cdot 10^{-3}$$

$$\Delta A_{CP}^{dir,exp} = [-1.57 \pm 0.29] \cdot 10^{-3} \quad \text{[LHCb 2019]}$$


$$A_{CP}(D^0 \rightarrow K^+ K^-) = [6.8 \pm 5.4(\text{stat}) \pm 1.6(\text{syst})] \cdot 10^{-4} \quad \text{[LHCb 2022]}$$

$$A_{CP}^{dir}(D^0 \rightarrow \pi^+ \pi^-) = [23.2 \pm 6.1] \cdot 10^{-4}$$

- Is the SM theoretical prediction in agreement?

Is it NP? [see e.g. 2210.16330]

- Weak sector (CKM parameters) probed by K & B physics



- Strong sector (hadronic uncertainties) problematic


Weak and strong, short and long distance

$$\mathcal{A}(D^0 \rightarrow f) = A(f) + i r_{CKM} B(f)$$


$$\mathcal{A}(\overline{D}^0 \rightarrow f) = A(f) - i r_{CKM} B(f)$$

$$a_{CP}^{dir} \approx 2 r_{CKM} \frac{|B(f)|}{|A(f)|} \cdot \sin \arg \frac{A(f)}{B(f)}$$

Tree topology



Penguin topology



From the short distance front:

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left[\sum_{i=1}^2 C_i(\mu) \left(\lambda_d Q_i^d(\mu) + \lambda_s Q_i^s(\mu) \right) - \lambda_b \left(\sum_{i=3}^6 C_i(\mu) Q_i(\mu) + C_{8g}(\mu) Q_{8g}(\mu) \right) \right]$$

$$\lambda_q = V_{cq}^* V_{uq}, \quad q = d, s, b.$$

$$|\lambda_d| \approx |\lambda_s| = \mathcal{O}(\lambda)$$

$$r_{CKM} = \text{Im} \frac{V_{cb}^* V_{ub}}{V_{cd}^* V_{ud}} \approx 6.2 \cdot 10^{-4}$$

Penguin operators

Current-current operators

$$Q_1^q = (\bar{q}c)_{V-A} (\bar{u}q)_{V-A}$$

$$Q_2^d = (\bar{q}_j c_i)_{V-A} (\bar{u}_i q_j)_{V-A}$$

$$(q = d, s)$$

$$Q_3 = (\bar{u}c)_{V-A} \Sigma_q (\bar{q}q)_{V-A}$$

$$Q_4 = (\bar{u}_j c_i)_{V-A} \Sigma_q (\bar{q}_i q_j)_{V-A}$$

$$Q_5 = (\bar{u}c)_{V-A} \Sigma_q (\bar{q}q)_{V+A}$$

$$Q_6 = (\bar{u}_j c_i)_{V-A} \Sigma_q (\bar{q}_i q_j)_{V+A}$$



$$C_{4,6} < 0.1 C_2, 0.03 C_1 \text{ (GIM mechanism at play)}$$

Weak and strong, short and long distance

$$\mathcal{A}(D^0 \rightarrow f) = A(f) + i r_{CKM} B(f)$$

$$\mathcal{A}(\overline{D}^0 \rightarrow f) = A(f) - i r_{CKM} B(f)$$

$$a_{CP}^{dir} \approx 2 r_{CKM} \frac{|B(f)|}{|A(f)|} \cdot \sin \arg \frac{A(f)}{B(f)}$$



From the short distance front:

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left[\sum_{i=1}^2 C_i(\mu) \left(\lambda_d Q_i^d(\mu) + \lambda_s Q_i^s(\mu) \right) - \lambda_b \left(\sum_{i=3}^6 C_i(\mu) Q_i(\mu) + C_{8g}(\mu) Q_{8g}(\mu) \right) \right]$$

$$\lambda_q = V_{cq}^* V_{uq}, \quad q = d, s, b.$$

$$|\lambda_d| \approx |\lambda_s| = \mathcal{O}(\lambda)$$

$$r_{CKM} = \text{Im} \frac{V_{cb}^* V_{ub}}{V_{cd}^* V_{ud}} \approx 6.2 \cdot 10^{-4}$$

Problem: hadronic matrix elements

$$\langle hh | Q_i | D^0 \rangle$$

Charm scale is special!

$$\Lambda_{\chi PT} \approx m_\rho < m_D = 1865 \text{ MeV}$$

$$\frac{\Lambda_{QCD}}{m_c} = \mathcal{O}(1)$$


See also: Khodjamirian, Petrov Phys. Lett. B, 774:235–242, 2017, Brod, Kagan, Zupan Phys. Rev. D, 86:014023, 2012, Schacht, Soni Phys. Lett. B,

825:136855, 2022

A way to look at the problem: rescattering

- Strong process, blind to the weak phase
- Isospin ($u \leftrightarrow d$) is a good symmetry of strong interactions
- In $I=0$, two channels:

$$S_{strong} = \begin{pmatrix} \pi\pi \rightarrow \pi\pi & \pi\pi \rightarrow KK \\ KK \rightarrow \pi\pi & KK \rightarrow KK \end{pmatrix}$$



Rescattering & what we learn about strong phases



- S matrix is **unitary**, as well as strong sub-matrix

- For $I=0$:
$$\begin{pmatrix} A(D \rightarrow \pi\pi) \\ A(D \rightarrow KK) \end{pmatrix} = \underbrace{\begin{pmatrix} \eta e^{i2\delta_1} & i\sqrt{1-\eta^2}e^{i(\delta_1+\delta_2)} \\ i\sqrt{1-\eta^2}e^{i(\delta_1+\delta_2)} & \eta e^{i2\delta_2} \end{pmatrix}}_{S_{\text{strong}}} \cdot \begin{pmatrix} A^*(D \rightarrow \pi\pi) \\ A^*(D \rightarrow KK) \end{pmatrix}$$

- The phases are related to the rescattering phases **which are known from data/nuclear experiments**
- Watson's theorem (elastic rescattering limit):
 $\arg A(D \rightarrow \pi\pi) = \arg A(\pi\pi \rightarrow \pi\pi) \bmod \pi$
- With inelasticities: more complicated, phase-shifts dependent on magnitudes of the amplitudes too

Magnitudes of matrix elements *without* rescattering

At the limit of $N_c \rightarrow \infty$, we are only left with the matrix elements from *factorisation*



(Same for $D \rightarrow KK$)

- Non-rescattering "bare" amplitudes:

$$T^B(D^0 \rightarrow \pi^+\pi^-) \propto \lambda_d C_1 \langle \pi^+\pi^- | Q_1 | D^0 \rangle_{fac} - \lambda_b (C_4 \langle \pi^+\pi^- | Q_4 | D^0 \rangle_{fac} + C_6 \langle \pi^+\pi^- | Q_6 | D^0 \rangle_{fac})$$

- Form factors are at the *non-rescattering* limit!
- Decay constant and form factor come from lattice and data (through χ PT)
- Internal gluon exchanges at each current are naturally included (but internal quark loops are suppressed)


Basic property of scattering amplitudes: analyticity

- Fundamental, model-independent property related to **causality**
- Cauchy's theorem:

$$A(s) = \frac{1}{2\pi i} \oint_C ds' \frac{A(s')}{s' - s} \text{ leads to}$$

$$\text{Re}A(s) = \frac{1}{\pi} \int_{s_{thr}}^{\infty} ds' \frac{\text{Im}A(s')}{s' - s}$$

(Dispersion relation)



- Unitarity of S-matrix & dispersion relation:

$$\underbrace{\text{Re}A(s)}_{\text{Re at a point}} = \frac{1}{\pi} \underbrace{\int_{s_{thr}}^{\infty} ds' \frac{\tan \delta(s')}{s' - s} \text{Re}A(s')}_{\text{integral of Re along the physical region}}$$

Analyticity & what we learn about magnitudes

- Integral equation, studied by **Muskhelishvili-Omnes**
- One subtraction: needs one piece of physical information
- **Single channel case** (& one subtraction at s_0), **physical** solution:

$$|A_I(s)| = \underbrace{A_I(s_0)}_{\text{ampl. when } \Omega = 1} \underbrace{\exp\left\{\frac{s-s_0}{\pi} PV \int_{4M_\pi^2}^{\infty} dz \frac{\delta_I(z)}{(z-s_0)(z-s)}\right\}}_{\text{Omnes factor } \Omega}$$

We need more than just large N_C !

$$|A_I(s = m_D^2)| = (\text{large } N_C \text{ result}) \times (\text{Omnes factor})_I$$

- More channels: Equally more solutions. **No analytical solution**


What we do

Summary of our method

- Factor out weak phases
- Flavour basis to isospin
- Isospin blocks:
 - $I=0$, unitarity with 2 channels: $\pi\pi$ and KK
 - $I=1$ with KK elastic rescattering
 - $I=2$ with $\pi\pi$ elastic rescattering
- Isospin amplitudes treated with dispersion relations calculated **numerically** (based on Moussallam et al. [hep-ph/9909292])
- Use inelasticity and phase-shift parametrisations [Pelaez et al., 1907.13162],[Pelaez et al., 2010.11222] up to energies $\sim m_D$ - extrapolate for higher & consider uncertainties
- For $I=1$ and 2, extract |Omnes factors| from Br's of $A(D^+ \rightarrow \pi^+\pi^0) \sim A_{I=2}, A(D^+ \rightarrow K^+\overline{K}^0) \sim A_{I=1}$, phases left unconstrained
- Decay-specific physical input: large N_C limit (for subtraction constant)

Choice of Omnes factors


For the isospin=0 channels we calculate numerically the Omnes matrix at $s = m_D^2$



	solution I	solution II	solution III
$\eta_0^0, \kappa_0^0 = 1$	$\Omega^{(0)} = \begin{pmatrix} 0.50 e^{+1.08i} & 1.01 e^{-1.03i} \\ 0.56 e^{-1.26i} & 0.29 e^{+2.87i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.39 e^{+1.64i} & 0.59 e^{-2.33i} \\ 0.51 e^{-1.25i} & 0.56 e^{+2.43i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.71 e^{+0.92i} & 1.35 e^{-0.87i} \\ 0.38 e^{-0.98i} & 0.42 e^{+0.85i} \end{pmatrix}$
$\eta_0^0 - \delta_0^0, \kappa_0^0 = 1$	$\Omega^{(0)} = \begin{pmatrix} 0.56 e^{+1.08i} & 0.61 e^{-1.73i} \\ 0.57 e^{-1.41i} & 0.58 e^{+2.85i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.42 e^{+1.75i} & 0.54 e^{-3.05i} \\ 0.51 e^{-1.25i} & 0.55 e^{+2.43i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.35 e^{+1.13i} & 0.74 e^{-0.87i} \\ 0.50 e^{-1.18i} & 0.55 e^{+2.48i} \end{pmatrix}$
$\eta_0^0 - \delta_0^0, \kappa_0^0 = 2$	$\Omega^{(0)} = \begin{pmatrix} 0.58 e^{+1.08i} & 0.64 e^{-1.74i} \\ 0.58 e^{-1.37i} & 0.61 e^{+2.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.43 e^{+1.54i} & 0.58 e^{-3.30i} \\ 0.52 e^{-1.05i} & 0.57 e^{+2.48i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.40 e^{+1.03i} & 0.80 e^{-0.30i} \\ 0.50 e^{-1.11i} & 0.50 e^{+2.53i} \end{pmatrix}$
$\eta_0^0 - \delta_0^0, \kappa_0^0 = 3$	$\Omega^{(0)} = \begin{pmatrix} 0.60 e^{+1.76i} & 0.66 e^{-1.74i} \\ 0.60 e^{-1.30i} & 0.63 e^{+2.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.44 e^{+1.58i} & 0.61 e^{-3.30i} \\ 0.52 e^{-1.17i} & 0.59 e^{+2.53i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.45 e^{+0.90i} & 0.80 e^{-0.53i} \\ 0.50 e^{-1.04i} & 0.57 e^{+2.58i} \end{pmatrix}$
sol. II: $ \eta_0^0 $	$\Omega^{(0)} = \begin{pmatrix} 2.03 e^{+1.39i} & 2.47 e^{-1.39i} \\ 0.37 e^{-0.38i} & 0.54 e^{+0.85i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 1.91 e^{+0.68i} & 2.78 e^{-2.35i} \\ 0.31 e^{-0.29i} & 0.45 e^{+0.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 2.20 e^{+0.63i} & 3.55 e^{-2.72i} \\ 0.35 e^{+0.03i} & 0.57 e^{+0.40i} \end{pmatrix}$
sol. C*: $ \eta_0^0 $	$\Omega^{(0)} = \begin{pmatrix} 1.83 e^{+1.39i} & 2.65 e^{-1.39i} \\ 0.34 e^{-0.48i} & 0.57 e^{+0.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 1.80 e^{+0.58i} & 3.11 e^{-2.36i} \\ 0.29 e^{-0.24i} & 0.49 e^{+0.84i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 2.09 e^{+0.32i} & 3.94 e^{-2.72i} \\ 0.32 e^{-0.03i} & 0.61 e^{+0.34i} \end{pmatrix}$



Choice of Omnes factors

For the isospin=0 channels we **calculate** numerically the Omnes matrix at $s = m_D^2$




	solution I	solution II	solution III
$\eta_0^0, \kappa_0^0 = 1$	$\Omega^{(0)} = \begin{pmatrix} 0.80 e^{+1.00i} & 1.01 e^{-1.00i} \\ 0.56 e^{-1.50i} & 0.29 e^{+0.87i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.39 e^{+1.04i} & 0.59 e^{-2.55i} \\ 0.51 e^{-1.50i} & 0.56 e^{+0.43i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.71 e^{+0.52i} & 1.35 e^{-0.87i} \\ 0.38 e^{-0.98i} & 0.42 e^{+0.85i} \end{pmatrix}$
$\eta_0^0 - \delta_0^0, \kappa_0^0 = 1$	$\Omega^{(0)} = \begin{pmatrix} 0.56 e^{+1.04i} & 0.61 e^{-1.75i} \\ 0.57 e^{-1.40i} & 0.58 e^{+0.85i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.42 e^{+1.75i} & 0.54 e^{-0.85i} \\ 0.51 e^{-1.50i} & 0.55 e^{+0.43i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.35 e^{+1.13i} & 0.74 e^{-0.87i} \\ 0.50 e^{-1.18i} & 0.55 e^{+0.48i} \end{pmatrix}$
$\eta_0^0 - \delta_0^0, \kappa_0^0 = 2$	$\Omega^{(0)} = \begin{pmatrix} 0.58 e^{+1.00i} & 0.64 e^{-1.74i} \\ 0.58 e^{-1.07i} & 0.61 e^{+0.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.43 e^{+1.54i} & 0.58 e^{-2.30i} \\ 0.52 e^{-1.05i} & 0.57 e^{+0.49i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.40 e^{+1.00i} & 0.80 e^{-0.30i} \\ 0.50 e^{-1.11i} & 0.50 e^{+0.53i} \end{pmatrix}$
$\eta_0^0 - \delta_0^0, \kappa_0^0 = 3$	$\Omega^{(0)} = \begin{pmatrix} 0.60 e^{+1.76i} & 0.66 e^{-1.74i} \\ 0.60 e^{-1.50i} & 0.63 e^{+0.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.44 e^{+1.50i} & 0.61 e^{-2.30i} \\ 0.52 e^{-1.17i} & 0.59 e^{+0.53i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 0.45 e^{+0.90i} & 0.80 e^{-0.55i} \\ 0.50 e^{-1.04i} & 0.57 e^{+0.58i} \end{pmatrix}$
sol. II: $ \delta_0^0 $	$\Omega^{(0)} = \begin{pmatrix} 2.03 e^{+1.50i} & 2.47 e^{-1.76i} \\ 0.37 e^{-0.30i} & 0.54 e^{+0.85i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 1.91 e^{+0.00i} & 2.78 e^{-2.55i} \\ 0.31 e^{-0.20i} & 0.45 e^{+0.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 2.20 e^{+0.00i} & 3.55 e^{-2.72i} \\ 0.35 e^{+0.00i} & 0.57 e^{+0.40i} \end{pmatrix}$
sol. C7: $ \delta_0^0 $	$\Omega^{(0)} = \begin{pmatrix} 1.83 e^{+1.50i} & 2.65 e^{-1.76i} \\ 0.34 e^{-0.80i} & 0.57 e^{+0.80i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 1.80 e^{+0.50i} & 3.11 e^{-2.36i} \\ 0.29 e^{-0.24i} & 0.49 e^{+0.24i} \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} 2.09 e^{+0.00i} & 3.94 e^{-2.72i} \\ 0.32 e^{-0.00i} & 0.61 e^{+0.34i} \end{pmatrix}$

- We examine the branching fraction predictions for the decays $\pi^+\pi^-$, $\pi^0\pi^0$, K^+K^- , $K^0\bar{K}^0$ based on each Omnes matrix separately
- Only a few of them give simultaneously correct Br values for all channels:





Choice of Omnes factors

For the isospin=0 channels we calculate numerically the Omnes matrix at $s = m_D^2$



	solution I	solution II	solution III
$\eta_0^0, \omega_0^0 = 1$	$\Omega^{(0)} = \begin{pmatrix} (0.90 \pm 1.00) & (1.01 \pm 1.00) \\ (0.50 \pm 1.00) & (0.50 \pm 2.07) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.39 \pm 1.60) & (0.50 \pm 2.55) \\ (0.51 \pm 1.30) & (0.56 \pm 2.45) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.71 \pm 0.52) & (1.35 \pm 0.47) \\ (0.38 \pm 0.90) & (0.42 \pm 0.45) \end{pmatrix}$
$\eta_0^0 - \delta\eta_0^0, \omega_0^0 = 1$	$\Omega^{(0)} = \begin{pmatrix} (0.50 \pm 1.80) & (0.61 \pm 1.75) \\ (0.57 \pm 1.41) & (0.58 \pm 2.00) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.42 \pm 1.75) & (0.54 \pm 2.40) \\ (0.51 \pm 1.30) & (0.55 \pm 2.45) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.35 \pm 1.13) & (0.74 \pm 0.47) \\ (0.50 \pm 1.10) & (0.55 \pm 0.48) \end{pmatrix}$
$\eta_0^0 - \delta\eta_0^0, \omega_0^0 = 2$	$\Omega^{(0)} = \begin{pmatrix} (0.58 \pm 1.80) & (0.64 \pm 1.74) \\ (0.58 \pm 1.37) & (0.61 \pm 2.00) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.43 \pm 1.60) & (0.58 \pm 2.30) \\ (0.52 \pm 1.35) & (0.57 \pm 2.40) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.40 \pm 1.00) & (0.80 \pm 0.30) \\ (0.50 \pm 1.11) & (0.50 \pm 0.53) \end{pmatrix}$
$\eta_0^0 - \delta\eta_0^0, \omega_0^0 = 3$	$\Omega^{(0)} = \begin{pmatrix} (0.00 \pm 1.70) & (0.66 \pm 1.74) \\ (0.00 \pm 1.30) & (0.63 \pm 2.20) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.44 \pm 1.50) & (0.61 \pm 2.30) \\ (0.52 \pm 1.17) & (0.59 \pm 2.35) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (0.45 \pm 0.90) & (0.90 \pm 0.33) \\ (0.50 \pm 1.00) & (0.57 \pm 0.58) \end{pmatrix}$
$\omega_0^0, \Omega^{(1)}[\eta_0^0]$	$\Omega^{(0)} = \begin{pmatrix} (2.01 \pm 1.20) & (2.47 \pm 1.70) \\ (0.37 \pm 0.52) & (0.54 \pm 2.00) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (1.91 \pm 0.00) & (2.78 \pm 2.30) \\ (0.31 \pm 0.23) & (0.45 \pm 2.30) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (2.20 \pm 0.00) & (3.55 \pm 0.72) \\ (0.35 \pm 0.02) & (0.57 \pm 0.40) \end{pmatrix}$
$\omega_0^0, \Omega^{(1)}[\eta_0^0]$	$\Omega^{(0)} = \begin{pmatrix} (1.83 \pm 1.20) & (2.65 \pm 1.70) \\ (0.34 \pm 0.40) & (0.57 \pm 2.00) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (1.90 \pm 0.50) & (3.11 \pm 2.30) \\ (0.29 \pm 0.24) & (0.49 \pm 2.34) \end{pmatrix}$	$\Omega^{(0)} = \begin{pmatrix} (2.09 \pm 0.02) & (3.94 \pm 0.72) \\ (0.32 \pm 0.02) & (0.61 \pm 0.34) \end{pmatrix}$

- We examine the branching fraction predictions for the decays $\pi^+\pi^-$, $\pi^0\pi^0$, K^+K^- , $K^0\bar{K}^0$ based on each Omnes matrix separately
- Only a few of them give simultaneously correct Br values for all channels:



Results

Rescattering quantified

With the branching fractions correctly reproduced

(old $D \rightarrow \pi\pi$, KK puzzle seems to be solved!)

the Omnes matrix looks like:

$$\Omega_{I=0} = \begin{pmatrix} 0.58e^{1.8i} & 0.64e^{-1.7i} \\ 0.58e^{-1.4i} & 0.61e^{2.3i} \end{pmatrix}$$

The **physical solution** is

$$\begin{pmatrix} \mathbf{A}(D \rightarrow \pi\pi) \\ \mathbf{A}(D \rightarrow KK) \end{pmatrix} = \Omega_{I=0} \cdot \begin{pmatrix} \mathbf{A}_{\text{factorisation}}(D \rightarrow \pi\pi) \\ \mathbf{A}_{\text{factorisation}}(D \rightarrow KK) \end{pmatrix}$$


It turns out:

Significant rescattering between the two final states!

penguin insertions \approx tree insertions

(of curr.-curr. operators, for $I=0$ reduced matrix elements)

CP asymmetries



charged meson channels
neutral meson channels

$$\Delta A_{CP}^{dir,exp} = (-1.57 \pm 0.29) \cdot 10^{-3}$$

With $\delta(I = 2, \pi\pi)$, $\delta(I = 1, KK)$
around the chosen values, we predict:

$$\Delta A_{CP}^{dir,theo} \sim 5 \cdot (10^{-4})!!$$

$$\text{and } a_{CP}^{dir}(D^0 \rightarrow \pi^+\pi^-) \approx 3 \cdot 10^{-4},$$


$$a_{CP}^{dir}(D^0 \rightarrow K^+K^-) \approx -2 \cdot 10^{-4}$$

$$a_{CP}^{dir} \approx 2 \underbrace{r_{CKM}}_{\sim 6 \cdot 10^{-4}} \underbrace{\frac{|B(f)|}{|A(f)|}}_{\sim 1/3} \cdot \underbrace{\sin \arg \frac{A(f)}{B(f)}}_{\sim 1}$$

NB: Short-distance penguins also not negligible for the CP asymmetries:
 C_6 small but annihilation insertion very large so that $C_6 \langle Q_6 \rangle_{fac} \sim C_1 \langle Q_1 \rangle_{fac}$

With fewer uncertain strong parameters (preliminary)

- $\pi\pi, KK$ inelasticity has large uncertainties
- Use only one low-energy strong phase for isospin 0: $\pi\pi + KK$ phase



- Assumption: 2-channel unitarity \rightarrow CPT/unitarity theorem also applying
- We manage to constrain:

$$0 < a_{CP}(\pi\pi)(0-0) \lesssim 5 \times 10^{-4}$$

$$-3 \times 10^{-4} \lesssim a_{CP}(KK)(0-0) < 0$$

- The CP asymmetry from $I = 2/0$ interference is not constrained, but would require very large values of isospin-0 Omnes matrix elements

Summary

- SM, data-driven approach that calculates the hadronic matrix elements deploying

Summary

- SM, data-driven approach that calculates the hadronic matrix elements deploying
 - 1 S-matrix unitarity and scattering amplitude analyticity; YES isospin, NO $SU(3)$ assumptions

Summary

- SM, data-driven approach that calculates the hadronic matrix elements deploying
 - ① S-matrix unitarity and scattering amplitude analyticity; YES isospin, NO $SU(3)$ assumptions
 - ② as much data as possible (rescattering, form factors and decay constants, Br's of D^+ decays)
- Strong rescattering data involves uncertainties → We keep the input that yields branching fractions **in reasonable agreement with experiment**

Summary

- SM, data-driven approach that calculates the hadronic matrix elements deploying
 - ① S-matrix unitarity and scattering amplitude analyticity; YES isospin, NO $SU(3)$ assumptions
 - ② as much data as possible (rescattering, form factors and decay constants, Br's of D^+ decays)
- Strong rescattering data involves uncertainties \rightarrow We keep the input that yields branching fractions **in reasonable agreement with experiment**
- **We still estimate the CP asymmetry for the $\pi^+\pi^-$ too small compared to the experimental value!**

Summary

- SM, data-driven approach that calculates the hadronic matrix elements deploying
 - ① S-matrix unitarity and scattering amplitude analyticity; YES isospin, NO $SU(3)$ assumptions
 - ② as much data as possible (rescattering, form factors and decay constants, Br's of D^+ decays)
- Strong rescattering data involves uncertainties \rightarrow We keep the input that yields branching fractions **in reasonable agreement with experiment**
- **We still estimate the CP asymmetry for the $\pi^+\pi^-$ too small compared to the experimental value!**
- CPV in $D^0 \rightarrow \pi^0\pi^0$ should be of similar magnitude (*could experiments look there?*)

Summary

- SM, data-driven approach that calculates the hadronic matrix elements deploying
 - ① S-matrix unitarity and scattering amplitude analyticity; YES isospin, NO $SU(3)$ assumptions
 - ② as much data as possible (rescattering, form factors and decay constants, Br's of D^+ decays)
- Strong rescattering data involves uncertainties \rightarrow We keep the input that yields branching fractions **in reasonable agreement with experiment**
- **We still estimate the CP asymmetry for the $\pi^+\pi^-$ too small compared to the experimental value!**
- CPV in $D^0 \rightarrow \pi^0\pi^0$ should be of similar magnitude (*could experiments look there?*)
- Future directions: different isospin-2 scenarios, more channels in isospin-0?
- But these are naively not expected to change the picture...

Thank you very much!

BACKUP

Isospin-2 and -1 fixing

$$\mathcal{A}(D^+ \rightarrow \pi^+ \pi^0) = \frac{3}{2\sqrt{2}} A_{I2}^\pi$$

$$\mathcal{A}(D^+ \rightarrow K^+ \overline{K}^0) = A_{I1}^K$$

We fix $|A_{I2}^\pi|$, $|A_{I1}^K|$ from the Br's and use them in e.g.

$$\mathcal{A}(D^0 \rightarrow \pi^+ \pi^-) = -\frac{1}{2\sqrt{3}} A_{I2}^\pi + \frac{1}{\sqrt{6}} A_{I0}^\pi$$

If $I=2$ elastic then $A_{I2}^\pi = \Omega_{I=2} A_{fac, I=2}$

If inelastic $A_{I2}^\pi = \Omega_{I=2} A_{fac, I=2} + (\text{mixing})$ but we use directly

$A_{I2}^\pi = |A_{I2}^\pi| \exp\{i\delta_{I=2}^{\pi\pi}\}$, phase left free

Naive estimate of final state interaction effects

$$\begin{pmatrix} A_{\pi\pi}^{I=0} \\ A_{KK}^{I=0} \end{pmatrix} = S_S^{1/2} \cdot \begin{pmatrix} A_{\pi\pi,\text{bare}}^{I=0} \\ A_{KK,\text{bare}}^{I=0} \end{pmatrix}$$

bare amplitudes: from factorisation (no strong phases)

Reproduces correctly Watson's theorem

What unitarity gives:

$$\begin{pmatrix} A_{\pi\pi}^{I=0} \\ A_{KK}^{I=0} \end{pmatrix} = S_S \cdot \begin{pmatrix} (A_{\pi\pi}^{I=0})^* \\ (A_{KK}^{I=0})^* \end{pmatrix}$$

No direct solution for the amplitudes, just relates them to the phases:

$$\arg A_{\pi\pi}^{I=0} = \delta_1 + \arccos \sqrt{\frac{(1+\eta)^2 - \left(\frac{|A_{KK}^{I=0}|}{|A_{\pi\pi}^{I=0}|}\right)^2 (1-\eta^2)}{4\eta}}$$

$$\arg A_{KK}^{I=0} = \delta_2 + \arccos \sqrt{\frac{(1+\eta)^2 - \left(\frac{|A_{\pi\pi}^{I=0}|}{|A_{KK}^{I=0}|}\right)^2 (1-\eta^2)}{4\eta}}$$

Numerical solution of 2-channel case

$$\begin{pmatrix} \text{Re}A^\pi(s) \\ \text{Re}A^K(s) \end{pmatrix} = \frac{s - s_0}{\pi} PV \int_{s_{thr}}^{\infty} ds' \frac{(\text{Re}T)^{-1}(\text{Im}T)(s')}{(s' - s)(s' - s_0)} \begin{pmatrix} \text{Re}A^\pi(s') \\ \text{Re}A^K(s') \end{pmatrix} + \begin{pmatrix} \text{Re}A_0^\pi(s_0) \\ \text{Re}A_0^K(s_0) \end{pmatrix}$$

- We discretise following the method from [Moussallam et al. hep-ph/9909292] into

$$\begin{pmatrix} \text{Re}A^\pi(s_i) \\ \text{Re}A^K(s_i) \end{pmatrix} = \frac{s_i - s_0}{\pi} \sum_j \hat{w}_j \frac{(\text{Re}T)^{-1}(\text{Im}T)(s_j)}{(s_j - s_i)(s_j - s_0)} \begin{pmatrix} \text{Re}A^\pi(s_j) \\ \text{Re}A^K(s_j) \end{pmatrix} + \begin{pmatrix} \text{Re}A_0^\pi(s_0) \\ \text{Re}A_0^K(s_0) \end{pmatrix}$$

- This creates an **invertible** matrix which gives a (discrete) solution
- Subtleties taken care of as in [Moussallam et al. hep-ph/9909292]
- To pick the *fundamental* solutions, we fix the vector at an unphysical point $s < 0$ and
 - check they behave as $\frac{1}{s}$ for large s
 - make sure the numerical determinant behaves as the (known) analytical determinant

Isospin decomposition

- $\pi\pi$ states can have isospin=0,2. KK can have isospin=0,1.

$$\begin{pmatrix} A(\pi^+\pi^-) \\ A(\pi^0\pi^0) \\ A(K^+K^-) \\ A(K^0\bar{K}^0) \end{pmatrix} = \begin{pmatrix} -\frac{1}{2\sqrt{3}} & -\frac{1}{\sqrt{6}} & 0 & 0 \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} & 0 & 0 \\ 0 & 0 & \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & -\frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} A_{\pi}^2 \\ A_{\pi}^0 \\ A_K^1 \\ A_K^0 \end{pmatrix}$$

$$\begin{pmatrix} A^\pi \\ A^K \end{pmatrix} = \begin{pmatrix} \Omega_{11} & \Omega_{12} \\ \Omega_{21} & \Omega_{22} \end{pmatrix} \begin{pmatrix} \text{Re}\lambda_d T^\pi + \dots \\ \text{Re}\lambda_s T^K + \dots \end{pmatrix}$$

$$\begin{pmatrix} B^\pi \\ B^K \end{pmatrix} = \begin{pmatrix} \Omega_{11} & \Omega_{12} \\ \Omega_{21} & \Omega_{22} \end{pmatrix} \begin{pmatrix} \text{Im}\lambda_d T^\pi + \sum_i \text{Im}\lambda_{d_i} P_i^\pi \\ \text{Im}\lambda_s T^K + \sum_i \text{Im}\lambda_{d_i} P_i^K \end{pmatrix}$$

Can consider either $\text{Im}\lambda_d = 0$ or $\text{Im}\lambda_s = 0$, not both simultaneously
 \Rightarrow In a_{CP}^{dir} there always exists a term $\sim T^\pi T^K$, both for $\pi\pi$ and for KK

Large N_C limit & effective operators

- $Q_1(i) = (\bar{d}_i c)_{V-A} (\bar{u} d_i)_{V-A}$, $Q_2(i) = (\bar{d} d)_{V-A} (\bar{u} c)_{V-A}$,
 $Q_{5,3} = (\bar{u} c)_{V-A} \sum_q (\bar{q} q)_{V\pm A}$,
 $Q_4 = \sum_q (\bar{u} q)_{V-A} (\bar{q} c)_{V-A}$, $Q_6 = -2 \sum_q (\bar{u} q)_{S+P} (\bar{q} c)_{S-P}$
- $C_1 = 1.18$, $C_2 = -0.32$, $C_3 = 0.011$, $C_4 = -0.031$, $C_5 = 0.0068$, $C_6 = -0.032$
($\mu = 2 \text{ GeV}$)
- $\lambda_d = V_{cd}^* V_{ud} \approx 0.22$
- $\overline{m}_c(2\text{GeV}) = 1.045\text{GeV}$
- Compare $m_D = 1865 \text{ MeV}$ to $\Lambda_{\chi PT} \approx m_\rho = 775 \text{ MeV}$