

# Pentaquarks at LHCb



*Ivan Polyakov*


*for LHCb collaboration*




*LHCb Implications Workshop,  
19-21 October 2022*

# The LHCb detector

- LHCb - forward spectrometer at LHC with excellent
  - momenta/mass,
  - vertex/time resolution
  - particle identification ( $K/\pi/p/\mu$ )



very powerful tool for heavy hadron spectroscopy  
 → contribute to major part of hadrons discovered at LHC



LHCb-FIGURE-2021-001

# First pentaquarks in $\Lambda_b \rightarrow J/\psi p K$


- In  $\Lambda_b \rightarrow J/\psi p K$  now 3 narrow pentaquarks are seen

PRL 115 (2015) 072001,  
PRL 122 (2019) 222001

State	$M$ [MeV]	$\Gamma$ [MeV]
$P_c(4312)^+$	$4311.9 \pm 0.7^{+6.8}_{-0.6}$	$9.8 \pm 2.7^{+3.7}_{-4.5}$
$P_c(4440)^+$	$4440.3 \pm 1.3^{+4.1}_{-4.7}$	$20.6 \pm 4.9^{+8.7}_{-10.1}$
$P_c(4457)^+$	$4457.3 \pm 0.6^{+4.1}_{-1.7}$	$6.4 \pm 2.0^{+5.7}_{-1.9}$

- Narrow (5-20 MeV)
- Note closeness to thresholds
- A wider state with  $M \sim 4380$  MeV and  $\Gamma \sim 200$  MeV to be confirmed with larger statistics
- Some of  $P_\psi$  are possibly seen in  $\Lambda_b \rightarrow J/\psi p \pi$

PRL 117 (2016) 082003





# $\Xi_b \rightarrow J/\psi \Lambda K$

Science Bulletin 66 (2021) 1278

- Use full Run1+2 (3+6fb<sup>-1</sup>) dataset
- Reconstruct  $\Lambda \rightarrow p\pi$  decay both in and outside of the VELO
- In total 1750 signal events, purity ~80%
- $\Xi^* \rightarrow \Lambda K$  contributions are clearly seen
- Full amplitude analysis firstly contributions in  $\Lambda K$  are examined

Run2 sample,  $\Lambda \rightarrow p\pi$  decay in VELO



State	$M_0$ (MeV)	$\Gamma_0$ (MeV)	$LS$ couplings	$J^P$ examined
$\Xi(1690)^-$	$1690 \pm 10$	$< 30$	4 (6)	$(1/2, 3/2)^\pm$
$\Xi(1820)^-$	$1823 \pm 5$	$24_{-10}^{+15}$	3 (6)	$3/2^-$
$\Xi(1950)^-$	$1950 \pm 15$	$60 \pm 20$	3 (6)	$(1/2, 3/2, 5/2)^\pm$
$\Xi(2030)^-$	$2025 \pm 5$	$20_{-5}^{+15}$	3 (6)	$5/2^\pm$
NR $\Lambda K^-$	-	-	4 (4)	$1/2^-$



# $\Xi_b \rightarrow J/\psi \Lambda K$ , amplitude

Science Bulletin 66 (2021) 1278

- 6D amplitude, follow helicity formalism

PRL 115 (2015) 072001

arXiv:1910.04566



$$\mathcal{M}_{\lambda_{\Xi_b}, \lambda_{\Lambda}, \Delta\lambda_{\mu}}^{\Xi_b^{*-}} \equiv \sum_n \sum_{\lambda_{\Xi_b^*}} \sum_{\lambda_{\psi}} \mathcal{H}_{\lambda_{\Xi_b^*}, \lambda_{\psi}}^{\Xi_b \rightarrow \Xi_b^* \psi} D_{\lambda_{\Xi_b}, \lambda_{\Xi_b^*} - \lambda_{\psi}}^{\frac{1}{2}}(0, \theta_{\Xi_b}, 0)^* \mathcal{H}_{\lambda_{\Lambda}, 0}^{\Xi_b^* \rightarrow K \Lambda} D_{\lambda_{\Xi_b^*}, \lambda_{\Lambda}}^{J_{\Xi_b^*}}(\phi_{\Lambda}, \theta_{\Xi_b^*}, 0)^* R_{\Xi_b^*}(m_{\Lambda K}) D_{\lambda_{\psi}, \Delta\lambda_{\mu}}^1(\phi_{\mu}, \theta_{\psi}, 0)^*$$

$$\mathcal{M}_{\lambda_{\Xi_b}, \lambda_{\Lambda}^{P_c}, \Delta\lambda_{\mu}^{P_c}}^{P_c^0} \equiv \sum_j \sum_{\lambda_{P_c^0}} \sum_{\lambda_{\psi}^{P_c}} \mathcal{H}_{\lambda_{P_c^0}, 0}^{\Xi_b \rightarrow P_c^0 K} D_{\lambda_{\Xi_b}, \lambda_{P_c^0}}^{\frac{1}{2}}(\phi_{P_c}, \theta_{\Xi_b}^{P_c}, 0)^* \mathcal{H}_{\lambda_{\psi}, \lambda_{\Lambda}^{P_c}}^{P_c^0 \rightarrow \psi \Lambda} D_{\lambda_{P_c^0}, \lambda_{\psi}^{P_c} - \lambda_{\Lambda}^{P_c}}^{J_{P_c^0}}(\phi_{\psi}, \theta_{P_c}, 0)^* R_{P_c^0}(m_{\psi \Lambda}) D_{\lambda_{\psi}, \Delta\lambda_{\mu}^{P_c}}^1(\phi_{\mu}^{P_c}, \theta_{\psi}^{P_c}, 0)^*$$




$$|\mathcal{M}|^2 = \sum_{\lambda_{\Xi_b}} \sum_{\lambda_{\Lambda}} \sum_{\Delta\lambda_{\mu}} \left| \mathcal{M}_{\lambda_{\Xi_b}, \lambda_{\Lambda}, \Delta\lambda_{\mu}}^{\Xi_b^{*-}} + e^{i\Delta\lambda_{\mu}\alpha_{\mu}} \sum_{\lambda_{\Lambda}^{P_c}} d_{\lambda_{\Lambda}^{P_c}, \lambda_{\Lambda}}^{\frac{1}{2}}(\theta_{\Lambda}) \mathcal{M}_{\lambda_{\Xi_b}, \lambda_{\Lambda}^{P_c}, \Delta\lambda_{\mu}}^{P_c^0} \right|^2$$

- $J^P$  determines allowed L, S values in  $\Xi_b \rightarrow P_{\psi} K$  and  $P_{\psi} \rightarrow J/\psi \Lambda$  decays and hence corresponding couplings:

$$H_{\lambda_B, \lambda_C}^{A \rightarrow BC} = \sum_L \sum_S \sqrt{\frac{2L+1}{2J_A+1}} B_{L,S} \left( \begin{array}{cc|c} J_B & J_C & S \\ \lambda_B & -\lambda_C & \lambda_B - \lambda_C \end{array} \right) \times \left( \begin{array}{cc|c} L & S & J_A \\ 0 & \lambda_B - \lambda_C & \lambda_B - \lambda_C \end{array} \right)$$

# $\Xi_b \rightarrow J/\psi \Lambda K$


- A need for one  $P_{\psi_s} \rightarrow J/\psi \Lambda$  was found
- Significance  $3.1\sigma$
- Two resonances are possible  
(analogous to  $P_{\psi}(4440)$  &  $P_{\psi}(4457)$ )
- $J^P$  examined are  $1/2^{\pm}, 3/2^{\pm}, 5/2^{\pm}$ , none is excluded



State	$M_0$ (MeV)	$\Gamma_0$ (MeV)	FF (%)
$P_{cs}(4459)^0$	$4458.8 \pm 2.9^{+4.7}_{-1.1}$	$17.3 \pm 6.5^{+8.0}_{-5.7}$	$2.7^{+1.9+0.7}_{-0.6-1.3}$

# $B_s \rightarrow J/\psi p \bar{p}$

- $B_{(s)}^0 \rightarrow J/\psi p \bar{p}$  decays were firstly observed in 2019
- Reanalyze the  $B_s$  decay with full Run1+2 data sample
- 800 signal events,  $\sim 85\%$  purity
- Amplitude fit:
  - no conventional intermediate states! thus only
    - $X \rightarrow p \bar{p}$ ,
    - $P_\psi^+ \rightarrow J/\psi p$
    - $P_\psi^- \rightarrow J/\psi \bar{p}$are considered on top of NR
- No B-tagging  $\rightarrow$   
 $P_\psi^+ \rightarrow J/\psi p$  and  $P_\psi^- \rightarrow J/\psi \bar{p}$  are fully symmetric



# $B_s \rightarrow J/\psi p \bar{p}$




Phys. Rev. Lett. 128 (2022) 062001

- No structures in  $p\bar{p}$  are seen
- Non-resonant proceeds with  $p\bar{p}$  in 1- (*S-waves in production & decay*)
- No evidence for  $P_\psi(4312) \rightarrow J/\psi p$  seen in  $\Lambda_b \rightarrow J/\psi p K$
- Found  $P_\psi \rightarrow J/\psi p$  with

$$M_{P_c} = 4337^{+7}_{-4} \text{ }^{+2}_{-2} \text{ MeV,}$$

$$\Gamma_{P_c} = 29^{+26}_{-12} \text{ }^{+14}_{-14} \text{ MeV,}$$

- Significances are  $3.1 - 3.7\sigma$
- $J^P$  examined are  $1/2^\pm, 3/2^\pm$  none is excluded







# $B \rightarrow J/\psi \Lambda \bar{p}$

- Previous amplitude analysis by CMS:
  - $B \rightarrow J/\psi \Lambda \bar{p}$  inconsistent with phase-space
  - can be explained with  $K^* \rightarrow \Lambda \bar{p}$  contributions

JHEP 12 (2019) 100



NEW!



# $B \rightarrow J/\psi \Lambda \bar{p}$

LHCb-PAPER-2022-031, in prep.

- Previous amplitude analysis by CMS:
  - $B \rightarrow J/\psi \Lambda \bar{p}$  inconsistent with phase-space
  - can be explained with  $K^* \rightarrow \Lambda \bar{p}$  contributions

JHEP 12 (2019) 100

- 4.6k signal events (x10 more than CMS had), 93% purity
- Reconstruct  $\Lambda \rightarrow p \pi$  decayed both in and outside of the VELO
- Amplitude analysis contributions:
  - non-resonant
  - $K^* \rightarrow \Lambda \bar{p}$
  - $P_\psi \rightarrow J/\psi \bar{p}$
  - $P_{\psi s} \rightarrow J/\psi \Lambda$




NEW!

# $B \rightarrow J/\psi \Lambda \bar{p}$

LHCb-PAPER-2022-031, in prep.

- Model with only  $NR(\Lambda \bar{p}) + K_4^*(2045), K_2^*(2250), K_3^*(2320)$  fails to describe data



**NEW!**




# $B \rightarrow J/\psi \Lambda \bar{p}$

LHCb-PAPER-2022-031, in prep.


- Add narrow  $P_{cs} \rightarrow J/\psi \Lambda$  state

$$M_{P_{cs}} = 4338.2 \pm 0.7 \pm 0.4 \text{ MeV} \text{ and } \Gamma_{P_{cs}} = 7.0 \pm 1.2 \pm 1.3 \text{ MeV}$$

- $15\sigma$  significance
- $1/2^-$  is preferred,  $1/2^+$  rejected at 90% CL
- Measured fit fractions:
  - $P_{\psi_s}$ :  $12.5 \pm 0.7 \pm 1.9$  %
  - NR( $J/\psi \bar{p}$ ):  $84.0 \pm 2.2 \pm 1.4$  %
  - NR( $\Lambda \bar{p}$ ):  $11.3 \pm 1.3 \pm 1.7$  %



# Discovered states




PRL 115 (2015) 072001,  
PRL 122 (2019) 222001

Science Bulletin 66 (2021) 1278  
Phys. Rev. Lett. 128 (2022) 062001

LHCB-PAPER-2022-031, in prep.

# Discovered states



	$J^P(X)$	$1/2^+$	$1/2^-$	$3/2^+$	$3/2^-$
$\Lambda_b/\Xi_b \rightarrow P_X K$	$1/2^+ \rightarrow X 0^-$	P-wave	S-wave	P-wave	D-wave
$B \rightarrow P_X \bar{p}$	$0^- \rightarrow X 1/2^+$	P-wave	S-wave	P-wave	D-wave

↑ suppressed due to low momenta in decay

# Not (yet) observations

- $\Lambda_b \rightarrow \eta_c[\rightarrow p\bar{p}]pK$ ,  $N_{\text{sig}} \sim 170$  PRD 102 (2020) 112012

$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow P_c(4312)^+ K^-) \times \mathcal{B}(P_c(4312)^+ \rightarrow \eta_c(1S)p)}{\mathcal{B}(\Lambda_b^0 \rightarrow \eta_c(1S)pK^-)} < 0.24$$

- $\Lambda_b \rightarrow \chi_{c1/2}[\rightarrow J/\psi\gamma]p\pi$ ,  $N_{\text{sig}} \sim 100/50$




- $\Lambda_b \rightarrow \chi_{c1/2}pK$ ,  $N_{\text{sig}} \sim 3100/1800$  JHEP 05 (2021) 095

- $\Lambda_b \rightarrow \chi_{c1}(3872)pK$ ,  $N_{\text{sig}} \sim 50$  JHEP 09 (2019) 028

- $\Lambda_b \rightarrow \Lambda_c p\bar{p}\pi$ ,  $N_{\text{sig}} \sim 900$  (only Run1) PLB 784 (2018) 101

- $\Lambda_b \rightarrow J/\psi p\pi$ ,  $N_{\text{sig}} \sim 2100$  (only Run1) PRL 117 (2016) 082003

*signals will likely appear with more statistics*



# Prospects

- More potential decay modes to look at (ideally all charged tracks):

- $\Lambda_b \rightarrow \chi_{c1} p K, \dots, \Lambda_b \rightarrow \eta_c p K, \dots$

- $\Lambda_b \rightarrow J/\psi p K_s \pi^-, \Lambda_b \rightarrow J/\psi \Lambda \pi^+ \pi^-, \Xi_b \rightarrow J/\psi \Lambda K^+ \pi^-(\pi^-), \dots$

- $B_s \rightarrow J/\psi \Lambda \bar{\Lambda}$

- $\Lambda_b/\Xi_b \rightarrow P_\psi [\rightarrow \Lambda_c \bar{D} \pi, \Xi_c \bar{D}, \dots] K, \dots$


- $P_\psi$  in prompt pp-collisions

*! can't promise we'll have enough statistic*

- Although... Run3(4) will give up to x5 (x10 for Run3&4) boost in statistics wrt current dataset



# Summary



- Latest results on pentaquarks:

- $P_{\psi s} \rightarrow J/\psi \Lambda$  in  $\Xi_b \rightarrow J/\psi \Lambda K$

Science Bulletin 66 (2021) 1278

- $P_\psi \rightarrow J/\psi p$  in  $B_s \rightarrow J/\psi p \bar{p}$

Phys. Rev. Lett. 128 (2022) 062001

- $P_{\psi s} \rightarrow J/\psi \Lambda$  in  $B \rightarrow J/\psi \Lambda \bar{p}$

LHCB-PAPER-2022-031, in prep.

- Stay tuned


# Appendix

# Interpretations

- Fit in both model frameworks:
  - compact multiquark**
    - genuine QCD state
    - size  $\sim 1\text{fm}$
  - molecular state**
    - two hadrons bound by  $\pi/\rho/\eta$  (QCD analog of “van der Waals” force)
    - are well separated (1-10fm)
    - natural closeness to thresholds
- Both suggest more of analogous states

see Richard, arxiv:1606.08593;  
Esposito, Pilloni, Polosa, arXiv:1611.07920

see Guo et al, arXiv:1705.00141




credit: Moonrunner Design

# Hope to learn from you

# Upgraded LHCb

- Major upgrades during last shutdown
  - Tracking&Vertexing
  - PID
  - Trigger



- Started to collect data in Run3 (2022-2025)
- Will give up to x5 (x10 for Run3&4) boost in statistics wrt current dataset
- More exciting results will follow

# $\Xi_b \rightarrow J/\psi \Lambda K$ , systematics

Science Bulletin 66 (2021) 1278

- Systematic uncertainties:
  - $J^P$  assignments for  $\Xi^*$  states
  - higher/all L, different d in Blatt-Weisskopf, different NR for  $\Lambda K$ , more  $\Xi^*$ ,
  - adding  $\Lambda \rightarrow p\pi$  helicity angle to consideration in amplitude
  - splitting into bins of  $\Xi^* \rightarrow \Lambda K$  helicity angle
  - or/and removing  $\Xi_b \rightarrow J/\psi \Sigma (\rightarrow \Lambda \gamma) K$  from sideband
  - limited statistics of simulation sample via efficiency

Source	$P_{cs}(4459)^0$			$\Xi(1690)^-$			$\Xi(1820)^-$			$\Xi^{*-}$ (1950)	$\Xi^{*-}$ (2030)	$NR$
	$M_0$	$\Gamma_0$	FF	$M_0$	$\Gamma_0$	FF	$M_0$	$\Gamma_0$	FF	FF	FF	FF
$J^P$	+4.7	+0.0	+0.1	+1.2	+14.0	+6.7	+0.8	+1.4	+4.2	+ 0.2	+0.0	+ 0.9
	-0.3	-5.7	-1.3	-0.1	- 0.9	-0.3	-0.2	-0.5	-0.3	- 9.4	-4.1	-11.2
Model	+0.7	+8.0	+0.7	+0.5	+ 1.8	+1.9	+1.0	+7.8	+6.9	+49.9	+3.8	+10.3
	-1.1	-2.0	-0.5	-0.4	-13.5	-8.9	-0.6	-8.2	-4.1	- 5.4	-1.6	- 6.4
$\Lambda$ decay	+0.0	+0.0	+0.0	+0.0	+ 0.2	+0.0	+0.0	+0.0	+0.0	+ 2.4	+0.0	+ 3.9
	-0.7	-4.7	-0.3	-0.4	- 0.0	-0.8	-0.5	-7.2	-4.1	- 0.0	-1.3	- 0.0
sWeights	+0.0	+0.3	+0.1	+0.1	+ 3.1	+1.4	+0.2	+2.2	+1.6	+ 0.7	+0.0	+ 0.0
	-0.2	-0.0	-0.0	-0.1	- 0.2	-0.0	-0.2	-1.5	-0.5	- 1.6	-0.2	- 2.7
Efficiency	+0.1	+0.0	+0.0	+0.1	+ 2.1	+0.8	+0.1	+1.1	+0.5	+ 2.3	+0.3	+ 1.1
	-0.1	-0.5	-0.1	-0.2	- 1.5	-1.3	-0.2	-0.3	-0.7	- 1.0	-0.2	- 0.9
Final	+4.7	+8.0	+0.7	+1.2	+14.0	+6.7	+1.0	+7.8	+6.9	+49.9	+3.8	+10.3
	-1.1	-5.7	-1.3	-0.4	-13.5	-8.9	-0.6	-8.2	-4.1	- 9.4	-4.1	-11.2

Table 3: Summary of absolute systematic uncertainties for the fit parameters. The units for masses ( $M_0$ ) and widths ( $\Gamma_0$ ) are MeV. The fit fraction in percent is denoted FF.

# $B_s \rightarrow J/\psi p \bar{p}$ , systematics

Phys. Rev. Lett. 128 (2022) 062001

- Estimated with pseudo-experiments generated according to alternative model, fit with baseline model

Source	$M_{P_c}$	$\Gamma_{P_c}$	$A(P_c)$	$f(P_c)$	$p$ (%)	$\sigma$
NR( $X$ ) model	0.1	1.4	0.013	6.4	0.003	4.2
$J^P(P_c)$ assignment	2	12	0.100	5.5	0.2	3.1
Efficiency	0.2	4	0.012	0.4	0.001	4.4
Background	0.1	2	0.001	0.7	0.001	4.3
Hadron radius	0.7	4	0.034	1.7	0.02	3.7
Fit bias	+0.2 -0.1	+5 -2	+0.040 -0.040	—	—	—
Total	2	14	0.11	8.6	—	3.1

**NEW!**

# $B \rightarrow J/\psi \Lambda \bar{p}$ , systematics

LHCB-PAPER-2022-031, in prep.

Source	$M_{P_{\psi s}^{\Lambda}}$	$\Gamma_{P_{\psi s}^{\Lambda}}$	$f_{P_{\psi s}^{\Lambda}}$	$f_{NR}(J/\psi \bar{p})$	$f_{NR}(\Lambda \bar{p})$
Hadron radius	0.1	0.4	0.3	0.2	0.2
$LS$ values	0.3	0.1	0.8	0.7	0.6
Breit–Wigner $P_{\psi}^{N-}$	0.1	0.9	0.8	...	...
$J^P(P_{\psi s}^{\Lambda})$ assignment	0.1	0.9	1.2	0.4	0.9
Fitting procedure	0.1	0.2	0.1	1.0	1.1
Efficiency	0.02	0.19	0.02	0.3	0.2
$\Lambda$ decay parameters	0.02	0.04	0.01	0.3	0.2
Background	0.01	0.05	0.96	0.4	0.7
Mass resolution	0.01	0.03	0.01	0.1	0.1
Total	0.4	1.3	1.9	1.4	1.7

# New naming convention proposed

arxiv:2206.15233

## round table (29 sept)

- To bring more order in the fast-growing list of exotic hadrons
- Preserve minimal change to existing names
- Create framework for future discoveries

Table 5: Summary of the impact of the exotic hadron naming scheme on various states, based on current knowledge of their properties. Quantum numbers that are not specified or marked “?” are unknown and the corresponding super-/sub-scripts not given. The current name indicated is that used in the PDG listings [16].

Minimal quark content	Current name	$I^{(G)}, J^{P(C)}$	Proposed name	Reference
$c\bar{c}$	$\chi_{c1}(3872)$	$I^G = 0^+, J^{PC} = 1^{++}$	$\chi_{c1}(3872)$	[24, 25]
$c\bar{c}u\bar{d}$	$Z_c(3900)^+$	$I^G = 1^+, J^P = 1^+$	$T_{\psi 1}^b(3900)^+$	[26–28]
$c\bar{c}u\bar{d}$	$X(4100)^+$	$I^G = 1^-$	$T_{\psi}(4100)^+$	[29]
$c\bar{c}u\bar{d}$	$Z_c(4430)^+$	$I^G = 1^+, J^P = 1^+$	$T_{\psi 1}^b(4430)^+$	[30, 31]
$c\bar{c}(s\bar{s})$	$\chi_{c1}(4140)$	$I^G = 0^+, J^{PC} = 1^{++}$	$\chi_{c1}(4140)$	[32–35]
$c\bar{c}u\bar{s}$	$Z_{cs}(4000)^+$	$I = \frac{1}{2}, J^P = 1^+$	$T_{\psi s 1}^{\theta}(4000)^+$	[7]
$c\bar{c}u\bar{s}$	$Z_{cs}(4220)^+$	$I = \frac{1}{2}, J^P = 1^?$	$T_{\psi s 1}(4220)^+$	[7]
$c\bar{c}c\bar{c}$	$X(6900)$	$I^G = 0^+, J^{PC} = ?^{?+}$	$T_{\psi\psi}(6900)$	[4]
$cs\bar{u}\bar{d}$	$X_0(2900)$	$J^P = 0^+$	$T_{cs 0}(2900)^0$	[5, 6]
$cs\bar{u}\bar{d}$	$X_1(2900)$	$J^P = 1^-$	$T_{cs 1}(2900)^0$	[5, 6]
$cc\bar{u}\bar{d}$	$T_{cc}(3875)^+$		$T_{cc}(3875)^+$	[8, 9]
$bb\bar{u}\bar{d}$	$Z_b(10610)^+$	$I^G = 1^+, J^P = 1^+$	$T_{\Upsilon 1}^b(10610)^+$	[36]
$c\bar{c}uud$	$P_c(4312)^+$	$I = \frac{1}{2}$	$P_{\psi}^N(4312)^+$	[3]
$c\bar{c}uds$	$P_{cs}(4459)^0$	$I = 0$	$P_{\psi s}^A(4459)^0$	[20]