


Combined analysis of meson mixings and EW precision observables in SM4

Nejc Kosnik
(in collaboration with Emi Kou)



Institut "Jožef Stefan", Ljubljana, Slovenija

Presented at "Portoroz 2011: The role
of heavy fermions in fundamental
physics"

13.4.2011

Motivation for another replication of fermions



- ◆ Relax tension with m_H direct bound
- ◆ 2 new CP violating phases for electroweak baryogenesis [Hou] $n_B/s \simeq 5 \times 10^{-10}$
- ◆ Large Yukawas of new fermions - possible dynamical explanation of EWSB

- ◆ Neutrinos are heavier than $m_Z/2$

- ◆ Current bounds

$m_{t'}$	> 338 GeV	CDF
$m_{b'}$	> 361 GeV	CMS
$m_{\ell'}$	> 100 GeV	LEP
$m_{\nu'}$	> 90 GeV	

Bounds depend on partial BR's, thus on CKM, PMNS



Framework

Fourth generation masses run as

$$300 \text{ GeV} < m_{t'}, m_{b'} < 600 \text{ GeV}$$

Higgs mass fixed at 117 GeV at this stage

$$100 \text{ GeV} < m_{\nu'}, m_{\ell'} < 600 \text{ GeV}$$

Higher values are perfectly possible but without improvement in the overall agreement with observables

↑
Direct lower bounds
from LEP, Tevatron, LHC

↑
Yukawa couplings
perturbativity

CKM matrix contains 3 new angles and 2 new phases

$$\begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) & A\lambda^3(\rho_1 - i\eta_1) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 & A\lambda^2(\rho_2 - i\eta_2) \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 - A^2\rho_3^2\lambda^2/2 & A\rho_3\lambda \\ A\lambda^3(\rho_1 - \rho_2 - i\eta_1 + i\eta_2) & -A\lambda^2(\rho_2 + i\eta_2) & -A\rho_3\lambda & 1 - (A\rho_3)^2\lambda^2/2 \end{pmatrix} + \mathcal{O}(\lambda^3)$$

Cabibbo angle power counting inspired by 3x3 unitarity measurements

$$|V_{ub'}|^2 = 1 - |V_{ud}|^2 - |V_{us}|^2 - |V_{ub}|^2 = 0.00001 \pm 0.0011,$$

$$|V_{cb'}|^2 = 1 - |V_{cd}|^2 - |V_{cs}|^2 - |V_{cb}|^2 = -0.002 \pm 0.027,$$

$$|V_{tb'}|^2 < 1 - |V_{tb}|^2, \quad |V_{tb}| = 0.88 \pm 0.07.$$

Electroweak precision observables

Oblique parameters: effect of heavy fermions in gauge boson self-energies


$$S_4 = \frac{2}{3\pi} \left[1 - \frac{1}{2} \log \frac{m_{t'}}{m_{b'}} - \frac{1}{2} \log \frac{m_{\ell'}}{m_{\nu'}} \right]$$

$$T_4 = \frac{3}{16\pi x_w (1-x_w) m_Z^2} [m_{t'}^2 + m_{b'}^2 - |V_{t'b'}|^2 F_T(m_{t'}^2, m_{b'}^2)]$$

$$+ \frac{1}{16\pi x_w (1-x_w) m_Z^2} [m_{\ell'}^2 + m_{\nu'}^2 - F_T(m_{\ell'}^2, m_{\nu'}^2)]$$

$$U_4 \simeq 0 \quad (\text{Lenz; Chanowitz}) \quad \mathfrak{S}$$


$$V_{t'b'} \approx 1 - V_{tb'}^2/2 \approx 1 - A^2 \rho_3^2 \lambda^2/2$$



Corresponds to points inside the 2σ contour in the S-T plane

$$\text{Cov}_{S,T} = \begin{pmatrix} 0.0081 & 0.006264 \\ 0.006264 & 0.0064 \end{pmatrix}$$

[S,T from Erler, Langacker]



Preferred small 3-4 mixing ($V_{tb'}$)

$$\begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) & A\lambda^3(\rho_1 - i\eta_1) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 & A\lambda^2(\rho_2 - i\eta_2) \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 - A^2 \rho_3^2 \lambda^2/2 & A\rho_3 \lambda \\ A\lambda^3(\rho_1 - \rho_2 - i\eta_1 + i\eta_2) & -A\lambda^2(\rho_2 + i\eta_2) & -A\rho_3 \lambda & 1 - (A\rho_3)^2 \lambda^2/2 \end{pmatrix}$$

Electroweak precision observables


Zbb nondecoupling vertex correction

$$R_b \equiv \Gamma(Z \rightarrow b\bar{b}) / \Gamma(Z \rightarrow \text{hadrons})$$

$$\Gamma(Z \rightarrow b\bar{b}) = \#m_Z(1 + \delta_b) \quad (\text{Bernabeu; Yanir})$$

$$\delta_b \approx 10^{-2} \left[\left(-\frac{m_t^2}{2m_Z^2} + 0.2\right)|V_{tb}|^2 + \left(-\frac{m_{t'}^2}{2m_Z^2} + 0.2\right)^2|V_{t'b}|^2 \right]$$

Probes 3-4 mixing and $m_{t'}$, ρ_3 must not be too large




V_{tb} measurement in single top production

not EWP per-se but directly probes ρ_3

$$V_{tb} = 1 - (A\rho_3\lambda)^2/2 \quad V_{tb} = 0.88 \pm 0.07$$

Requires ρ_3 not too small.



Meson mixing observables

Probe of CKM and mass scales

$$\begin{pmatrix}
 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) & A\lambda^3(\rho_1 - i\eta_1) \\
 -\lambda & 1 - \lambda^2/2 & A\lambda^2 & A\lambda^2(\rho_2 - i\eta_2) \\
 A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 - A^2\rho_3^2\lambda^2/2 & A\rho_3\lambda \\
 A\lambda^3(\rho_1 - \rho_2 - i\eta_1 + i\eta_2) & -A\lambda^2(\rho_2 + i\eta_2) & -A\rho_3\lambda & 1 - (A\rho_3)^2\lambda^2/2
 \end{pmatrix} \Delta m_D$$

ϵ_K $\Delta m_s \sin 2\beta_s$ $\Delta m_d \sin 2\beta$

$(m_{t'})$ ← EWP → $(m_{b'})$


K and B mixing observables are theoretically reliable and sensitive to $m_{t'}$, e.g.

$$\xi_{sd}^2 \frac{m_{B_s}}{m_{B_d}} \left| \frac{\eta_t(V_{tb}V_{ts}^*)^2 S_0(x_t) + \eta_{t'}(V_{t'b}V_{t's}^*)^2 S_0(x_{t'}) + 2\eta_{tT}V_{tb}V_{ts}^*V_{t'b}V_{t's}^* S_0(x_t, x_{t'})}{\eta_t(V_{tb}V_{td}^*)^2 S_0(x_t) + \eta_{t'}(V_{t'b}V_{t'd}^*)^2 S_0(x_{t'}) + 2\eta_{tT}V_{tb}V_{td}^*V_{t'b}V_{t'd}^* S_0(x_t, x_{t'})} \right| = \frac{\Delta m_s}{\Delta m_d} \Big|_{\text{exp}}$$

D meson mixing is sensitive to $m_{b'}$, however difficult to assign probabilistic significance due to long-distance background

A global view

$m_t, m_{b'}, m_{\text{leptons}}, \rho_3$



$m_{b'}$
 $\rho_1, \eta_1, \rho_2, \eta_2$

CKM4

m_t
 $\rho_1, \eta_1, \rho_2, \eta_2, \rho_3$

Global fit

To quantify the impact of meson mixing observables.

Similar analyses done by [Dighe 2010, Lenz 2010]

Observables (15): $S = 0.03 \pm 0.09$, $T = 0.07 \pm 0.08$ EWP and/or driven by ρ_3
 $R_b = 0.216 \pm 0.001$
 $V_{tb} = 0.88 \pm 0.07$

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \end{pmatrix} \gamma$$

Tree-level quantities, primarily sensitive to CKM3 (PDG values)

$$\epsilon_K, \sin 2\beta$$
$$\Delta m_s, \frac{\Delta m_s}{\Delta m_d}$$

FCNC observables, very sensitive to new CKM parameters

Write down Gaussian χ^2 for each observable as $[o(y) - o_{\text{exp}}]^2 / \sigma^2$

Mass splitting in charm sector is treated as a “kinematical” constraint

$$|M_{12,D}^{b'}| < 3|M_{12,D}^{\text{exp}}|$$

Note that “3” is arbitrary (Lenz; Golowich)

Global fit parameters

Model parameters (13): $m_{t'}$, $m_{b'}$, $m_{\nu'}$, $m_{\ell'}$ λ , A , ρ , η , $\rho_{1,2,3}$, $\eta_{1,2}$

Theoretical (nuisance) parameters:

$$\eta_c = 1.43(23)$$

$$\eta_{ct} = 0.47(4)$$

$$\eta_t = 0.5765(65)$$

$$\hat{B}_K = 0.725(26)$$

$$f_K = 156.1(8) \text{ MeV}$$

$$\kappa_\epsilon = 0.94(2)$$

$$\eta_B = 0.55(1)$$

$$\xi = 1.237(32)$$

$$f_{B_s} \sqrt{\hat{B}_{B_s}} = 270(30) \text{ MeV} \quad [\text{Lubicz, Tarantino}]$$

Theoretical parameters freely slide within their allowed ranges and do not contribute to χ^2 . Similar to CKMFitter's RFit, except that we do not add statistical tail. (preliminary)

D mixing theoretical parameters' errors are irrelevant when compared to arbitrariness in interpretation of the experimental Δm_D .

Interpretation of fit results

1. Global minimum of χ^2 , χ^2_{\min} , determines the overall quality of the fit.

$$n_{\text{DOF}} = 15 - 13 = 2 \text{ degrees of freedom}$$

$$\begin{aligned} \chi^2_{n_{\text{DOF}}=2} &\leq 2.3 && 1\sigma \\ &\leq 6.2 && 2\sigma \\ &\leq 11.8 && 3\sigma \end{aligned}$$

Assumption of parabolic (Gaussian) behavior around minimum. To improve, resort to MC and determine confidence levels pseudoexperimentally.

2. Assuming model is correct, we find allowed range of its model parameter “ y_1 ” by considering

$$\Delta\chi^2(y_1) = \min_{\{y_2, y_3, \dots\}} [\chi^2(y_1, y_2, \dots) - \chi^2_{\min}]$$

$\Delta\chi^2(y_1)$ at “best” value of y_1 is 0,

$$N\text{-}\sigma \text{ region} = \{y_1 ; \Delta\chi^2(y_1) \leq N^2\}$$


Global minimum

(preliminary)

$$\chi^2_{\min} = 8.60$$

$$= 2.84 + 2.14 + 1.89 + 1.46 + \dots$$

$$V_{tb} \quad R_b \quad V_{cs} \quad S, T$$



2.5 σ fluctuation

Significance of deviation is expected to decrease once we include additional observables.

$$m_{t'} \approx 325 \text{ GeV}$$

$$m_{b'} \approx 305 \text{ GeV}$$

$$m_{\nu'} \approx 100 \text{ GeV}$$

$$m_{\ell'} \approx 190 \text{ GeV}$$

$$\lambda \approx 0.22515$$

$$A \approx 0.802$$

$$\rho \approx 0.14$$

$$\eta \approx 0.40$$

$$\rho_1 \sim 0.3$$

$$\eta_1 \sim 1.4$$



$$\rho_2 \sim -0.1$$

$$\eta_2 \sim 0.3$$

$$\rho_3 \sim 0.3$$

Mass splittings (see also Lenz's talk)

4th generation doublets can both be degenerate




Compare to commonly used optimal point, without CKM angles


$$m_{\ell'} - m_{\nu'} \approx 30 - 60 \text{ GeV}$$

$$m_{t'} - m_{b'} \approx 50 \text{ GeV}$$

[Kribs, 2007]




CKM elements predictions



$$1 - (A\rho_3\lambda)^2/2, \quad 1 - \lambda^2/2 \sim 0.97$$


Expansion of CKM stable.



$$A\lambda^2(\rho_2 + i\eta_2)$$

Important for direct searches,
assuming $\text{Br}(t' \rightarrow b W) = 1$

Prediction of $\sin 2\beta_s$



Conclusion

- ◆ EW data favors splitting in both quark and leptonic sectors
- ◆ Crucial degree of freedom is the 3-4 mixing, allowing much wider range of masses and splittings, and opening portal to flavor physics
- ◆ Flavor observables are coupled to EW observables via 3-4 mixing and quark masses.

Conclusion

- ◆ Minimal set of relevant observables ($n_{\text{DOF}} = 2$) strongly constrains CKM elements.
- ◆ Phase in B_s mixing is constrained
- ◆ Study of constraints in $(m_{t'}, V_{t'q})$ and $(m_{b'}, V_{qb'})$ planes is underway