



Version 11.0

Electromagnetic Physics

Vladimir Ivantchenko (CERN & Princeton University)
Geant4 Advanced Course




- Electromagnetic physics (EM) overview
- Main Gamma processes
- Main charged particle processes
- Secondary production thresholds
- EM physics constructors
- User interface to EM physics
- Special EM topics


ELECTROMAGNETIC (EM) PHYSICS OVERVIEW

- Photon processes
 - γ conversion into e+e- pair
 - Compton scattering
 - Photoelectric effect
 - Rayleigh scattering
 - *Gamma-nuclear interaction in hadronic sub-library*
- Electron and positron processes
 - Ionization
 - Coulomb scattering
 - Bremsstrahlung
 - Production of e+e- pair
 - *Nuclear interaction in hadronic sub-library*
 - Positron annihilation
- Suitable for HEP & many other Geant4 applications with electron and gamma beams

HEP calorimeter



Medical linac




Located in \$G4INSTALL/sources/processes/electromagnetic

- **Standard**
 - γ , e up to 100 TeV
 - hadrons up to 100 TeV
 - ions up to 100 TeV
- **Muons**
 - up to 1 PeV
 - energy loss propagator
- **X-rays**
 - X-ray and optical photon production processes
- **High-energy**
 - processes at high energy ($E > 10 \text{ GeV}$)
 - physics for exotic particles
- **Polarisation**
 - simulation of polarised beams
- **Optical**
 - optical photon interactions
- **Low-energy**
 - Livermore library γ , e- from 10 eV up to 1 GeV
 - Livermore library based polarized processes
 - PENELOPE 2008 code rewrite , γ , e- , e+ from 250 eV up to 6 GeV
 - hadrons and ions up to 1 GeV
 - atomic de-excitation (fluorescence + Auger)
- **DNA**
 - Geant4 DNA modes and processes
 - Micro-dosimetry models for radiobiology
 - from 0.025 eV to 10 MeV
 - many of them material specific (water)
 - Chemistry in liquid water
- **Adjoint**
 - sub-library for reverse Monte Carlo simulation from the detector of interest back to source of radiation
- **Utils** : general EM interfaces and helper classes

- The uniform coherent approach for all EM packages
 - low energy and high energy models may work together
- A physical interaction or process is described by a process class
 - For example: **G4ComptonScattering**
 - Assigned to Geant4 particle types in Physics List
 - Three EM base processes:
 - **G4VEmProcess**
 - **G4VEnergyLossProcess**
 - **G4VMultipleScattering**
- A physical process can be simulated according to several models
 - each model being described by a model class
 - Naming scheme : « G4**ModelName****ProcessName**Model »
 - For example: **G4LivermoreComptonModel**
 - Models can be assigned to certain **energy ranges** and **G4Regions**
 - Inherit from **G4VEmModel** base class
- **Model classes provide the computation of**
 - Cross section and stopping power
 - Sample selection of atom in compound
 - Final state (kinematics, production of secondaries, ...)

- The scalability of Geant4 application in the MT mode depends on how effectively data management is performed
- **Shared EM physics data:**
 - tables for cross sections, stopping powers and ranges are kept by processes
 - Differential cross section data are kept by models
 - Material properties are in material data classes
 - EM parameters established for Physics Lists in the `G4EmParameters` class



Tables are filled by Master and have read-only access in run time

In this scheme number of threads is not limited

Main Gamma Processes

- Photo-effect is the main process for absorption of low-energy gamma
 - Rayleigh scattering should not be neglected if an accurate dosimetry simulation is needed
- At high energy gamma conversion dominates
- Gammas may be absorbed by nuclei due to giant dipole resonance
 - Producing neutrons, protons, and gamma






Photo-electric effect – example of gamma process

In the photo-electric absorption process a **photon is absorbed** by an atom and an **electron is emitted** with an energy:

$$E_{\text{photoelectron}} = E_{\gamma} - B_{\text{shell}}(Z_i) \quad (1)$$

The atom, left in an excited state with a vacancy in the ionized shell, decays to its ground state through a cascade of radiative and non-radiative transitions with the **emission of characteristic x-rays** and **Auger and Coster-Kronig electrons**.



Primary gamma may be polarized, photoelectron angular distribution will be affected

- Atomic de-excitation is initiated by other EM physics interactions:
 - photoelectric effect, ionisation (by e- or ions PIXE), Compton scattering,...
 - these interactions leave the target atom in an excited state
- The EADL (Evaluated Atomic Data Library) contains transition probabilities:
 - radiative transition characteristic X-ray emission (fluorescence photon emission)
 - Auger e- emission: initial and final vacancies are in different shells
 - Coster-Kronig e- emission: initial and final vacancies are in the same shells
- Due to a common interface, the atomic de-excitation is compatible with both the standard and the low-energy EM physics categories:
 - can be enabled and controlled by UI command (before initialization):

```
/process/em/fluor true  
/process/em/auger true  
/process/em/pixe true  
  
/run/initialize
```

- fluorescence transition is active by default in some EM physics constructor while others (Auger, PIXE) not

Gamma EM processes in the standard output

- Geant4 standard EM interactions for:
 - photon (γ) interactions (example):

```
phot: for gamma SubType= 12 BuildTable= 0
=====
LambdaPrime table from 200 keV to 100 TeV in 61 bins
===== EM models for the G4Region DefaultRegionForTheWorld =====
LivermorePhElectric : Emin=      0 eV  Emax=    100 TeV  AngularGenSauterGavrila  FluoActive
```


```
compt: for gamma SubType= 13 BuildTable= 1
=====
Lambda table from 100 eV to 1 MeV, 7 bins per decade, spline: 1
LambdaPrime table from 1 MeV to 100 TeV in 56 bins
===== EM models for the G4Region DefaultRegionForTheWorld =====
Klein-Nishina : Emin=      0 eV  Emax=    100 TeV
```

```
conv: for gamma SubType= 14 BuildTable= 1
=====
Lambda table from 1.022 MeV to 100 TeV, 18 bins per decade, spline: 1
===== EM models for the G4Region DefaultRegionForTheWorld =====
BetheHeitler : Emin=      0 eV  Emax=     80 GeV  AngularGenUrban
BetheHeitlerLPM : Emin=    80 GeV  Emax=    100 TeV  AngularGenUrban
```

```
Rayl: for gamma SubType= 11 BuildTable= 1
=====
Lambda table from 100 eV to 100 keV, 7 bins per decade, spline: 0
LambdaPrime table from 100 keV to 100 TeV in 63 bins
===== EM models for the G4Region DefaultRegionForTheWorld =====
LivermoreRayleigh : Emin=      0 eV  Emax=    100 TeV  CullenGenerator
```

Main Charged Particle Processes

- At low energies ionisation dominates for e-
 - For e+ annihilation dominates at very low energy
- Above critical energy bremsstrahlung is the main process
 - Radiation energy loss exceed ionization energy loss
 - Process of e+e- pair production has much less cross section
- Difference between electrons and positrons increased for low energy
 - Is practically negligible above critical energy



Simulation of a step of a charged particle

- Values of **mean dE/dx , range, cross section of δ -electron production, and bremsstrahlung** are **pre-computed at initialisation stage** of Geant4 and are stored in a **G4PhysicsTable**
- **At run time** for each simulation step, a spline interpolation of tables is used to get **mean energy loss**
- At each step, a sampling of the **energy loss fluctuation** is performed
 - The interface to a fluctuation model is **G4VEmFluctuationModel**
- The cross sections of δ -electron production and bremsstrahlung are used to **sample production above the threshold T_{cut}** at **PostStep**
- If atomic de-excitation is active, then **fluorescence and Auger electron production** is sampled **AlongStep** and **PostStep**

Hadron and ion ionisation



- Bethe-Bloch formula with corrections used for $E > 2$ MeV


$$-\frac{dE}{dx} = 4\pi N_e r_0^2 \frac{z^2}{\beta^2} \left(\ln \frac{2m_e c^2 \beta^2 \gamma^2}{I} - \frac{\beta^2}{2} \left(1 - \frac{T_c}{T_{\max}} \right) - \frac{C}{Z} + \frac{G - \delta - F}{2} + zL_1 + z^2L_2 \right)$$

- C – shell correction
- G – Mott correction
- δ – density correction
- F – finite size correction
- L_1 - Barkas correction
- L_2 - Bloch correction
- Nuclear stopping
- Ion effective charge



- Bragg peak parameterizations for $E < 2$ MeV
 - ICRU'49, ICRU'73, ICRU'90, and NIST databases

- Scaling relation for heavy particles:

- $S_h(E) = S(E * M_p / M_h) * Q_h^2$,
- M_h, Q_h - hadron mass and charge
- Applicable to any charged particle including exotics and all ions
- This is possible, because dE/dx depend mainly on β




- **Urban model** based on a simple model of particle-atom interaction
 - Atoms are assumed to have only two energy levels E_1 and E_2
 - Particle-atom interaction can be:
 - an excitation of the atom with energy loss $E = E_1 - E_2$
 - an ionization with energy loss distribution $g(E) \sim 1/E^2$
- **PAI model** uses photo absorption cross section data
 - Energy transfers are sampled with production of secondary e^- or γ
 - Relativistic model
 - Very slow model, should be applied for sensitive region of detector





Multiple Coulomb scattering

- **Coulomb scattering**: elastic scattering of charged particles on the atomic potential
- **Event-by-event modelling of elastic scattering** is feasible only if the mean number of interactions per track is below few hundred
- **this limits the applicability of the detailed simulation model only for electrons with relatively low kinetic energies**
 - up to 100 keV or thin targets



- detailed simulation becomes very inefficient, high energy particle transport simulation codes employ condensed history simulation model
 - **multiple scattering (MSC)** model is a solution
 - each track is simulated considering many elastic scattering at a step
- **A summary effects of high number of elastic interactions is in**
 - angular deflection of the particle
 - spatial displacement of the track post step point
 - increased effective track length


- The algorithm performs simulation of many elastic scatterings at a step of a particle
 - The physics processes and the geometry select the step length; MSC performs the $t \leftrightarrow z$ transformation only
 - Sampling of scattering angle (θ , Φ)
 - Computing of displacement and relocation of particle AlongStep
- To provide accurate simulation on geometry interface between different materials MSC step limitation is applied
 - Simple
 - UseSafety
 - UseSafetyPlus
 - UseDistanceToBoundary
- Other step limit parameters:
 - **RangeFactor** – is the most important
 - Geometry factor
 - Safety factor
 - Skin
 - Lambda limit
- Default MSC parameters are optimized for
 - Accurate simulation of EM showers
 - HEP sampling calorimeters
 - Accurate simulation of shielding




Secondary production thresholds

Secondary production threshold for bremsstrahlung

- **Bremsstrahlung photon emission:**
 - low energy photons (k) will be emitted with high rate DCS $\sim 1/k$
 - generation and tracking of all these low energy photons would not be feasible (CPU time)
 - but low energy photons has a very small absorption length
 - If the detector spacial resolution is worst than this length then the followings are *equivalent*:
 - *a*: generating and tracking these low energy photons till all their energy will be deposited
 - *b*: or just depositing the corresponding energy at the creation point (at a trajectory point)
 - note, that we think in energy scale at the model level that translates to length (spacial) at the transport level
 - a secondary production threshold might be introduced (either in energy or length)




22 27. Passage of particles through matter



Secondary production threshold technique

- Introduce secondary photon production threshold:
 - *secondary photons*, with initial energy below a gamma production threshold ($k < E_{\gamma}^{\text{cut}}$), are not generated
 - the corresponding energy (that would have been taken away from the primary) is accounted as *CONTINUOUS* energy loss of the primary particle along its trajectory
- Electron makes a step with a given length L , one can compute the mean energy loss (due to sub-threshold photon emissions) along the step as $L \times dE/dx$ (would be true only if $E = \text{const}$ along the step)
- *Secondary photons*, with initial energy above a gamma production threshold ($k > E_{\gamma}^{\text{cut}}$), are generated (*DISCRETE*)
- the emission rate is determined by the corresponding (restricted) cross section (σ)




$$\frac{dE}{dx}(E, E_{\gamma}^{\text{cut}}, Z) = \mathcal{N} \int_0^{E_{\gamma}^{\text{cut}}} k \frac{d\sigma}{dk}(E, Z) dk$$

$$\sigma(E, E_{\gamma}^{\text{cut}}, Z) = \int_{E_{\gamma}^{\text{cut}}}^E \frac{d\sigma}{dk}(E, Z) dk$$


- Secondary production thresholds in Geant4:
 - user needs to provide them in length (the default value of 0.7 [mm] for the reference physics lists)
 - its proper value application dependent (size of the sensitive volume, CPU)
 - UI command: `/run/setCut 0.1 mm`
 - `/run/setCutForAGivenParticle e- 0.1 mm`
 - internally translated to energies at initialisation (depending on material and particle type)
 - the corresponding energy has a minimum value: default 1 keV but the user can set it
 - UI command: `/cuts/setLowEdge 500 eV`
 - production threshold defined for **gamma**, **e⁻**, **e⁺** and **proton** secondary particle types
 - **gamma** production threshold is used in bremsstrahlung while the **e⁻** in ionization
 - **e⁺** production threshold might be used in case of e-/e+ pair production
 - **proton** production threshold is used as a kinetic energy threshold for nuclear recoil in case of elastic scattering of *all hadrons and ions*
 - **gamma and e⁻** production thresholds might be used (optionally: `/process/em/applyCuts true`) in all discrete EM interactions producing such secondaries - Compton, Photoelectric, etc.
 - it's not mandatory to use production thresholds
 - however, high energy physics simulation would not be feasible without them !

Example demonstrating importance of cuts



cut [μm]	mean E_{dep}	rms E_{dep}	prod. thres. [keV]		mean num. sec.	
			γ	e^-	γ	e^-
1	1.54423	0.000573911	0.99	0.99	0.0006811	0.1018230
2	1.54443	0.000583879	0.99	2.9547	0.0006843	0.0316897
5	1.54882	0.000605834	0.99	13.1884	0.0006857	0.0068261
10	1.56717	0.000665733	0.99	31.9516	0.0006730	0.0028232
20	1.58734	0.000743473	1.08038	47.8191	0.0006651	0.0018811
50	1.62223	0.000912408	1.67216	80.7687	0.0006557	0.0011304
100	1.65893	0.001108240	2.32425	121.694	0.0006518	0.0007536
200	1.69338	0.001342180	3.2198	187.091	0.0006465	0.000477
500	1.74642	0.001774670	5.00023	337.972	0.0006184	0.0002617
1000	1.78751	0.002219870	6.95018	548.291	0.0006054	0.0001622
2000	1.83440	0.002861020	9.66055	926.09	0.0005786	9.3e-05
5000	1.90700	0.004243030	14.9521	2074.3	0.0005427	4.07e-05
10000	1.97378	0.006036600	20.6438	4007.59	0.000521	2.22e-05

Compute the mean of the energy deposit in the target: E_0 - primary, E_f - final energy



Golden rule:

For transport in solid/liquid media cut in range should be below minimal geometry size

EM PHYSICS CONSTRUCTORS

- A Physics list is the mandatory user class making the general interface between the physics the user needs and the Geant4 kernel
- List of particles: for which EM physics processes are defined
 - $\gamma, e^{\pm}, \mu^{\pm}, \pi^{\pm}, K^{\pm}, p, \Sigma^{\pm}, \Xi^{-}, \Omega^{-}, \text{anti}(\Sigma^{\pm}, \Xi^{-}, \Omega^{-})$
 - $\tau^{\pm}, B^{\pm}, D^{\pm}, D_s^{\pm}, \Lambda_c^{+}, \Sigma_c^{+}, \Sigma_c^{++}, \Xi_c^{+}, \text{anti}(\Lambda_c^{+}, \Sigma_c^{+}, \Sigma_c^{++}, \Xi_c^{+})$
 - d, t, He3, He4, Genericlon, anti(d, t, He3, He4)
 - 12 light hyper- and anti-hyper- nuclei
- The **G4ProcessManager** of each particle maintains a **list of processes**
- Geant4 provides several configurations of EM physics lists called **constructors** (**G4VPhysicsConstructor**) in the **physics_lists** library of Geant4
- These constructors can be included into a **modular Physics list** in a user application (**G4VModularPhysicsList**)

Geant4 standard EM Physics Constructors for HEP applications

- Description of Coulomb scattering:
 - e^{\pm} : Urban - MSC model below 100 [MeV] and the Wentzel - WVI + Single scattering (mixed simulation) model above 100 [MeV]
 - muon and hadrons: Wentzel - WVI + Single scattering (mixed simulation) model
 - ions: Urban - MSC model
- Different MSC stepping algorithms and/or parameters: speed v.s. accuracy

Constructor	Components	Comments
<code>G4EmStandardPhysics</code>	Default: nothing or <code>_EM0</code> (QGSP_BERT, FTFP_BERT,...)	for ATLAS and other HEP simulation applications
<code>G4EmStandardPhysics_option1</code>	Fast: due to simpler MSC step limitation, cuts used by photon processes (FTFP_BERT_EMV)	similar to one used by CMS; good for crystals but not good for sampling calorimeters (with more detailed geometry)
<code>G4EmStandardPhysics_option2</code>	Experimental: similar to option1 with updated photoelectric model but no-displacement in MSC (FTFP_BERT_EMX)	similar to one used by LHCb

Combined Geant4 EM Physics Constructors

- The primary goal is more the physics accuracy over the speed
- Combination of standard and low-energy EM models for more accurate physics description
- More accurate models for e^{\pm} MSC (Goudsmit-Saunderson(GS)) and more accurate stepping algorithms (compared to HEP)
- Stronger continuous step limitation due to ionisation (as others given per particle groups)
- Recommended for more accuracy sensitive applications: medical (hadron/ion therapy), space

Constructor	Components	Comments
<code>G4EmStandardPhysics_option3</code>	Urban MSC model for all particles	proton/ion therapy
<code>G4EmStandardPhysics_option4</code>	most accurate combination of models (particle type and energy); GS MSC model with Mott correction and error-free stepping for e^{\pm})	the goal is to have the most accurate EM physics description
<code>G4EmLivermorePhysics</code>	Livermore models for e^{-} , γ below 1 GeV and standard above; same GS MSC for e^{\pm} as in option4)	accurate Livermore based low energy e^{-} and γ transport
<code>G4EmPenelopePhysics</code>	PENELOPE models for e^{\pm} , γ below 1 GeV and standard above; same GS MSC for e^{\pm} as in option4)	accurate PENELOPE based low energy e^{-} , e^{+} and γ transport

Experimental Geant4 EM Physics Constructors

- Supposed to be used only by the developers for validations and model developments
- The main difference is in the description of the Coulomb scattering (GS, WVI, SS)

Constructor	Components	Comments
<code>G4EmStandardPhysicsGS</code>	standard EM physics and the GS MSC model for e^{\pm} with HEP settings	may be considered as an alternative to EM0 i.e. for HEP
<code>G4EmStandardPhysicsWVI</code>	WentzelWVI + Single Scattering mixed simulation model for Coulomb scattering	high and intermediate energy applications
<code>G4EmStandardPhysicsSS</code>	single scattering (SS) model description of the Coulomb scattering	validation and verification of the MSC and mixed simulation models
<code>G4EmLowEPPhysics</code>	Monarsh University Compton scattering model, 5D gamma conversion model, WVI-LE model	testing some low energy models
<code>G4EmLivermorePolarized</code>	polarized gamma models	a (polarized) extension of the Livermore physics models

USER INTERFACE TO EM PHYSICS

- EM parameters of any EM physics list may be modified at initialization of Geant4 using C++ interface to the `G4EmParameter` class or via UI commands
- Example of interfaces of `G4EmParameters`:
 - `SetMuHadLateralDisplacement()`
 - `SetMscMuHadRangeFactor()`
 - `SetMscMuHadStepLimitType()`
- Corresponding UI commands:
 - `/process/msc/MuHadLateralDisplacement`
 - `/process/msc/RangeFactorMuHad`
 - `/process/msc/StepLimitMuHad`
- Some other UI commands:
 - `/process/em/deexcitationIgnoreCut true`
 - `/process/eLoss/UseAngularGenerator true`
 - `/process/em/lowestElectronEnergy 50 eV`
 - `/process/em/lowestMuHadEnergy 100 keV`
 -

- Geant4 UI commands to define cuts and other EM parameters
- **G4EmCalculator**
 - easy access to cross sections and stopping powers (TestEm0)
- **G4EmParameters**
 - C++ interface to EM options alternative to UI commands
- **G4EmSaturation**
 - Birks effect (recombination effects)
- **G4ElectronIonPair**
 - sampling of ionisation clusters in gaseous or silicon detectors
- **G4EmConfigurator**
 - add models per energy range and geometry region
- **G4NIELCalculator**
 - Helper class allowing computation of NIEL at a step, which should be added in user stepping actions or sensitive detector (TestEm1)

How to extract Physics ?

- Possible to retrieve Physics quantities using a **G4EmCalculator** object
- Physics List should be **initialized**
- Example for retrieving the total cross section of a process with name **procName**, for **particle** and material **matName**

```
#include "G4EmCalculator.hh"
```

```
...
```

```
G4EmCalculator emCalculator;
```

```
G4Material* material =
```

```
  G4NistManager::Instance()->FindOrBuildMaterial(matName);
```

```
G4double density = material->GetDensity();
```

```
G4double massSigma = emCalculator.ComputeCrossSectionPerVolume  
  (energy, particle, procName, material)/density;
```

```
G4cout << G4BestUnit(massSigma, "Surface/Mass") << G4endl;
```

- A good example: [\\$G4INSTALL/examples/extended/electromagnetic/TestEm0](#)
Look in particular at the [RunAction.cc](#) class

SPECIAL EM TOPICS



PHYSICAL STAGE
step-by-step modelling of
physical interactions of incoming
& secondary ionising radiation
with biological medium
(liquid water)

- Excited **water molecules**
- Ionized **water molecules**
- **Solvated electrons**

Physico-chemical/chemical stage

- **Radical** species production
- Diffusion
- Mutual chemical interactions


Geometrical models
DNA strands, chromatin fibres, chromosomes, whole cell nucleus, cells...
for the prediction of damage resulting from direct and indirect hits



Special EM topics: EM models per region

- Special EM models can be set to be used only in a given detector `G4Region`
- Example to use Geant4-DNA physics in a given detector region on the top of the standard EM physics:
 - the **G4EmConfigurator** can be used to add Geant4-DNA models
 - the DNA models are used only in the region B. for energies below 10 MeV
 - makes possible CPU and physics performance optimisation
 - the more accurate CPU intense simulation is done only in the region of interest
 - UI commands are available from Geant4 10.2 that allow easy configuration of some models per-region on the top of any EM constructor:


```
/process/em/AddPAIRegion proton MYREGION pai  
/process/em/AddMicroElecRegion MYREGION  
/process/em/AddDNARegion MYREGION opt0
```



Special EM topics: EM models per region



- Special EM models can be set to be used only in a given detector `G4Region`
- Example to use Geant4-DNA physics in a given detector region on the top of the standard EM physics:

the **microdosimetry** extended example:



Quantum entanglement in positron annihilation

(arXiv: 2012.04939v1)



- There is angular correlation for Compton scattering of two photons in PET device
- Geant4 method how simulate quantum effects has been developed by J. Allison
- The developed method may be potentially used in HEP

THANK YOU