

# Neutrino Radiation for a realistic Collider

C. Carli, C. Ahdida, D. Calzolari,  
G. Lerner, G. Lacerda, A. Lechner,  
K. Skoufaris ...

Muon Collider annual meeting, 13<sup>th</sup> October 2022





# Content

- Introduction - Neutrino Radiation Issue
- Analytical estimates
  - ◆ Absorbed dose per decay
  - ◆ approximation by Gaussian
- FLUKA Simulations of doses due to neutrino interactions
- Extrapolation to collider lattice
- Mitigation by “Wobbling”
- Summary and Outlook

# Introduction

## Neutrino Radiation Issue


- Radiation due to neutrino beam reaching the earth surface
  - ◆ Narrow radiation “cone” for a short piece of the machine
  - ◆ Showers from neutrinos interacting close to earth surface generate dose seen at surface
  - ◆ Matter in front (“shielding”) does not help but makes situation even worse
  
- Strong increase of maximum dose with muon energy
  - ◆ Cross sections about proportional to energy
  - ◆ Typical energy per interaction of neutrino with matter proportional to muon energy
  - ◆ Opening of radiation cone inversely proportional to muon energy



# Analytical estimates absorbed dose per decay

Following approach by B.J. King,  
(see, e.g., arXiv:physics/9908017)

- Consider say decay  $m^- \rightarrow e^- + \bar{n}_e + n_m$
- In muon rest system: anisotropic neutrino distribution with known energy distribution
- Lorentz boost to obtain neutrino direction and energies in lab system (Lorentz factor  $g$ )
- Sum over
  - Both neutrino types
  - cross sections of possible interactions times fraction of energy deposited per interaction in shower
- Gives



$$\Delta D = \underbrace{\left( 0.410 \cdot 10^{-39} \frac{\text{m}^2}{\text{TeV}} \frac{E_{\mu}^2}{4} \right)}_{1.10 \cdot 10^{-28} \text{ Gy m}^2} \frac{1}{m_u} \frac{1}{L_s^2} \frac{\langle E_v^2 \rangle}{\langle E_v^{i2} \rangle} \frac{1}{N} \frac{dN}{d\Omega} = \left( 1.10 \cdot 10^{-28} \text{ Gy m}^2 \right) \frac{4\gamma^4}{\pi L_s^2} \frac{1}{\left( 1 + \gamma^2 (d/L_s)^2 \right)^4}$$

with  $\frac{\langle E_n^2 \rangle}{\langle E_n^{i2} \rangle} = 4g^2 / \left( 1 + g^2 J^2 \right)^2$  the ratio between the square of the neutrino energy in the lab and the rest system  
 and  $N^{-1} dN/dW = (g^2/\rho) / \left( 1 + g^2 J^2 \right)^2$  the angular distribution of neutrinos

- The decay  $m^+ \rightarrow e^+ + n_e + \bar{n}_m$  gives (almost) identical final result despite different cross sections and fractions

# Analytical estimates approximation by Gaussian

- Absorbed dose reasonably well described by Gaussian



$$DD \gg \left(1.10 \times 10^{-28} \text{ Gy m}^2\right) \frac{4g^4}{\rho L_s^2} \frac{1}{\left(1 + g^2 \left(d/L_s\right)^2\right)^4}$$

$$\approx \left(1.10 \cdot 10^{-28} \text{ Gy m}^2\right) \frac{4g^4}{\rho L_s^2} e^{-3g^2(d/L_s)^2}$$

- Rms opening angle of neutrino radiation cone of

$$J_{rms} = 1 / (\sqrt{6}g)$$

- Assuming Gaussian for the beam divergence
  - Folding of divergence from muon decay process and beam divergence simple
- Almost suitable as source term to estimate doses generated by neutrino interactions



Dose per muon decay at distance  $L_s = 100$  km and  $\gamma = 47\,322$ , i.e., 10 TeV com energy

# FLUKA simulations of doses due to neutrino interactions

- Motivation: improvement and check of analytical derivations
  - ◆ Possible widening of neutrino radiation cone due to lateral extension of shower
  - ◆ Effective dose instead of absorbed dose

## ▪ Ansatz

$$DH \approx D\hat{H} e^{-d^2/2S_{DH}^2}$$

$$= w_{R,eff} \cdot \left(1.104 \cdot 10^{-28} \text{ Gy m}^2\right) \frac{4g^4}{\rho(L_s^2 + 6g^2S_s^2)} \exp\left(-\frac{d^2}{2(L_s^2/6g^2 + S_s^2)}\right)$$


$$D\hat{H} \approx w_{R,eff} \cdot \left(1.104 \cdot 10^{-28}\right) 4g^4 / \left(\rho(L_s^2 + 6g^2S_s^2)\right)$$

$$S_{DH} \approx \sqrt{L_s^2 / (6g^2) + S_s^2}$$

$$D\hat{H} S_{DH}^2 \approx w_{R,eff} \cdot \left(1.104 \cdot 10^{-28}\right) 2g^2 / (3\rho)$$

- ◆ with  $w_{R,eff}$  to convert absorbed dose  $D$  to dose equivalent  $H = w_{R,eff} D$
- ◆ and  $S_s$  the rms radial extension of the shower (assumed as well to be Gaussian?!)
- Result of fitting “by hand” to FLUKA results for 5 TeV muons (green lines in plot)
  - ◆  $w_{R,eff} = 1.3 \text{ Sv/Gy}$  and  $S_s = 0.12 \text{ m}$
  - ◆ Results may be different for other energies
  - ◆ Shower extension  $S_s$  neglected for further studies (slightly pessimistic)

FLUKA studies by G. Lerner et al.




# Extrapolation to a collider lattice

- Taking details of lattice into account:

- ◆ Twiss gamma functions  $g_H(s)$  and  $g_V(s)$  and derivative of dispersion  $D'(s)$
- ◆ Physical rms emittances  
 $e_H = e_V = 25 \text{ mm} / g \gg 0.528 \text{ nm}$
- ◆ Rel. momentum spread  $s_p / p = 10^{-3}$

- Extrapolation of argument of exponential



$$-3g^2 (J_V^2 + J_H^2) = \frac{-J_V^2}{2 \times \frac{1}{6g^2}} + \frac{-J_H^2}{2 \times \frac{1}{6g^2}} \quad \text{D} \quad -\frac{J_V^2}{2S_{J_V}^2} - \frac{(J_H - \hat{J}(s))^2}{2S_{J_H}^2}$$

$$S_{J_H}^2 = \frac{1}{6g^2} + e_H g_H(s) + \left( \frac{s_p}{p} \cdot D'(s) \right)^2$$

$$S_{J_V}^2 = \frac{1}{6g^2} + e_V g_V(s)$$

- ◆ Without mitigation measures gives

$$\frac{dH}{dt} = \left( 1.104 \cdot 10^{-28} \text{ Gy m}^2 \right) w_{R,eff} \frac{4g^4 f_r N_m}{\rho L_s^2 C} \int ds \frac{1/(6g^2)}{S_{J_H} \cdot S_{J_V}} \exp \left[ -\frac{(J_H - \hat{J}(s))^2}{2S_{J_H}^2} - \frac{J_V^2}{2S_{J_V}^2} \right]$$

with  $N_m$  the number of muons per bunch,  $f_r$  the repetition rate and  $C$  the circumference



# Numerical evaluations

## Simple cases

- Divergence of muon beam neglected, peak dose rate

- ◆ Straight section  $D_S$ , using  $C = 2\rho g E_{rm} / (c e \bar{B})$  and  $L_s \gg 2R_e h$  with  $R_e \gg 6.38 \times 10^6$  m the earth radius

$$\frac{dH_S}{dt} = \left( 6.85 \cdot 10^{-22} \frac{\text{Sv}}{\text{T}} \right) f_r N_m \left( \frac{E}{5\text{TeV}} \right)^3 \frac{D_S \bar{B}}{h} \quad \text{for beam energies around } E \gg 5\text{TeV}$$

- ◆ Bending magnet – integration w.r.t  $s$  using  $J_H - \hat{J}(s) = (eB / (gE_{rm} / c)) s$

$$\frac{dH_B}{dt} = \left( 6.85 \cdot 10^{-22} \frac{\text{Sv}}{\text{T}} \right) f_r N_m \left( \frac{E}{5\text{TeV}} \right)^3 \frac{\bar{B}}{h} \int ds \exp \left( -3 \left( \frac{eBc}{E_{rm}} s \right)^2 \right) = \left( 2.47 \cdot 10^{-22} \text{Sv m} \right) f_r N_m \left( \frac{E}{5\text{TeV}} \right)^3 \frac{\bar{B}}{h B}$$

- ◆ Integrated peak equivalent dose per muon beam for one year operation (5000 h =  $18 \cdot 10^6$  s) without mitigation measures

$N_m = 1.8 \times 10^{12}$  muons per bunch,  
 $f_r = 5\text{Hz}$  repetition,  
 $E \gg 5\text{TeV}$  beam energy,  
 $C = 10000\text{km}$  circumference and  
 $\bar{B} = 10.42\text{T}$  average field

h (m)	$L_s$ (km)	$H_S$ (mSv) for $\Delta s = 0.3$ m	$H_B$ (mSv) for $B = 8$ T
100	35.7	3.5	0.52
200	50.5	1.75	0.26
500	79.7	0.70	0.105
784	100	0.45	0.067


Contribution only from of  
the two muon beams

Mitigation mandatory!!




# Numerical evaluations equivalent dose from arc cell at 100 km

- Integrals evaluated for present (work in progress) 10 TeV com collider arc half cell
  - ◆ In collider mid-plane as function of  $\mathcal{J}_H$  (i.e.,  $\mathcal{J}_V = 0$ ) for one year (5000 h operation)



# Mitigation by “Wobbling”

- Wobbling of machine in vertical direction – part of MAP proposal?
  - ◆ Time-dependent mechanical deformation of ring around arc (including chromatic compensation, matching section and FMC arc cells)
  - ◆ High precision movement system
  - ◆ Impact on optics?
- For 10 TeV com collider with 10 km circumference and say 4.8 km arcs



Vertical bend  
 $\pm 16.7 \text{ Tm}$

- ◆ Combination of pieces of parabola – two pieces with opposite curvature one period
  - ◆ Say 8 periods 660 m long periods generating angles between  $-1 \text{ mrad}$  and  $+1 \text{ mrad}$
  - ◆ Magnetic field (average) bending in vertical  $\pm 0.11 \text{ T}$
  - ◆ Excursion (maximum total)  $\pm 150 \text{ mm}$
  - ◆ Replaces vertical Gaussian angle distribution with rms opening of  $\approx 0.0086 \text{ mrad}$  by about rectangular distribution within  $\pm 1 \text{ mrad}$
- => About two order of magnitude reduction of peak dose rates

# Summary and Outlook

- Dose generated at the earth surface due to showers generated by neutrino interactions is a serious issue
  - ◆ Muon decays generate narrow cone swept by bendings in bending plane
  - ◆ In particular high doses in direction of (short) straights sections
  - ◆ Numerical estimates based on “source term” from FLUKA simulations  
(=> Comparison with MARS results to be done?)
- Time-dependent vertical deformation of the whole arc proposed as mitigation measure
  - ◆ For a 10 TeV com collider angle variations (linear up and down) in a range  $\pm 1$  mrad would allow to gain about two orders of magnitude
  - ◆ To be combination with other mitigation measures
    - Collider installed deep underground with suitable positioning
    - Appropriate orientation towards uncritical areas (possibly owned by facility)
- Feasibility to be studied
  - ◆ High precision movement system
  - ◆ Impact on beam dynamics in particular in chromatic compensation and matching sections (vertical dispersion, orbit stability ...)