




# **3D structure of jet-induced diffusion wake in an expanding quark-gluon plasma**

Zhong Yang


Tan Luo, Wei Chen, Longgang Pang and Xin-nian Wang

ATHIC 2023 Hiroshima

# Jet in heavy-ion collisions




**QGP(quark-gluon plasma):** A deconfined strongly interacting matter that behaves like a perfect fluid



# Jet-induced medium response

Jet-induced medium response in the form of Mach-cone-like excitation.



- Width of front wake of Mach cone is related with viscous properties of QGP medium;
- Mach cone angle is sensitive to EoS.


$$\sin\theta = \frac{c_s}{v}$$

R.B.Neufeld. PRC79,054909(09')

# Medium modifications of gamma-jets at LHC

## Jet Profile


$$\rho(r) = \frac{1}{\delta r} \frac{\sum_{\text{jets}} \sum_{r < r_{\text{trk}} < r + \delta r} (p_T^{\text{trk}} / p_T^{\text{jet}})}{\sum_{\text{jets}} \sum_{r_{\text{trk}} < R} (p_T^{\text{trk}} / p_T^{\text{jet}})}$$



Luo, Cao, He & Wang, arXiv:1803.06785

**Jet-induced medium response can contribute to enhancement of soft hadrons within the jet cone**

## Jet fragmentation Function



Chen, Cao, Luo, Pang & Wang, arXiv: 2005.09678

# Medium response and soft gluon radiation


**Medium response** leads to enhancement of soft hadrons in the direction of jet. (Jet shape,  $I_{\{AA\}}...$ )

**Medium-induced gluon radiation** has the similar effect.

Medium response:  $\delta f(p) \sim e^{-p \cdot u/T}$

Medium-induced gluon radiation:  $\omega \approx \lambda^2 \hat{q}/2 \sim T$

It is difficult to separate their contribution to enhancement of soft hadrons.



**Diffusion wake:** an unambiguous part of the jet-induced medium response. It can lead to depletion of soft hadrons in the opposite direction of the jet.

# LBT: Linear Boltzmann Transport


$$p_1 \partial f_1 = - \int dp_2 dp_3 dp_4 (f_1 f_2 - f_3 f_4) |M_{12 \rightarrow 34}|^2 (2\pi)^4 \delta^4 \left( \sum_i p^i \right) + \textit{inelastic}$$

Medium-induced gluon(HT):

$$\frac{dN_g}{dz d^2 k_{\perp} dt} \approx \frac{2C_A \alpha_s}{\pi k_{\perp}^4} P(z) \hat{q} (\hat{p} \cdot u) \sin^2 \frac{k_{\perp}^2 (t - t_0)}{4z(1-z)E}$$

Tracked partons:


- Jet shower partons
- Thermal recoil partons
- Radiated gluons
- Negative partons(Back reaction induced by energy-momentum conservation)




# LBT: Jet-induced medium response

Shock wave: propagation of recoil particles

(a)  $t=4$  fm/c



(b)  $t=8$  fm/c



He, Luo, Wang & Zhu, PRC91 (2015) 054908

Diffusion wake: propagation of negative partons

# CoLBT-hydro model

1. LBT for energetic partons(jet shower and recoil)
2. Hydrodynamic model for bulk and soft hadrons: CLVisc
3. Sorting jet partons according to a cut-off parameter  $p_{cut}^0$

hard partons:  $p \partial f(p) = -C(p) \quad (p \cdot u > p_{cut}^0)$

soft and negative partons:

$$j^\nu = \sum_i p_i^\nu \delta^{(4)}(x - x_i) \theta(p_{cut}^0 - p \cdot u)$$


4. Updating medium information by solving the hydrodynamic equation with source term

$$\partial_\mu T^{\mu\nu}(x) = j^\nu(x)$$

5. The final hadron spectra:
  - (1) hadronization of hard partons within a parton recombination model
  - (2) jet-induced hydro response via Cooper-Frye freeze-out



# CoLBT-hydro: Jet-induced medium response




Chen, Yang, He, Ke, Pang & Wang, PRL 127 (2021) 8, 082301


**The Mach-cone-like jet-induced medium response including the diffusion wake is clearly seen in the right panel.**

# Gamma-jet substructure within CoLBT-hydro

## Jet asymmetry




## Jet profile



Yang, Luo, Chen, Pang, Wang, PRL 130 (2023) 5,052301

CoLBT-hydro model can describe both jet energy loss and its redistribution in QGP

# Azimuthal distribution of soft hadrons at RHIC





It is the signal of diffusion wake which leads to the depletion of soft hadrons in the  $\gamma$  direction

Chen, Cao, Luo, Pang & Wang, PLB777(2018)86

# Azimuthal distribution of soft hadrons at LHC

Mixed event MPI (Initial Multiple parton interaction) subtraction:



$$\frac{dN_{MPI}^{hZ}}{d\phi} \approx \frac{dN_{mix}^{hZ}}{d\phi} - \int_1^\pi \frac{d\phi'}{\pi} \left( \frac{dN^{hZ}}{d\phi} - \frac{dN^{hZ}}{d\phi} \Big|_{\phi=1} \right)$$





Chen, Yang, He, Ke, Pang & Wang, PRL 127 (2021) 8, 082301

# Motivation to study 3D structure of DW

- (1) The previous studies of diffusion wake focus on the azimuthal angle.**
- (2) The jet is a 3D observable, thus the diffusion wake should also have a 3D structure.**



# 3D structure of diffusion wake




$$\Delta\eta = \eta_h - \eta_{\text{jet}}$$

$$\Delta\phi = \phi_h - \phi_{\text{jet}}$$


Yang, Luo, Chen, Pang, Wang, Phys.Rev.Lett., 2023,130(5):052301

**Diffusion wake valley(DF-wake valley):**a valley is formed on top of the MPI ridge due to the depletion of soft hadrons by jet-induced diffusion wake.

# 3D structure of diffusion wake



# 3D structure of diffusion wake



Yang, Luo, Chen, Pang, Wang, Phys.Rev.Lett., 2023,130(5):052301


**Double Gaussian fitting:** 
$$F(\Delta\eta) = \int_{\eta_{j1}}^{\eta_{j2}} d\eta_j F_3(\eta_j) (F_2(\Delta\eta, \eta_j) + F_1(\Delta\eta))$$

$$F_1(\Delta\eta) = A_1 e^{(-\Delta\eta^2/\sigma_1^2)}$$

$$F_2(\Delta\eta, \eta_j) = A_2 e^{-(\Delta\eta + \eta_j)^2/\sigma_2^2}$$



# Sensitivity to Jet energy loss




Yang, Luo, Chen, Pang, Wang, Phys.Rev.Lett., 2023,130(5):052301

Longer propagation length and larger jet energy loss leads to deeper DF-W valley.

The MPI ridge has a very weak and non-monotonic dependence on  $x_{jy}$  due to the non-monotonic dependence of the propagation length on  $x_{jy}$  for minijets from MPI.


# Sensitivity to shear viscosity



Yang, Luo, Chen, Pang, Wang, *Phys.Rev.Lett.*, 2023,130(5):052301


Competition between increased radial flow and negative longitudinal pressure in the shear correction of the energy momentum tensor leads to a a slightly smaller MPI ridge and a deeper DF-wake valley in viscous hydro than in an ideal hydro.

# Sensitivity to equation of state




Yang, Luo, Chen, Pang, Wang, Phys.Rev.Lett., 2023,130(5):052301

The effective speed of sound is higher in eosq than s95.



# The signal in gamma-hadron correlation




$$\Delta\eta = \eta_h - \eta_\gamma$$

$\gamma$  as reference

Using gamma as reference can also get double peak structure in rapidity distribution.

Competition  $\left\{ \begin{array}{l} \rightarrow \text{Smearing effect caused by the difference between jet and gamma} \\ \rightarrow \text{Jets with larger energy loss are taken into account (No cut of jet pt in gamma-hadron event)} \end{array} \right.$

# The signal in Z-hadron correlation



$$p_T^Z > 30 \text{ GeV}/c$$



$$p_T^{\text{jet}} > 15 \text{ GeV}/c$$

$$|\eta_z| < 2.4$$

$$|\eta_{\text{jet}}| < 1.6$$

We can get the signal of the diffusion wake in **jet-hadron correlation**, but not in **Z-hadron correlation**.

# The smearing effect in Z-jet and gamma-jet



The differences of  $\eta$  and  $\phi$  between trigger and jet in Z-jet events are larger than that in  $\gamma$ -jet events.

**Using  $\gamma$ -hadron correlation to find signal of diffusion wake is a good choice for experimental group.**

# Summary



- 1. Jet-induced medium response can help us glean QGP properties.**
- 2. With MPI subtraction, we can get signal of diffusion wake at LHC.**
- 3. There is a unique signal of DF-wake in rapidity distribution of jet-hadron correlation.**
- 4. By double Gaussian fit method, we studied DF-wake valley's sensitivity to jet energy loss, shear viscosity and EoS.**
- 5. Using gamma-hadron correlation is a good choice to look for the signal of diffusion wake.**

**Thanks for your attention**






# Back up

# 3D structure of diffusion wake after ML selection




**Jet initial positions are selected by the ML associated 2D jet tomography**

# Energy density and quiver plot



# Energy density and quiver plot



# Medium response and soft gluon radiation

Medium response:  $\delta f(p) \sim e^{-p \cdot u/T}$


Medium-induced gluon radiation:

Formation time:  $\tau_f = \frac{2\omega}{k_T^2} \quad k_T^2 \approx \tau_f \hat{q} \quad \tau_f \approx \sqrt{2\omega/\hat{q}}$


Mean-free-path limits the formation time:  $\tau_f \leq \lambda \sim 1/T \quad \hat{q} \sim T^3$   
 $\omega \approx \lambda^2 \hat{q}/2 \sim T$

**It is difficult to separate contribution to enhancement of soft hadrons from medium-induced soft gluon radiation or medium response.**

# Equation of state



$$c_s^2 = \frac{\partial p}{\partial \epsilon}$$



# MPI Subtraction

- (1) We first calculate the uniform correlation between  $Z/\gamma$  in one event and hadrons from another similar  $Z/\gamma$ -jet event.
- (2) We assume the effect of the diffusion wake on the total  $Z/\gamma$ -hadron yield in the mixed events is negligible.
- (3) Contributions from jets to the  $Z/\gamma$ -hadron correlation in these mixed events, which are assumed to be the same as the integrated  $Z/\gamma$ -hadron yield within an angle  $|\Delta\phi| > 1$  in  $Z/\gamma$ -jet events in addition to the MPI background.

