

CKM global fits and new physics in meson mixing



VNIVERSITAT
DE VALÈNCIA



“This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101031558”

Luiz VALE SILVA (IFIC, UV – CSIC)


**[Charles, Descotes-Genon, Ligeti, Monteil, Papucci, Trabelsi, LVS,
PRD 102, 056023 (2020), arXiv:2006.04824]**

6th FCC Physics Workshop, Krakow, 25/01/2023

The Standard Model (SM) and Beyond

- **Flavour physics** played a central role in the **formulation of the SM** (new fermion generations, manifestation of CP Violation, etc.)
- Many **flavour observables** enjoy the status of **precision physics**, thanks to progress in different fronts (e.g. QCD inputs)
- **Flavour physics** can play a **leading role** in addressing the questions left open by the SM, and reveal New Physics sectors

Hierarchies in the spectrum of particles and CKM matrix:




	d	s	b
u	■	■	·
c	■	■	■
t	·	■	■

Reach to New Physics (NP)

- **Low-energy observables** → probe energies much beyond the reach of direct searches
- The bounds on non-SM contributions **shape NP candidates**
- If deviation seen, possible **NP manifestation!**


Different sectors:



[European Strategy Update 2020, arXiv:1910.11775]

Current status of flavour

- **Overall agreement w/ the SM, but some existing tensions (e.g., incl. vs excl. $|V_{xb}|$)**




[CKMfitter update as of Spring '21]

- **Flavour** is one of the **main physics cases for future experiments**
- **Future data will guide the field**, testing present anomalies and possibly revealing new ones

NP in B meson mixing

- **HERE:** address present and future bounds on NP in $|\Delta B|=2$, and discuss future limitations

- $|\Delta B|=2$: NP competes with **suppressions in the SM (GIM/loop)**, and enjoy the status of **precision physics**



- **Not discussing D meson mixing**

[K meson mixing: PRD 89, 033016 (2014), [arxiv:1309.2293](#)]

- **Combine projections for future data:** need global fit including “tree” and “loop” observables

[see e.g. [CKMfitter](#)]

NP in B meson mixing

- **NP in $|\Delta B|=2$:**

h_d and h_s set sizes

Bag parameters,
↓ decay constants

$$M_{12} = M_{12}^{\text{SM}} \times (1 + h e^{2i\sigma})$$

NP parameters

- **Assumptions:**

- No NP in $|\Delta F|=1$:

tree level in SM (γ , $|V_{ub}|$, $|V_{cb}|$, ...) free of NP

- NP is short-distance

- Unitarity of the CKM 3x3 matrix

- Unrelated NP in B_d and B_s systems


[See: PRD 89, 033016 (2014),
[arxiv:1309.2293](https://arxiv.org/abs/1309.2293)]

- SMEFT: **four-quark operators** of different chiral structures

Present status of NP in B meson mixing

- **Agreement with the SM ($h_d=h_s=0$) at $\sim 1\sigma$**
- **Allowed size for NP at the level of $O(20\%)!$**
- **Extractions of ρ and η** (Wolfenstein parm.) degrade by factor ~ 3

Status as of Summer '19



Black dot: best fit point;
 σ_d and σ_s are unconstrained


New era of flavour ahead

	LHC era			HL-LHC era	
	Run 1 (2010-12)	Run 2 (2015-18)	Run 3 (2021-24)	Run 4 (2027-30)	Run 5+ (2031+)
ATLAS, CMS	25 fb ⁻¹	150 fb ⁻¹	300 fb ⁻¹	→	3000 fb ⁻¹
LHCb	3 fb ⁻¹	9 fb ⁻¹	23 fb ⁻¹	50 fb ⁻¹	*300 fb ⁻¹

* assumes a future LHCb upgrade to raise the instantaneous luminosity to $2 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

[↑ See [arXiv:1808.08865](https://arxiv.org/abs/1808.08865)]

- Expression of interest for an **LHCb Upgrade II**
- **Belle II**: 50x the Belle and nearly 100x the BaBar data sets; ongoing discussions about **upgrade**



Benchmarks for the future

- **Phase I: LHCb-upgrade I** 50/fb, & **Belle II** 50/ab
- **Phase II: LHCb-upgrade II** 300/fb, & **Belle II upgrade** 250/ab
- **Phase III: Phase II + FCC-ee**

FCC-ee: initial phase of FCC; operates at different EW thresholds:
 5×10^{12} Z bosons, 10^8 WW pairs, $>10^6$ Higgses, $>10^6$ tt pairs

Attribute	$\Upsilon(4S)$	pp	Z^0	Particle species	B^0	B^+	B_s^0	Λ_b	B_c^+	$c\bar{c}$	$\tau^-\tau^+$
All hadron species		✓	✓	Yield ($\times 10^9$)	310	310	75	65	1.5	600	170
High boost		✓	✓	[FCC Physics Opportunities, Conceptual Design Report] [Flavour cases: EPJPlus 136, 837 arXiv:2106.01259 , and EPJPlus 136, 912 arXiv:2106.12168]							
Enormous production cross-section		✓									
Negligible trigger losses	✓		✓								
Low backgrounds	✓		✓								
Initial energy constraint	✓		(✓)								

Experimental and theoretical inputs

	Central values	Uncertainties				Reference Phases I–III
		Current [28]	Phase I	Phase II	Phase III	
$ V_{ud} $	0.97437	± 0.00021	id	id	id	[28]
$ V_{us} f_+^{K \rightarrow \pi}(0)$	0.2177	± 0.0004	id	id	id	[28]
$ V_{cd} $	0.2248	± 0.0043	± 0.003	id	id	[40,41]
$ V_{cs} $	0.9735	± 0.0094	id	id	id	[28,40,41]
Δm_d [ps ⁻¹]	0.5065	± 0.0019	id	id	id	[17]
Δm_s [ps ⁻¹]	17.757	± 0.021	id	id	id	[17]
$ V_{cb} _{\text{SL}} \times 10^3$	42.26	± 0.58	± 0.60	± 0.44	id	[29]
$ V_{cb} _{W \rightarrow cb} \times 10^3$	± 0.17	[34–36]
$ V_{ub} _{\text{SL}} \times 10^3$	3.56	± 0.22	± 0.042	± 0.032	id	[29]
$ V_{ub} / V_{cb} $ (from Λ_b)	0.0842	± 0.0050	± 0.0025	± 0.0008	id	[30]
$\mathcal{B}(B \rightarrow \tau \nu) \times 10^4$	0.83	± 0.24	± 0.04	± 0.02	± 0.009	[29,34]
$\mathcal{B}(B \rightarrow \mu \nu) \times 10^6$	0.37	...	± 0.03	± 0.02	id	[29]
$\sin 2\beta$	0.680	± 0.017	± 0.005	± 0.002	± 0.0008	[29,30,34]
α [°] (mod 180°)	91.9	± 4.4	± 0.6	id	id	[29]
γ [°] (mod 180°)	66.7	± 5.6	± 1	± 0.25	± 0.20	[29,30,34]
β_s [rad]	-0.035	± 0.021	± 0.014	± 0.004	± 0.002	[30,34]
$A_{\text{SL}}^d \times 10^4$	-6	± 19	± 5	± 2	± 0.25	[14,17,34,37]
$A_{\text{SL}}^s \times 10^5$	3	± 300	± 70	± 30	± 2.5	[14,17,34,37]
\bar{m}_t [GeV]	165.30	± 0.32	id	id	± 0.020	[28,34]
$\alpha_s(m_Z)$	0.1185	± 0.0011	id	id	± 0.00003	[28,34]
$f_+^{K \rightarrow \pi}(0)$	0.9681	± 0.0026	± 0.0012	id	id	[30]
f_K [GeV]	0.1552	± 0.0006	± 0.0005	id	id	[30]
f_{B_s} [GeV]	0.2315	± 0.0020	± 0.0011	id	id	[30]
B_{B_s}	1.219	± 0.034	± 0.010	± 0.007	id	[30]
f_{B_s}/f_{B_d}	1.204	± 0.007	± 0.005	id	id	[30]
B_{B_s}/B_{B_d}	1.054	± 0.019	± 0.005	± 0.003	id	[30]
$\tilde{B}_{B_s}/\tilde{B}_{B_d}$	1.02	± 0.05	± 0.013	id	id	[30,42,43]
\tilde{B}_{B_s}	0.98	± 0.12	± 0.035	id	id	[30,42,43]
η_B	0.5522	± 0.0022	id	id	id	[44]

[See for refs.: PRD 102, 056023 (2020), arXiv:2006.04824
and also EPJPlus 136, 837 arXiv:2106.01259,
and EPJPlus 136, 912 arXiv:2106.12168]

Experimental and theoretical inputs




- **Caveat:** experimental sensitivity studies for FCC still in progress...
- $|V_{ub}|_{SL}$ & $|V_{cb}|_{SL}$: dominance of stat. Uncs. @ Phase II (Belle II U.)
 $|V_{cb}|$ accuracy $\sim 0.4\%$ @ FCC-ee: $W \rightarrow bc$ [Schune, Monteil]
[incl. vs excl. inputs and impact on extraction of NP in $|\Delta F|=2$: De Bruyn, Fleischer, Malami, van Vliet]
[recent B_c to $\tau\nu$ @ Tera-Z: Zheng, Xu, Cao, Yu, Wang, Prell, Cheung, Ruan; Amhis, Hartmann, Helsen, Hill, Sumensari]
- **Future stat. accuracy in angles** ($\alpha, \beta, \beta_s, \gamma$) $\sim < 1^\circ$
At this level of exp. accuracy: need for theo. studies of isospin breaking corrections, penguin pollution, etc.
[recent dedicated α @ Tera-Z (B^0 to $\pi^0\pi^0$): Wang, Descotes-Genon, Deschamps, Li, Chen, Zhu, Ruan]
[recent Tera-Z studies of β_s , etc.: Aleksan, Oliver, Perez; Aleksan, Oliver, Perez; Li, Ruan, Zhao]
- Phases I and II uncs. for **Lattice QCD** (decay constants, bag parameters) $< 1\%$

Literature discusses Lattice QCD projections up to Phase II

Future improvements

- SM reference: shift the central values
- Compared to Current, **improvement by factor >3 (5) at Phase I (II)**




Sensitivities	Summer 2019	Phase I	Phase II
h_d	0.26	0.073	0.049
h_s	0.12	0.065	0.044



Bottlenecks

Necessary improvements beyond current expectations
for enhancing sensitivity to NP:

- **Lattice QCD** (also short-distance QCD corrections)
- $|V_{cb}|$, overall normalization (Wolfenstein parameter A)
- **Individual impacts on h_d and h_s : O(20-30)%**



Future reach to NP in B meson mixing

$$\frac{C_{ij}^2}{\Lambda^2} (\bar{q}_{i,L} \gamma_\mu q_{j,L})^2, \quad \longrightarrow \quad h \simeq 1.5 \frac{|C_{ij}|^2 (4\pi)^2}{|\lambda_{ij}^t|^2 G_F \Lambda^2} \simeq \frac{|C_{ij}|^2}{|\lambda_{ij}^t|^2} \left(\frac{4.5 \text{ TeV}}{\Lambda} \right)^2,$$



$$\sigma = \arg(C_{ij} \lambda_{ij}^{t*}),$$

- In the absence of suppressions: **NP scale >> TeV**
- Possible flavour and loop suppressions: **alleviate bounds on NP**

Couplings	NP loop order	Sensitivity for Summer 2019 [TeV]		Phase I Sensitivity [TeV]		Phase II Sensitivity [TeV]	
		B_d mixing	B_s mixing	B_d mixing	B_s mixing	B_d mixing	B_s mixing
$ C_{ij} = V_{ti} V_{tj}^* $ (CKM-like)	Tree level	9	13	17	18	20	21
	One loop	0.7	1.0	1.3	1.4	1.6	1.7
$ C_{ij} = 1$ (No hierarchy)	Tree level	1×10^3	3×10^2	2×10^3	4×10^2	2×10^3	5×10^2
	One loop	80	20	2×10^2	30	2×10^2	40

Discovery prospects

B meson mixing observables also provide potential **discovery for NP**



Conclusions

- Flavour physics: crucial in **shaping the SM**, but also in looking for **candidates of NP**
- **$|\Delta B|=2$** : only one flavour aspect of future experimental and theoretical progress
- Allowed NP in B meson mixing **still large**: bounds will largely improve
- Identified future limitations in Phase II:
LQCD and $|V_{cb}|$
- **Complementarity**: FCC-ee addresses $|V_{cb}|$ bottleneck

BACK UP!

Experimental quantities vs. theoretical ones

Observables considered in the fit that are modified by NP in $|\Delta B|=2$:

$$\Delta_q = |\Delta_q| e^{i2\Phi_q^{\text{NP}}}$$

parameter	prediction in the presence of NP
Δm_q	$ \Delta_q^{\text{NP}} \times \Delta m_q^{\text{SM}}$
2β	$2\beta^{\text{SM}} + \Phi_d^{\text{NP}}$
$2\beta_s$	$2\beta_s^{\text{SM}} - \Phi_s^{\text{NP}}$
2α	$2(\pi - \beta^{\text{SM}} - \gamma) - \Phi_d^{\text{NP}}$
$\Phi_{12,q} = \text{Arg}\left[-\frac{M_{12,q}}{\Gamma_{12,q}}\right]$	$\Phi_{12,q}^{\text{SM}} + \Phi_q^{\text{NP}}$
A_{SL}^q	$\frac{\Gamma_{12,q}}{M_{12,q}^{\text{SM}}} \times \frac{\sin(\Phi_{12,q}^{\text{SM}} + \Phi_q^{\text{NP}})}{ \Delta_q^{\text{NP}} }$
$\Delta\Gamma_q$	$2 \Gamma_{12,q} \times \cos(\Phi_{12,q}^{\text{SM}} + \Phi_q^{\text{NP}})$

Different representation

$$\langle B_q | \mathcal{H}_{\Delta B=2}^{\text{SM+NP}} | \bar{B}_q \rangle \equiv \langle B_q | \mathcal{H}_{\Delta B=2}^{\text{SM}} | \bar{B}_q \rangle \times (\text{Re}(\Delta_q) + i \text{Im}(\Delta_q))$$

$$\text{Re}(\Delta_q) + i \text{Im}(\Delta_q) = r_q^2 e^{i2\theta_q} = 1 + h_q e^{i\sigma_q}$$

Soares & Wolfenstein, PRD 47, 1021 (1993)
 Deshpande, Dutta & Oh, PRL77, 4499 (1996)
 Silva & Wolfenstein, PRD 55, 5331 (1997)
 Cohen et al., PRL78, 2300 (1997)
 Grossman, Nir & Worah, PLB 407, 307 (1997)

