



# Measurements of $B_{(s)} \rightarrow \mu\mu$ decays

Dmytro Kovalskyi for the CMS Collaboration



# Why $B_{(s)} \rightarrow \mu\mu$ ?



- Standard Model Prediction
  - $B_s \rightarrow \mu\mu: (3.66 \pm 0.14) \times 10^{-9}$
  - $B^0 \rightarrow \mu\mu: (1.03 \pm 0.05) \times 10^{-10}$
  - Beneke, Bobeth, arXiv:1908.07011
- Unique rare  $b \rightarrow s \ell \ell$  decays
  - Sensitive to New Physics effects
  - Non perturbative hadronic contributions enter via  $B_{(s)}$  decay constant
  - Well known from Lattice QCD
- Note
  - SM predictions scale as  $|V_{cb}|^2$
  - Difference between inclusive and exclusive measurements of  $|V_{cb}|$  leads to additional uncertainty
  - Buras, Venturini, arXiv:2203:11960




# Current Results



- Latest combination of ATLAS, CMS, LHCb results is based on RunI and partial Run2 data - [BPH-20-003](#)
- Full RunI+Run2 results by LHCb - [PRL 128\(2022\) 041801](#)

# Rare B decay Anomalies



Altmannshofer, Stangl, arXiv:2103.13370

- Multiple discrepancies are observed in rare B decays
  - $3.1\sigma$  Lepton Flavour Universality violation in  $R(K)$  and  $R(K^*)$
  - $2-3\sigma$  discrepancies in branching fraction and angular observables
- Global fits provide good description of data with the Wilson coefficients  $C_9$  and  $C_{10}$  of the vector and pseudo-vector operators  $O_9$  and  $O_{10}$  in the effective 4-fermion interaction
  - Only  $O_{10}$  operator contributions to  $B_{(s)} \rightarrow \mu\mu$

# Branching Fraction Measurement



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = \mathcal{B}(B^+ \rightarrow J/\psi K^+) \times \frac{N_{B_s^0 \rightarrow \mu^+ \mu^-}}{N_{B^+ \rightarrow J/\psi K^+}} \times \frac{\epsilon_{B^+ \rightarrow J/\psi K^+}}{\epsilon_{B_s^0 \rightarrow \mu^+ \mu^-}} \times \frac{f_u}{f_s}$$

or  $\left\{ = \mathcal{B}(B_s^0 \rightarrow J/\psi \phi) \times \frac{N_{B_s^0 \rightarrow \mu^+ \mu^-}}{N_{B_s^0 \rightarrow J/\psi \phi}} \times \frac{\epsilon_{B_s^0 \rightarrow J/\psi \phi}}{\epsilon_{B_s^0 \rightarrow \mu^+ \mu^-}} \right\}$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = \mathcal{B}(B^+ \rightarrow J/\psi K^+) \times \frac{N_{B^0 \rightarrow \mu^+ \mu^-}}{N_{B^+ \rightarrow J/\psi K^+}} \times \frac{\epsilon_{B^+ \rightarrow J/\psi K^+}}{\epsilon_{B^0 \rightarrow \mu^+ \mu^-}} \times \frac{f_u}{f_d} = 1$$

external  
B production  
fraction ratio

- Branching Fraction are normalized using  $B^+ \rightarrow J/\psi K^+$  and  $B_s \rightarrow J/\psi \phi$  decays
  - Most systematic effects cancel in the ratio
  - $B^+ \rightarrow J/\psi K^+$  provides nominal normalization
    - $B_s \rightarrow J/\psi \phi$  - alternative normalization with different systematics that can benefit from independent BF measurement of  $B_s \rightarrow J/\psi \phi$

# Event Preselection




Selection	$B_s^0 \rightarrow \mu^+ \mu^-$	$B^+ \rightarrow J/\psi K^+$	$B_s^0 \rightarrow J/\psi \phi$
B candidate mass [ GeV ]	[4.90,5.90]	[4.90,5.90]	[4.90,5.90]
Blinding window [ GeV ]	[5.15,5.50]		
$p_{T\mu}$ [ GeV ]	> 4	> 4	> 4
$ \eta_\mu $	< 1.4	< 1.4	< 1.4
3D SV displacement significance	> 6	> 4	> 4
$p_{T\mu\mu}$ [ GeV ]	> 5	> 7	> 7
$\mu\mu$ SV probability	> 0.025	> 0.1	> 0.1
$J/\psi$ candidate mass [ GeV ]		[2.9,3.3]	[2.9,3.3]
Kaon $p_T$ [ GeV ]		> 1	> 1
Mass-constrained fit probability		> 0.025	> 0.025
2D $\mu\mu$ pointing angle [rad]		< 0.4	< 0.4
$\phi$ candidate mass [ GeV ]			[1.01, 1.03]

- Selection requirements are as loose as possible
  - Provide more data to MultiVariate Analysis (MVA)
  - Limited by trigger requirements
- Normalization channel selection is optimized to match kinematics of signal

# Dominant Contributions




3-body and partial decays




Muons from the same  
B hadron




Signal  $B_s \rightarrow \mu\mu$





Combinatorial Background



Muons originate from  
different B hadrons




# Muon Fakes



- Double muon fakes from  $B \rightarrow hh$  - non-trivial background
  - Looks like signal
  - Rate is comparable to  $B^0 \rightarrow \mu\mu$
  - $B \rightarrow K\pi$  and  $B_s \rightarrow KK$  are dominant contribution
- Primary source of fakes
  - Pion and kaon decays in flight to muon and neutrino
  - Other contributions are negligible and easy to reject
- Used MVA based muon identification
  - Detect minor imperfections in the muon candidate trajectory
  - Factor of 2-3 better rejection of fakes than the standard muon selection
    - Kaon decays are easier to reject
- Fake rates are measured in  $K_s \rightarrow \pi\pi$  and  $\phi \rightarrow KK$  control samples
  - Simulated reasonably well: ~25% systematic per hadron

# Multivariate Analysis

# Multivariate Analysis




- New multivariate analysis (MVA<sub>B</sub>) used to suppress the dominant backgrounds
  - Trained with signal MC and mass sideband data with the XGBoost package (advanced gradient boosting algorithm)
- Most discriminating variables
  - Pointing angles:  $\alpha_{2D}, \alpha_{3D}$
  - Impact parameter and its significance:  $\delta_{3D}, \delta_{3D}/\sigma(\delta_{3D})$
  - Flight length and its significance:  $\ell_{3D}/\sigma(\ell_{3D})$
  - Isolation for B candidate and muons
  - Dimuon vertex quality

# MVA in Data




CMS *Simulation Preliminary*




- MVA mismodeling can be a major source of systematics
  - Need a data control sample
  - $B^+ \rightarrow J/\psi K^+$  is the best candidate
- MVA is trained to rejected  $\mu\mu K$  events
  - Extra track, wrong pointing angle etc
- Need to use correct input to get signal-like response
  - $\mu\mu K$ : pointing angle, impact parameter
  - $\mu\mu$ : vertex probability, displacement, isolation (ignore kaon)

# $B^+ \rightarrow J/\psi K^+$ Event Selection




- The pointing angle is one of the most powerful discriminators
- The distribution has a non-trivial dependence on the decay kinematics
  - Need to match  $B^+ \rightarrow J/\psi K^+$  phase space to  $B_s \rightarrow \mu\mu$
- The agreement between  $\alpha(\mu\mu)$  and  $\alpha(\mu\mu K)$  is getting better when the kaon is soft
  - Leads to larger background, but it's manageable thanks to large number of events
    - Use sPlot technique to extract distributions in Data

# Flight length significance




Default



Scaled

- Soft kaon requirement gives matching distributions for all input variables but the flight length significance
- It correlates with the dimuon mass
  - Smaller mass  $\rightarrow$  smaller opening angle  $\rightarrow$  larger uncertainty on vertex position along the trajectory  $\rightarrow$  smaller significance
- Scaling flight length significance by 1.6 provides a decent matching



# XGBoost Reweighting



- With the special  $B^+ \rightarrow J/\psi K^+$  control sample we can correct for mismodeling in MC simulation
- Trained XGBoost classifier on MC vs Data with sWeights
  - Capture the difference between MC and Data
  - Use  $w = \frac{\text{Prob}_{\text{Data}}}{\text{Prob}_{\text{MC}}}$  as a weight to reweight  $B_{(s)} \rightarrow \mu\mu$  MC samples

# Branching Fraction Measurements

# Unbinned ML Fit



- 2D unbinned ML fit in mass and its uncertainty
- 16 categories
  - rapidity of forward muon: [0,0.7] and [0.7,1.4]
  - data taking period: 2016a, 2016b, 2017, 2018
  - MVA: loose and tight
- Shapes
  - Signal - fixed CrystalBall with mass scale and resolution corrected using J/ $\psi$  and Y(IS)
  - Combinatorial - 1st order polynomial with free floating parameters
  - Semileptonic - Gaussian with floating parameters


# Postfit Event Yields



Data set	Channel	$N(B_s^0)$	$N(B^0)$	$N(\text{comb})$	$N(\text{peak})$	$N(\text{semi})$	$N(\text{total})$	Data
$\text{MVA}_B > 0.99$								
2016a	0	5.3	0.2	2.8	0.2	6.0	14.5	16
2016a	1	9.4	0.4	16.2	0.4	9.9	36.3	35
2016b	0	6.3	0.3	1.7	0.2	7.9	16.4	12
2016b	1	9.9	0.4	8.6	0.4	13.3	32.6	32
2017	0	23.5	1.0	51.4	0.8	29.6	106.3	114
2017	1	33.9	1.3	89.6	1.4	44.0	170.2	165
2018	0	34.5	1.4	64.8	1.3	38.4	140.4	143
2018	1	50.0	2.0	151.0	2.5	50.9	256.4	252
$0.99 > \text{MVA}_B > 0.9$								
2016a	0	4.8	0.2	118.0	0.2	8.4	131.6	132
2016a	1	8.9	0.4	324.8	0.4	16.5	351.0	352
2016b	0	5.6	0.2	107.6	0.2	10.9	124.5	126
2016b	1	9.2	0.4	257.1	0.4	18.2	285.3	287
2017	0	15.2	0.6	637.7	0.7	26.4	680.6	683
2017	1	21.7	0.9	1430.5	1.1	44.3	1498.5	1498
2018	0	23.3	1.0	936.2	1.2	52.5	1014.2	1017
2018	1	34.2	1.4	2222.5	1.8	79.7	2339.6	2340

- Postfit event yields by categories
  - Channel 0 and I refer to forward muon rapidity ranges of [0.0,0.7] and [0.7,1.4]
  - Expected number of  $B^0$  events is still very small


# Efficiency Correction



- Use  $B^+ \rightarrow J/\psi K^+$  events to model  $B_{(s)} \rightarrow \mu\mu$  decays in Data and MC
  - Additional selection requirements
    - Kaon  $P_T < 1.5 \text{ GeV}$
    - Rescaled flight length significance
- Data/MC correction factors
  - Ratio method: efficiency ratio between weighted data/MC
  - XGBoost: train a XGBoost classifier to reweight MC to match to the data


Method	MVA <sub>B</sub> >0.9			MVA <sub>B</sub> >0.99		
	2016	2017	2018	2016	2017	2018
Ratio	1.011±0.013	0.939±0.007	0.903±0.008	1.058±0.019	0.891±0.008	0.885±0.010
XGBoost	0.991±0.008	0.949±0.003	0.917±0.002	1.008±0.011	0.905±0.004	0.908±0.002

# $B^+ \rightarrow J/\psi K^+$ Yield Fits



- Two different signal models are used to extract signal normalization in unbinned maximum likelihood fits
  - Nominal model is built using analytical functions
  - Alternative one is using non-parametric signal model convolved with a resolution model
- The difference between the two estimates is taken as systematics (1%)

# $B_s \rightarrow J/\psi\varphi$ Yield Fits



- $B_s \rightarrow J/\psi\varphi$  is used as an alternative normalization for  $B_s$  BF
  - At the moment it's primarily a cross-check
- Data is fitted to two different signal models, taking the difference as systematics

# BF Systematics




Effect	$\text{BF}(B_s \rightarrow \mu^+ \mu^-)$	$\text{BF}(B^0 \rightarrow \mu^+ \mu^-)$
Trigger efficiency	2–4%	
Pileup	1%	
Vertex quality	1%	
MVA <sub>B</sub> correction	2–3%	
Tracking efficiency	2.3%	
J/ $\psi$ K <sup>+</sup> shape	1%	
Fit bias	2.2%	4.5%
$f_s/f_u$ ratio	3.5%	-

- Signal efficiency is correlated with the  $B_s$  lifetime
  - $B_s \rightarrow \mu\mu$  branching fractions are measured assuming the SM lifetime value
  - For alternative hypothesis scale BF using the following scale factor


$$\alpha_{\text{BF}} = 1.577 - 0.358 \tau$$

- $\tau$  is in ps
  - Example:  $\alpha_{\text{BF}}(1.61) = 1.0$ ,  $\alpha_{\text{BF}}(1.43) = 1.065$

# $f_s/f_u$ ratio in BF fit



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = \mathcal{B}(B^+ \rightarrow J/\psi K^+) \frac{N_{B_s^0 \rightarrow \mu^+ \mu^-}}{N_{B^+ \rightarrow J/\psi K^+}} \frac{\epsilon_{B^+ \rightarrow J/\psi K^+} f_u}{\epsilon_{B_s^0 \rightarrow \mu^+ \mu^-} f_s},$$




- $f_s/f_u = 0.231 \pm 0.008$
- Based on  $P_T$ -dependent results from LHCb
  - PRD 104 (2021) 032005
  - Integrate with the effective  $P_T$  distribution
  - Previous measurement used  $0.252 \pm 0.032$
- Resulting BF can be rescaled:
  - One can use a different  $f_s/f_u$  value
  - Treated as an external uncertainty
    - not as a constrained nuisance parameter

# $B_s \rightarrow \mu\mu$ BF Result



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = [3.83^{+0.38}_{-0.36} \text{ (stat)}^{+0.19}_{-0.16} \text{ (syst)}^{+0.14}_{-0.13} (f_s/f_u)] \times 10^{-9}$$


Alternative using  $B_s \rightarrow J/\psi \phi$ :  $\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = [3.95^{+0.39}_{-0.37} \text{ (stat)}^{+0.27}_{-0.22} \text{ (syst)}^{+0.21}_{-0.19} \text{ (BF)}] \times 10^{-9}$





# $B^0 \rightarrow \mu^+ \mu^-$ Search



$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = [0.37^{+0.75}_{-0.67} \text{ (stat)}^{+0.08}_{-0.09} \text{ (syst)}] \times 10^{-10}$$





# Sensitivity to $B^0 \rightarrow \mu\mu$




- The main challenge with  $B^0 \rightarrow \mu\mu$  is the combinatorial background
- It will require more data and analysis improvements to reach discovery level

# Contour Plots






# Lifetime Measurement

# Effective Lifetime Measurement



- In the absence of CP violation only the heavy  $B_s$  state decays into dimuon
  - Different composition of states may be allowed by New Physics.
- Plot shows decay time projection
- Efficiency correction
  - Decay time efficiency derived from MC
  - Corrected by  $B^+ \rightarrow J/\psi K^+$  data to mitigate the bias from tight  $MVA_B$  requirement.
  - The residual bias and the difference between  $B_s \rightarrow \mu\mu$  and  $B^+ \rightarrow J/\psi K^+$  are considered as a systematic uncertainty.

# Lifetime Systematics




Effect	2016a	2016b	2017	2018
Efficiency modeling		<b>0.01 ps</b>		
Scanning over different gen lifetime sample		<b>0.01 ps</b>		
Decay time mis-modeling	<b>0.10 ps</b>	<b>0.06 ps</b>	<b>0.02 ps</b>	<b>0.02 ps</b>
Lifetime bias	<b>0.04 ps</b>	<b>0.04 ps</b>	<b>0.05 ps</b>	<b>0.04 ps</b>
<b>Total</b>	<b>0.11 ps</b>	<b>0.07 ps</b>	<b>0.05 ps</b>	<b>0.04 ps</b>

- Dominant systematics comes from a strong correlation between  $\text{MVA}_B$  and decay time, which are hard to model well
  - Corrections can be derived in data and uncertainty is mostly limited by the size of the control sample

# Effective Lifetime



$$\tau = 1.83^{+0.23}_{-0.20} \text{ (stat)}^{+0.04}_{-0.04} \text{ (syst) ps}$$




# Summary



- Studies of rare  $B_{(s)} \rightarrow \mu^+ \mu^-$  decays provides a unique tool to explore and understand rare B decay anomalies
  - Theoretically clean
  - Sensitive to the same processes
- CMS finalized analysis of  $140 \text{ fb}^{-1}$  data collected during LHC Run-2
  - All results are consistent with SM predictions
- Relative uncertainty on  $\text{BF}(B_s \rightarrow \mu^+ \mu^-)$  has been reduced to 11%
  - The best single measurement to date
- Statistical uncertainties dominate in all measurements
  - Good perspectives for further improvements with Run-3 data
- A conference note and other materials are available at
  - [https://cms-results.web.cern.ch/cms-results/public-results/preliminary-results/  
BPH-21-006/index.html](https://cms-results.web.cern.ch/cms-results/public-results/preliminary-results/BPH-21-006/index.html)

# Back Up

# Analysis Overview



# Event Display

