Azimuthal anisotropic flow of identified hadrons in Au + Au collisions at BES-II energies at STAR

Prabhupada Dixit
(For the STAR Collaboration)
Indian Institute of Science Education and Research (IISER) Berhampur

Critical Point and Onset of Deconfinement (CPOD), 2022

Supported in part by the
U.S. DEPARTMENT OF ENERGY | Office of Science

Outline

- **Introduction & motivation**
- **STAR detectors**
- **Analysis details**
- **Results**
  - ✯ p_T dependence of v_n
  - ✯ Centrality dependence
  - ✯ NCQ Scaling
  - ✯ $v_3/v_2^{3/2}$ ratio
- **Summary**

Introduction & motivation





Elliptic flow coefficient (v_2) : Initial spatial anisotropy (dominant source) + Event-by-event fluctuations

Triangular flow coefficient (v_3) : Event-by-event fluctuations in the overlap region

$$\frac{dN}{d(\phi - \Psi_n)} = N_0 \left[1 + \sum_{n=1}^{\infty} 2v_n \cos(\phi - \Psi_n) \right]$$

$$v_n = \langle\langle \cos n(\phi - \Psi_n) \rangle\rangle$$

Importance of v_2 and v_3

☞ Sensitive to the initial state and transport properties of the medium.

☞ Measurements of v_2 and v_3 are important to constrain the models. C. Shen *et al* JPG 38 (2011) 124045

Introduction & motivation

Results from RHIC BES-I

STAR Phys. Rev. C 93 (2016) 14907



☞ ϕ mesons seem to deviate from the NCQ scaling at $\sqrt{s_{NN}} < 19.6$ GeV.

☞ But statistics is not significant to draw any conclusion.

★ High Statistics data from BES-II enable us to measure v_2 and v_3 of multi-strange hadrons and φ mesons with high precession specifically at low energy regime.

STAR detectors and particle identification



Time Projection Chamber (TPC)



Time of Flight (ToF)



- Full azimuthal coverage
- Excellent particle identification capability

BES-II upgrades

- iTPC upgrade: Larger pseudorapidity coverage (-1.5 < η < 1.5)
- Better dE/dx and momentum resolution.
- Better track quality.

Reconstruction of (multi-)strange hadrons and ϕ mesons







Data set information for this analysis:

- System: Au+Au
- Year: 2019 (BES-II data)
- Collision energy: 19.6 and 14.6 GeV
- #Events: ~380M (19.6 GeV) & ~400M (14.6 GeV)
- Source of systematic uncertainty: Variation of analysis cuts e.g. collision vertex selection cuts, particle identification cuts, quality track selection cuts etc.

Analysis details

The nth order flow coefficient is given by

$$v_n = \langle\langle \cos n(\phi - \Psi_n) \rangle\rangle$$



- The reaction plane of the collision can not be determined directly from the experiment.
- The event plane is used as a proxy for the reaction plane.

Event plane resolution

$$R_n = \langle \cos n(\Psi_n - \Psi_R) \rangle$$

Experimentrally, $R_n(sub) = \sqrt{\langle \cos n(\Psi_A - \Psi_B) \rangle}$



Event plane determination

$$\Psi_n = \frac{1}{n}\tan^{-1}\left(\frac{Q_y}{Q_x}\right)$$

$$Q_x = \sum_i w_i \cos(n\phi_i)$$

$$Q_y = \sum_i w_i \sin(n\phi_i)$$

To minimize non-flow correlation

★ **Sub-event plane method is used to calculate v_n.**

★ **η gap of 0.1 is taken between two sub-event planes Ψ_A ($-1.5 < \eta < -0.05$) and Ψ_B ($0.05 < \eta < 1.5$).**

★ **To calculate v_n of a particle in negative η region, event plane from positive η side is used and vice versa.**

The weight factor $w_i = p_T \times \phi$-weight.

ϕ-weight: accounts for the azimuthal acceptance correction of the detectors.

Analysis details

v_n measurements for (multi-)strange hadrons and ϕ mesons

Invariant mass method

N. Borghini and J.-Y. Ollitrault, Phys. Rev. C 70, 064905 (2004)



Event plane method

A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C 58, 1671 (1998)



$$v_n^{S+B}(M_{inv}) = \langle cos\left[n(\phi-\psi_n)\right]\rangle = v_n^S\frac{S}{S+B}(M_{inv}) + v_n^B\frac{B}{S+B}(M_{inv})$$

$$\frac{dN}{d(\phi-\psi_n)} = \frac{N_0}{2\pi}\left(1+2v_n\cos n(\phi-\psi_n)\right)$$

$$v_n^B(M_{inv}) = p_0 + p_1 M_{inv}$$

Results: p_T dependence of v_2 @19.6 GeV



☞ Mass ordering observed in the low p_T region ($p_T < 1.5$ GeV/c) : Radial flow

☞ Baryon to meson separation observed in the high p_T region : Quark coalescence

☆ **The statistical errors are reduced by a factor of ~3 compared to BES-I.**

Results: Centrality dependence of v_2 @19.6 GeV



☞ **Strong centrality dependence of v_2 → Spatial anisotropy is a dominant cause for v_2**

Results: Centrality dependence of v_2 @14.6 GeV



☞ Strong centrality dependence of v_2 → Spatial anisotropy is a dominant cause for v_2

Results: Centrality dependence of v_3 @19.6 GeV



☞ Weak centrality dependence of v_3 → Event-by-event fluctuation is a dominant cause for v_3

Results: NCQ scaling in v_2 @19.6 GeV



☞ The scaling for v_2 holds within 20% for particles and within 10% for anti-particles (except at low p_T for $\bar{\Lambda}$ and $\bar{p}$)

Partonic collectivity in the initial stage of the system and hadronization via coalescence.

Results: NCQ scaling in v_3 @19.6 GeV





⋆ p_0 is the parameter from the simultaneous 0th order polynomial fit to the ratios.

☞ The modified scaling for v_3 holds within 30% for particles and within 15% for anti-particles.

PHENIX Phys. Rev. C 93, 051902 (2016)

Results: NCQ scaling in v_2 @14.6 GeV





- ☞ The scaling for v_2 holds within 15% for the (multi-)strange hadrons except low p_T $\bar{\Lambda}$.
- ☞ ϕ mesons are following the NCQ scaling at 14.6 GeV.
- ☞ The rising trend in the K_S^0 v_2 at $(m_T - m_0)/n_q > 1$ GeV/c^2 may arise due to the non-flow contribution. Non-flow estimation is underway.

☆ See S. Zhou's talk for light hadrons v_2 at 14.6 GeV

Results: $v_3/v_2^{3/2}$ ratio @19.6 GeV

Correlation between v_3 and $v_2^{3/2}$

Previous RHIC measurements suggests that $v_n \propto v_2^{n/2}$

STAR Phys. Rev. Lett. 92, 062301 (2004)
PHENIX Phys. Rev. Lett. 105, 062301 (2010)
STAR Phys. Rev. C 103, 064907 (2021)



☞ The ratio $v_3/v_2^{3/2}$ shows non-trivial p_T dependence.

☞ $v_3/v_2^{3/2}$ ratios are sensitive to the initial state fluctuations and transport properties of the medium.

E. Retinskaya et al. Phys. Rev. C 89, 014902 (2014)

Summary

☞ Using high statistics BES-II data, precise measurements of v_2 of identified hadrons in 19.6 and 14.6 GeV Au+Au collisions have been presented, with improved statistical significance by a factor of 3 compared to BES-I.

☞ New results of v_2 and v_3 of (multi-)strange hadrons and ϕ mesons are presented.

p_T dependence of v_2

☞ Confirmation of usual trend of mass ordering in v_2 at low p_T and baryon-meson separation at high p_T in low energies at 19.6 GeV using strange and multi-strange hadrons.

Centrality dependence of v_n

☞ Strong centrality dependence of v_2 : initial spatial anisotropy is a dominant cause for v_2.

☞ Weak centrality dependence of v_3: event-by-event fluctuation is a dominant cause for v_3.

NCQ scaling

☞ The NCQ scaling holds for both particles and anti-particles.

☞ The scaling holds for ϕ mesons at 14.6 GeV.

☞ The scaling suggests the collectivity in the partonic phase of the system and hadronization via quark coalescence.

$v_3/v_2^{3/2}$ ratio

☞ The ratio shows weak dependence of p_T above $p_T > 1.0$ GeV/c.

☞ Can be used to constrain the initial state fluctuations and η/s of the medium.

Thank you …