

# CP violation and determination of the “flat” (b,s) unitarity triangle at FCC-ee


R. Aleksan (CEA/IRFU), L. Oliver (IJCLab), E.Perez (CERN)

FCC Flavor Physics  
Workshop  
12/9/2022

$$V_{ub}^* V_{us} + V_{cb}^* V_{cs} + V_{tb}^* V_{ts} \stackrel{?}{=} 0$$

$$\lambda^4, \quad \lambda^2, \quad \lambda^2$$

$$V_{CKM} = \begin{bmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{bmatrix}$$



$$\alpha_s = \arg \left( -\frac{V_{ub}^* V_{us}}{V_{tb}^* V_{ts}} \right), \quad \beta_s = \arg \left( -\frac{V_{tb}^* V_{ts}}{V_{cb}^* V_{cs}} \right), \quad \gamma_s = \arg \left( -\frac{V_{cb}^* V_{cs}}{V_{ub}^* V_{us}} \right) \approx (67^\circ, 1^\circ, 111^\circ)$$

$$B_s \rightarrow D_s K$$

$$B_s \rightarrow J/\psi \phi$$

$$D^\pm \rightarrow D^0 (\bar{D}^0) K^\pm$$

- CP violation and determination of the bs "flat" unitarity triangle at FCC-ee, <https://arxiv.org/abs/2107.02002>
- Study of CP violation in  $B^\pm \rightarrow D^0 (\bar{D}^0) K^\pm$  at FCC-ee <https://arxiv.org/abs/2107.05311>

# Detector response

## ➤ Modelisation of the detector response :

- Detailed description of tracks, accounting for multiple scattering

$$\text{Acceptance : } |\cos \theta| < 0.95$$

$$\text{Track } p_T \text{ resolution : } \frac{\sigma(p_T)}{p_T^2} = 2. \times 10^{-5} \oplus \frac{1.2 \times 10^{-3}}{p_T \sin \theta}$$


$$\text{Track } \phi, \theta \text{ resolution : } \sigma(\phi, \theta) \mu\text{rad} = 18 \oplus \frac{1.5 \times 10^3}{p_T \sqrt[3]{\sin \theta}}$$

$$\text{Vertex resolution : } \sigma(d_{\text{Im}}) \mu\text{m} = 1.8 \oplus \frac{5.4 \times 10^1}{p_T \sqrt{\sin \theta}}$$

$$\text{Vertex resolution : } \langle \sigma(d_{\text{Im}}) \rangle \text{ bachelor } K \text{ in } D_s K$$

$$\langle \sigma(d_{\text{Im}}) \rangle \simeq 10 \mu\text{m}$$

$$\text{Calorimeter resolution : } \frac{\sigma(E)}{E} = \frac{3 \times 10^{-2}}{\sqrt{E}} \oplus 5 \times 10^{-3}$$




## ➤ For vertexing Full MC events + response of the IDEA detector with DELPHES

- Genuine vertex fitting

$\beta_S = \arg\left(-\frac{V_{tb}^* V_{ts}}{V_{cb}^* V_{cs}}\right)$ : CP violation with  $B_s \rightarrow J/\psi\phi \rightarrow \mu^+ \mu^- K^+ K^-$

$\int L dt = 150 ab^{-1}$

$\sim 6 \cdot 10^6 B_s(\bar{B}_S)$  evts @ FCCee



$$\Gamma(B_s(t) \rightarrow f_{CP}) = |\langle f_{CP} | B_s \rangle|^2 e^{-\Gamma t} \{1 - (1 - 2\omega)\eta_f \sin \phi_{CP} \sin \Delta m t\}$$

$$\Gamma(\bar{B}_s(t) \rightarrow f_{CP}) = |\langle f_{CP} | \bar{B}_s \rangle|^2 e^{-\Gamma t} \{1 + (1 - 2\omega)\eta_f \sin \phi_{CP} \sin \Delta m t\}$$

$V_{cb} V_{cs}^*$



$\frac{V_{tb} V_{ts}^*}{V_{tb}^* V_{ts}}$

$V_{cb}^* V_{cs}$

$\omega = \text{wrong tagging}$

$$\phi_{CP} = 2\beta_S(+\pi) \approx 2^\circ (SM)$$

	LEP	BaBar	LHCb
$\epsilon(1 - 2\omega)^2$	25-30%	30%	6%




$\beta_S = \arg\left(-\frac{V_{tb}^* V_{ts}}{V_{cb}^* V_{cs}}\right)$  : CP violation with  $B_S \rightarrow J/\psi\phi \rightarrow \mu^+ \mu^- K^+ K^-$

CKM:  $\beta_S \approx 1^\circ$

PDG:  $\beta_S = (0.60 \pm 0.89)^\circ$

$\sigma(d_{flight}) \approx 20 \mu m$



However for  $B_S \rightarrow J/\psi\phi$


PDG		
$\Gamma_L/\Gamma$	$0.527 \pm 0.008$	CP = +
$\Gamma_{  }/\Gamma$	$0.228 \pm 0.007$	CP = +
$\Gamma_{\perp}/\Gamma$	$0.245 \pm 0.004$	CP = -

In HQS,  $\Gamma_{||} = \Gamma_{\perp} \Rightarrow \mathcal{A}^{mix} = \mathcal{A}_L^{mix}$

Angular analysis required (tbd)  
 Otherwise additional  $\sin\Phi$  term  
 amplitude dilution by factor  $\sim 0.5$   
 Slightly reduced sensitivity (can be compensated using  $J/\psi \rightarrow e^+ e^-$  and other modes e.g.  $J/\psi\eta$ )

Mean B flight distance  $\approx 3000 \mu m$

Should  $\Gamma_L/\Gamma = 1$




$\delta(\sin\phi_{CKM}) = \delta(\sin 2\beta_S) \approx 1.2 \times 10^{-3} \cong \delta(\beta_S) \approx 3.5^\circ \times 10^{-2} (stat.)$

# Effect of penguins in $B_s \rightarrow J/\psi\phi$

$$\mathcal{I} = (\mathcal{T} + \mathcal{E})^2 \left[ |V_{cs} V_{cb}^*|^2 \frac{V_{tb} V_{ts}^*}{V_{cb} V_{cs}^*} + |V_{tb} V_{ts}^*|^2 \left( \frac{\mathcal{P}_t}{(\mathcal{T} + \mathcal{E})} \right) \right]^2$$

$$\mathcal{P}_t \rightarrow (\mathcal{P}_t - \mathcal{P}_c) \times \left[ 1 + \left( \frac{V_{us}^* V_{ub}}{V_{ts}^* V_{tb}} \right) \frac{\mathcal{P}_u - \mathcal{P}_c}{\mathcal{P}_t - \mathcal{P}_c} \right]$$




Uncertainty on  $\phi_{CKM}$  is dominated by FCC statistical error as long as  $\mathcal{R} = \frac{\mathcal{P}_t}{\mathcal{T}} < \sim 3\%$

Important input from theory is very important

# Mistag and $B_s$ Mixing Measurement with $B_s \rightarrow D_s \pi$

## Golden channel





**Expect  $\sim 14 \cdot 10^6$  evts** with very small Background (mainly combinatorics)

- No direct decay  $B_s \rightarrow D_s^+ \pi^-$ , i.e. Flavour specific decay
- no CP violation in this mode.


$$\Gamma(\bar{B}_s \rightarrow D_s^+ \pi^-) \propto e^{-\Gamma t} [(1 - \omega) \cos^2 \Delta m t / 2 + \omega \sin^2 \Delta m t / 2]$$

$$\Gamma(B_s \rightarrow D_s^+ \pi^-) \propto e^{-\Gamma t} [\omega \cos^2 \Delta m t / 2 + (1 - \omega) \sin^2 \Delta m t / 2]$$



$$\delta(\Delta m_{B_s})_{stat} \approx (5 \times 10^{-4}) 10^{12} \hbar s^{-1} \text{ [LHCb: } (6.0 \times 10^{-3}) 10^{12} \hbar s^{-1} \text{]}$$

$$\delta(\omega)_{stat} = 1.4 \times 10^{-4}$$




**Wrong tagging measured very precisely**

$\alpha_s = \arg\left(-\frac{V_{ub}^* V_{us}}{V_{tb}^* V_{ts}}\right)$ : CP violation with  $B_s \rightarrow D_s^\pm K^\mp \rightarrow \phi \pi^\pm K^\mp \rightarrow K^+ K^- \pi^\pm K^\mp$


Expect  $\sim 10^6$  evts with very small Background

R.A. , I. Dunietz, B. Kayser Z. Phys. C54, 653 (1992)



$V_{cb} V_{us}^*$

$\frac{V_{tb} V_{ts}^*}{V_{tb}^* V_{ts}}$



$V_{ub}^* V_{cs}$

$$\alpha_s = \arg\left(-\frac{V_{ub}^* V_{us}}{V_{tb}^* V_{ts}}\right)$$

$$\phi_{CKM} = -(\alpha_s - \beta_s) (+\pi)$$

$$\begin{aligned} \Gamma(B_s(t) \rightarrow f) &= |<f|B_s>|^2 e^{-\Gamma t} \{ [1 - \omega(1 - \rho^2)] \cos^2 \frac{\Delta mt}{2} \\ &+ [\rho^2 + \omega(1 - \rho^2)] \sin^2 \frac{\Delta mt}{2} \\ &- (1 - 2\omega)\rho \sin \phi_{CP}^+ \sin \Delta mt \} \\ \Gamma(\bar{B}_s(t) \rightarrow f) &= |<f|B_s>|^2 e^{-\Gamma t} \{ [\rho^2 + \omega(1 - \rho^2)] \cos^2 \frac{\Delta mt}{2} \\ &+ [1 - \omega(1 - \rho^2)] \sin^2 \frac{\Delta mt}{2} \\ &+ (1 - 2\omega)\rho \sin \phi_{CP}^+ \sin \Delta mt \} \\ \Gamma(B_s(t) \rightarrow \bar{f}) &= |<f|B_s>|^2 e^{-\Gamma t} \{ [\rho^2 + \omega(1 - \rho^2)] \cos^2 \frac{\Delta mt}{2} \\ &+ [1 - \omega(1 - \rho^2)] \sin^2 \frac{\Delta mt}{2} \\ &- (1 - 2\omega)\rho \sin \phi_{CP}^- \sin \Delta mt \} \\ \Gamma(\bar{B}_s(t) \rightarrow \bar{f}) &= |<f|B_s>|^2 e^{-\Gamma t} \{ [1 - \omega(1 - \rho^2)] \cos^2 \frac{\Delta mt}{2} \\ &+ [\rho^2 + \omega(1 - \rho^2)] \sin^2 \frac{\Delta mt}{2} \\ &+ (1 - 2\omega)\rho \sin \phi_{CP}^- \sin \Delta mt \} \end{aligned}$$

- No penguin pollution
- $\rho = |\lambda_f| = \left| \frac{q\langle f|\bar{B}_s\rangle}{p\langle f|B_s\rangle} \right|$
- There is a strong phase  $\delta$   
 $\phi_{CP}^\pm = \phi_{CKM} \pm \delta$
- 4 time dependent distributions



$\phi_{CKM}$   
(with 2-fold ambiguity)

$$\sin^2 \phi_{CKM} = \frac{1 + \sin \phi_{CP}^+ \sin \phi_{CP}^- \pm \sqrt{(1 - \sin \phi_{CP}^+)^2 (1 - \sin \phi_{CP}^-)^2}}{2}$$

Note:  $\Delta\Gamma_s$  neglected , which helps remove ambiguity



$\alpha_S = \arg\left(-\frac{V_{ub}^* V_{us}}{V_{tb}^* V_{ts}}\right)$ : CP violation with  $B_s \rightarrow D_s^\pm K^\mp \rightarrow \phi \pi^\pm K^\mp \rightarrow K^+ K^- \pi^\pm K^\mp$





Charged final state only

	unit	value
acceptance	%	86
$\sigma(m_{D_s})$	MeV	$\sim 2.1$
$\sigma(m_{B_s})$	MeV	$\sim 5.6$


To be compared to

$\sigma(m_{B_s})_{LHCb} \approx 17 \text{ MeV}$






$$\alpha_S = \arg\left(-\frac{V_{ub}^* V_{us}}{V_{tb}^* V_{ts}}\right): \text{CP violation with } B_S \rightarrow D_S^\pm K^\mp \rightarrow \phi \pi^\pm K^\mp \rightarrow K^+ K^- \pi^\pm K^\mp$$





- Tracking resolution **crucial** to reduce background
- Combinatoric background to be added (but expected to be relatively small)
- A realistic PID (ToF + dE/dx) enough



$\alpha_s = \arg\left(-\frac{V_{ub}^* V_{us}}{V_{tb}^* V_{ts}}\right)$  : Measurement of CP violation with  $B_s \rightarrow D_s K$

PDG:  $\gamma = (71.1^{+4.6}_{-5.3})^\circ$

$\int L dt = 150 \text{ ab}^{-1}$



$\delta(\rho) \approx 3.2 \times 10^{-3} (\text{stat.})$



$\delta(\sin^2 \phi_{CKM}) \approx \delta(\sin^2 \alpha_s) \approx 5 \times 10^{-3} (\text{stat.}) \cong \delta(\alpha_s) \approx 0.4^\circ (\text{stat.})$

Potential statistical gain of factor 4-5 with  $D_s^\pm \rightarrow K^{*0} K^\pm, \phi \rho^\pm, \dots$  but background needs to be studied (see backup)+  
 Additionnal potential gain (another factor  $\sim 2$ ) with  $B_s \rightarrow D_s^{*\pm} K^\mp, D_s^\pm K^{*\mp}, D_s^{*\pm} K^{*\mp}$ , most modes including  $\gamma(s)$

$$\gamma_S = \arg\left(-\frac{V_{cb}^* V_{cs}}{V_{ub}^* V_{us}}\right): \text{Direct CP violation with } B^\pm \rightarrow D^0(\bar{D}^0)K^\pm, (D^0 \rightarrow K^+K^-, K_S\pi^0)$$

well-known method to measure the  $\gamma$  angle of the “usual” UT

Gronau, London ; Gronau , Wyler



With a final state  $f$  that is accessible to both  $D^0$  and  $\bar{D}^0$ : interference, and CPV.

$$\Gamma(B^+ \rightarrow f_{(D)}K^+) \neq \Gamma(B^- \rightarrow f_{(D)}K^-) \Rightarrow \text{Asymmetry } \mathcal{A}_{CP}^\pm$$

$$\phi_{CKM} = \pi + \gamma_S$$

$$\mathcal{A}_{CP}^\pm = \frac{\pm 2\mathcal{R} \sin \Delta \cos \gamma_S}{1 + \mathcal{R}^2 \mp \mathcal{R} \cos \Delta \cos \gamma_S}$$

$$\mathcal{R}^2 = \frac{Br(B^+ \rightarrow D^0 K^+)}{Br(B^+ \rightarrow \bar{D}^0 K^+)}$$

$\mathcal{R}$  already known to 5%,  
can be much improved  
with  $D^0$  semi-leptonic  
decays

$\Delta$  = strong phase difference. PDG:  $-130^\circ \pm 5^\circ$

Combination of  $\mathcal{A}_{CP}^+$  ( $K^+K^-$ ) and  $\mathcal{A}_{CP}^-$  ( $K^0\pi^0$ ) gives  $\Delta$  and  $\gamma_S$  (8-fold ambiguity)

$\gamma_S = \arg\left(-\frac{V_{cb}^* V_{cs}}{V_{ub}^* V_{us}}\right)$ : **Direct CP violation with  $B^\pm \rightarrow D^0(\bar{D}^0)K^\pm, (D^0 \rightarrow K^+K^-, K_S\pi^0)$**



✦  $B^+ \rightarrow (K^+K^-)_D K^+$ : Relatively easy thanks to excellent mass resolution and PID :  
 $\sigma(D^0) \sim 2 \text{ MeV}$  and  $\sigma(B^+) \sim 6 \text{ MeV}$

✦  $B^+ \rightarrow (K_S\pi^0)_D K^+$ : more challenging :  
 ➤ **Displaced pion tracks from  $K_S$  decay** : Up to O(1m) from the IP. Requires a **large enough tracker** **<efficiency>** **> 90% up to 1.5m feasible**  
 ➤ **Excellent photon energy resolution** : Requires **Crystal like Calorimeter**

$$\sigma(K_S) \sim 2.5 \text{ MeV}$$

$$\sigma(B^+) \sim 20 \text{ MeV} \text{ with } \frac{\sigma_{E\gamma}}{E_\gamma} = \frac{3\%}{\sqrt{E}}$$

$$\sigma(B^+) \sim 80 \text{ MeV} \text{ with } \frac{\sigma_{E\gamma}}{E_\gamma} = \frac{15\%}{\sqrt{E}}$$



$\gamma_S = \arg\left(-\frac{V_{cb}^* V_{cs}}{V_{ub}^* V_{us}}\right)$ : **Direct CP violation with  $B^\pm \rightarrow D^0(\bar{D}^0)K^\pm, (D^0 \rightarrow K^+K^-, K_S\pi^0)$**

$$\int L dt = 150 \text{ ab}^{-1}$$



$B^+ \rightarrow \bar{D}^0 K^+ \rightarrow K^+ K^- K^+$	$\sim 5.8 \cdot 10^5$
$B^+ \rightarrow D^0 K^+ \rightarrow K^+ K^- K^+$	$\sim 5.7 \cdot 10^3$
$B^+ \rightarrow \bar{D}^0 K^+ \rightarrow K_S \pi^- K^+$	$\sim 1.2 \cdot 10^6$
$B^+ \rightarrow D^0 K^+ \rightarrow K_S \pi^- K^+$	$\sim 1.2 \cdot 10^4$

Asymmetries are sizable. E.g. with  $\Delta = -130^\circ$  and  $\gamma_S = 108^\circ$  (SM) :

$$\mathcal{A}_{CP}^+(K^+K^-) \approx -15\% \quad \text{and} \quad \mathcal{A}_{CP}^+(K_S\pi^0) \approx 14\%$$

with **expected statistical uncertainties of  $\sim 0.1\%$**  (absolute, accounting for approx. acceptance and efficiencies), which corresponds to  $\sigma_{\gamma_S} \approx 2.8^\circ$

(uncertainty on  $\gamma_S$  depends on the value of  $\Delta$  – ranges between  $< 1^\circ$  to a few  $^\circ$ )


Possible improvements with additional modes, e.g.  $D \rightarrow K_S \eta, B^+ \rightarrow D^0 K^{*+}$

Measurement of  $\gamma_S$  to  $1^\circ - 2^\circ$  within reach.

# Conclusions

- FCC should enable a precise measurement of the three angles of the “flat” (b,s) UT :  
 with  $B_s \rightarrow J/\psi \phi$  ,  $D_s K$  and  $B^\pm \rightarrow D^0(\overline{D}^0)K^\pm$

$\sigma(\gamma_s) \approx \pm 1^\circ$  via  $B^\pm \rightarrow D^0(\overline{D}^0)K^\pm \rightarrow K^+K^-K^\pm, K_s\pi^0K^\pm$   
 with possible improvement with other modes



$\sigma(\beta_s) \approx \pm 0.035^\circ$  via  $B_s \rightarrow J/\psi \phi \rightarrow \mu^+\mu^-K^+K^-$   
 with possible improvement with other modes  
 • 25x better than the current precision

$\sigma(\alpha_s) \approx \pm 0.4^\circ$  via  $B_s \rightarrow D_s^\pm K^\mp \rightarrow K^+K^-\pi^\pm K^\mp$   
 with possible improvement with other modes

- Simple relation between the 3 phases directly measured in these three processes :

$$-\phi_{D_s K} + \phi_{J/\psi \phi} + \phi_{D^0 K} = 0 \pmod{\pi}$$

should hold in the Standard Model.

## Conclusions

➤ These modes are excellent showcase for the determination of constraints on detector

✦ **Excellent tracking and vertexing resolution**,  $\frac{\sigma(p_T)}{p_T^2} \leq 2. \times 10^{-5} \oplus \frac{1.2 \times 10^{-3}}{p_T \sin\Theta}$

✦ **Excellent calorimetry resolution**, ideally  $\frac{\sigma(E)}{E} \lesssim \frac{5 \times 10^{-2}}{\sqrt{E}} \oplus 5 \times 10^{-3}$

Allows to use  
many other  
decay mode !!!

✦ **Excellent PId resolution** ➤  $3 \sigma$   $K/\pi$  separation up to 25 GeV (covers also K tagging),  
Ideally up to 35 GeV

✦ **Excellent Ks reconstruction** ( crucial for many flavour analyses )



**Additional Slides**

In SM , only few other possible diagrams with same CKM element as tree diagram

- ⇒ well defined CKM angle measured
- ⇒ no direct CP violation expected





Figure 5: Exchange (sub-dominant) diagrams for  $\bar{B}_s \rightarrow D_s^+ K^-$

## Expected number of events

$E_{\text{cm}} = 91.2 \text{ GeV}$ and $\int L = 150 \text{ ab}^{-1}$			
$\sigma(e^+e^- \rightarrow Z)$ nb	number of Z	$f(Z \rightarrow \bar{B}_s)$	Number of produced $\bar{B}_s$
$\sim 42.9$	$\sim 6.4 \cdot 10^{12}$	0.0159	$\sim 1 \cdot 10^{11}$
$\bar{B}_s$ decay Mode	Decay Mode	Final State	Number of $\bar{B}_s$ decays
nonCP eigenstates			
$D_s^+ \pi^-$	$D_s^+ \rightarrow \phi \pi$	$K^+ K^- \pi^+ \pi^-$	$\sim 6.9 \cdot 10^6$
$D_s^+ \pi^-$	$D_s^+ \rightarrow \phi \rho$	$K^+ K^- \pi^+ \pi^- \pi^0$	$\sim 12.9 \cdot 10^6$
$D_s^+ K^-$	$D_s^+ \rightarrow \phi \pi$	$K^+ K^- \pi^+ K^-$	$\sim 5.2 \cdot 10^5$
$D_s^+ K^-$	$D_s^+ \rightarrow \phi \rho$	$K^+ K^- \pi^+ K^- \pi^0$	$\sim 9.8 \cdot 10^5$
$D^0 \phi$	$D^0 \rightarrow K \pi$	$K^- \pi^+ K^+ K^-$	$\sim 6.1 \cdot 10^4$
$D^0 \phi$	$D^0 \rightarrow K \rho$	$K^- \pi^+ K^+ K^- \pi^0$	$\sim 1.7 \cdot 10^5$
CP eigenstates			
$J/\psi \phi$	$J/\psi \rightarrow \mu^+ \mu^-$	$\mu^+ \mu^- K^+ K^-$	$\sim 3.2 \cdot 10^6$
$\phi \phi$	$\phi \rightarrow K^+ K^-$	$K^+ K^- K^+ K^-$	$\sim 4.8 \cdot 10^5$

(To be x 2 for  $B_s$ )

# TPC





# Inclusion of « standard and modest » PID (dE/dx and ToF)

Somewhat conservative PID

- Resolution  $\sigma\left(\frac{dE}{dx}\right) = 5\%$
- Resolution  $\sigma(ToF) = 20\text{ps} (\cong 6\text{mm})$
- ToF Detector location : 2m from IP

Probability of  $\pi$  misidentification as K with  $\varepsilon(K)=50\%$





# Inclusion of « improved » dE/dx and ToF




Resolution  $\sigma\left(\frac{dE}{dx}\right) = 4\%$

Resolution  $\sigma(ToF) = 10\text{ps} (\cong 3\text{mm})$

Detector location : 2m from IP



# Energy spectrum of $\gamma$ from $D_s^- \rightarrow \phi \rho^- \rightarrow (K^+ K^-)_\phi (\pi^- \pi^0)_\rho$





# Inclusion of neutrals for $B_S \rightarrow D_S K$ reconstruction

e.g. could potentially increase statistics (x 3) by adding  $D_S^\pm \rightarrow \phi \rho^\pm$   $\frac{D_S^\pm \rightarrow \phi \rho^\pm}{D_S^\pm \rightarrow \phi \pi^\pm} \approx 1.9$

More generally many physics topics (such as flavor physics) would benefit by using neutrals

⇒ Significant advantage compared to LHCb ⇒ constraint on calorimeter and PId

With very good calorimeter resolution (Xtal type)



$$\sigma(D_S^\pm(\phi\pi^\pm)K^\mp) \approx 5.6\text{MeV} \rightarrow \sigma(D_S^\pm(\phi\rho^\pm)K^\mp) \approx 20\text{MeV}$$

⇒ Background  $D_S^\pm(\phi\rho^\pm)\pi^\mp$  huge


⇒ Excellent PId mandatory

Much worse with LAr type Cal.

$$\sigma(D_S^\pm(\phi\rho^\pm)K^\mp) \approx 36.\text{MeV}$$



# Effect of dE/dx and ToF




« Irreducible bkg », only mass resolution can beat it

**Excellent calorimetry (Xtal like) is also mandatory**



Other backgrounds have to be added  
dE/dx + simple ToF probably not enough unless

- beyond state-of-the-art is achieved for dE/dx and ToF
- or addition of a dedicated PID system







# Inclusion of neutrals for $B_s \rightarrow D_s K$ reconstruction


Assuming HGCal like calorimeter with  $\frac{\delta E}{E} = \frac{0.15}{\sqrt{E}} \oplus 0.005$



Xtal type to HGCal Type :  $\sigma(D_s^\pm(\phi\rho^\pm)K^\mp) \approx 20MeV \rightarrow 51MeV$



Candidates / (5 MeV/c<sup>2</sup>)



[LHCb, JHEP 05 (2015) 019]

