

# Constraints on axion-like polarization oscillations in the cosmic microwave background with POLARBEAR

Jake Spisak on behalf of the POLARBEAR Collaboration  
PASCOS: Dark Matter IV Parallel Session  
June 27th, 2023

Based on arxiv <https://arxiv.org/abs/2303.08410>

# Axions (Axion-Like Particles) and Cosmic Birefringence

- Axion defined for this talk: generic light pseudoscalar  $\phi$ 
  - Also sometimes called 'axion-like particle'
  - Masses of interest: ( $m \approx 10^{-19} - 10^{-22}$  eV)

- Model:

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m_\phi^2 \phi^2 - \frac{g_{\phi\gamma}}{4} \phi F_{\mu\nu} \tilde{F}^{\mu\nu}$$

- Key observable: cosmic birefringence
  - Rotates linearly polarized light

$$\beta = \frac{g_{\phi\gamma}}{2} (\phi(\vec{x}_{\text{abs}}, t_{\text{abs}}) - \phi(\vec{x}_{\text{emit}}, t_{\text{emit}}))$$





Image: Yuto Minami

# Axions as Fuzzy Dark Matter

- Ultralight means large de Broglie wavelength: “Fuzzy dark matter”

$$\lambda_{\text{dB}} \equiv \frac{2\pi}{mv} = 0.48 \text{ kpc} \left( \frac{10^{-22} \text{ eV}}{m} \right) \left( \frac{250 \text{ km/s}}{v} \right)$$

- Effects on small scale structure
  - Small halo formation is cut off
  - Smooth density at halo center



Hui, Annu. Rev. Astron. Astrophys. 2021

# Axion Signal in the CMB

- Oscillating classical field description:

$$\phi(\vec{x}, t) = \phi_0(\vec{x}, t) \sin(m_\phi t + \theta(\vec{x}))$$


Oscillation period: days-months

- Cosmic birefringence:

$$\beta = \frac{g_{\phi\gamma}}{2} (\phi(\vec{x}_{\text{abs}}, t_{\text{abs}}) - \phi(\vec{x}_{\text{emit}}, t_{\text{emit}}))$$

- CMB polarization angle rotation: (Federreke et. al., PRD 2019)

$$\beta_{\text{CMB}}(t) = \frac{g_{\phi\gamma}\phi_0}{2} \sin(m_\phi t + \theta)$$



0.41  $\mu\text{K}$  -160 160  $\mu\text{K}$

ESA/Planck

13.8 billion  
years



# Axion Signal in the CMB

- Oscillating classical field description:

$$\phi(\vec{x}, t) = \phi_0(\vec{x}, t) \sin(m_\phi t + \theta(\vec{x}))$$

Oscillation period: days-months

- Cosmic birefringence:


$$\beta = \frac{g_{\phi\gamma}}{2} (\phi(\vec{x}_{\text{abs}}, t_{\text{abs}}) - \phi(\vec{x}_{\text{emit}}, t_{\text{emit}}))$$

Averaged out:  
washout effect

- CMB polarization angle rotation: (Federreke et. al., PRD 2019)

$$\beta_{\text{CMB}}(t) = \frac{g_{\phi\gamma}\phi_0}{2} \sin(m_\phi t + \theta)$$

CMB  
emitted  
over  
10,000's  
of years



ESA/Planck

13.8 billion  
years



# Axion Signal in the CMB

- Oscillating classical field description:

$$\phi(\vec{x}, t) = \phi_0(\vec{x}, t) \sin(m_\phi t + \theta(\vec{x}))$$

Oscillation period: days-months

- Cosmic birefringence:


$$\beta = \frac{g_{\phi\gamma}}{2} (\phi(\vec{x}_{\text{abs}}, t_{\text{abs}}) - \phi(\vec{x}_{\text{emit}}, t_{\text{emit}}))$$

- CMB polarization angle rotation: (Federreke et. al., PRD 2019)

$$\beta_{\text{CMB}}(t) = \frac{g_{\phi\gamma}\phi_0}{2} \sin(m_\phi t + \theta)$$

Sinusoidal rotation effect: **this work**

CMB absorbed  
by POLARBEAR 1



0.41 μK -160 160 μK


ESA/Planck

13.8 billion  
years



# POLARBEAR-1 Observations

- POLARBEAR-1: CMB telescope in Atacama Desert
  - 150 GHz
  - 3.5 arc-min resolution
  - Took data 2012-2016
- Use 2 years of data: 2012-2014
- 3 small patches:
  - 'Observation': staring at 1 patch, up to 8 hours long
- 515 total observations



POLARBEAR sky patches

POLARBEAR  
Ap.J. 2014

# Estimating an Angle for Each Observation

- Under small rotation angle  $\alpha$ , the maps are:

$$E_{\ell m}^{\text{obs}} = E_{\ell m}^{\text{CMB}} - 2\alpha B_{\ell m}^{\text{CMB}}$$


$$B_{\ell m}^{\text{obs}} = 2\alpha E_{\ell m}^{\text{CMB}} + B_{\ell m}^{\text{CMB}}$$

- Correlate single observation B map with coadded E maps to estimate angle

$$C_{\ell, \text{obs}}^{EB} = 2\alpha_{\text{obs}} C_{\ell, \text{CMB}}^{EE} + \mathcal{O}(\alpha^2)$$



Observed Calculate Known





# Estimating an Angle for Each Observation

- Under small rotation angle  $\alpha$ , the maps are:

$$E_{\ell m}^{\text{obs}} = E_{\ell m}^{\text{CMB}} - 2\alpha B_{\ell m}^{\text{CMB}}$$

$$B_{\ell m}^{\text{obs}} = 2\alpha E_{\ell m}^{\text{CMB}} + B_{\ell m}^{\text{CMB}}$$

carries all the info


- Correlate single observation B map with coadded E maps to estimate angle

$$C_{\ell, \text{obs}}^{EB} = 2\alpha_{\text{obs}} C_{\ell, \text{CMB}}^{EE} + \mathcal{O}(\alpha^2)$$




Observed Calculate Known


One observation:  $\alpha = 0.74^\circ$



# Angle Timestream



# Results: No Detection



POLARBEAR 2023

- Test for presence of signal:  
$$\Delta\chi^2 \equiv \chi^2(A = 0) - \chi^2(A^{\text{mle}}, f^{\text{mle}}, \theta^{\text{mle}})$$
- Compare to a simulated distribution
- $\sigma_{\text{PTE}} = 1.7$ : no significant detection
- Place 95% upper confidence limit on sinusoid amplitude  $A_{95}$  across frequency range

# Stochastic Nature of the Axion Field

Recall:

$$A = \frac{g_{\phi\gamma}\phi_0}{2}$$


Axion field  
amplitude at  
telescope

Problem:  $\phi_0$  is unknown

- Model as Rayleigh distribution


$$P(\phi_0) = \frac{2\phi_0}{\phi_{DM}^2} e^{-\frac{\phi_0^2}{\phi_{DM}^2}}$$

$\frac{1}{2}\phi_{DM}^2 m_\phi^2 = 0.3 \text{ GeV/cm}^3$



Local axion field amplitude variation vs.  
coherence time  $\tau_c \gtrsim 6000$  years

# Constraints on Axion-Photon Coupling



- Marginalize over unknown  $\phi_0$  amplitude
- Assuming axion is all the DM: median 95% upper confidence limit

$$g_{\phi\gamma} < (2.4 \times 10^{-11} \text{ GeV}^{-1}) \times \left( \frac{m_{\phi}}{10^{-21} \text{ eV}} \right)$$

- First CMB analysis of this kind to incorporate stochastic effect of local axion field

# Conclusion

- Axion-photon coupling generates cosmic birefringence
- We searched for a sinusoidal oscillation of the CMB polarization angle using the POLARBEAR telescope.
- No signal found: placed constraints on the coupling vs axion mass in the ( $m \approx 10^{-19} - 10^{-22}$  eV) range

BACKUP


# Signal Estimation

- Estimate sinusoidal signal  $A \sin(2\pi f t + \theta)$  using likelihood:

$$\mathcal{L}(A, f, \theta) \propto e^{-\frac{1}{2} \chi^2(A, f, \theta)}$$

- Frequency range:

$$\frac{1}{50} \text{ days}^{-1} \leq f \leq 0.45 \text{ days}^{-1}$$




Large example signal in simulated data



# Largest Systematic Issue: Half Wave Plate Noise

- Half wave plate (HWP): rotates polarization of incoming light
- HWP was rotated in  $11.25^\circ$  increments between observations during 1st year
- Error at each increment:  $\sigma_{\text{HWP}} = 0.56^\circ$
- Causes low-frequency noise
- **Mitigation: minimum frequency used is  $1/50 \text{ days}^{-1}$**



# Null Test Results

$$T_{\text{null}}(f) \equiv \frac{|A_{\text{null}}(f)|^2}{\sigma (\Re(A_{\text{null}}(f)))^2}$$

$$A_{\text{null}}(f) \equiv (A_1^{\text{mle}} e^{i\theta_1^{\text{mle}}} - A_2^{\text{mle}} e^{i\theta_2^{\text{mle}}})(f)$$

TABLE I. The five null test PTE values used in the pass criteria #1.

PTE statistic	Description	PTE
$\max_{t,f} T_{\text{null}}$	Spurious axion signal	0.032
$\sum_{t,f} T_{\text{null}}$	Total chi-square	0.062
$\max_t \sum_f T_{\text{null}}$	Bad test	0.060
$\max_f \sum_t T_{\text{null}}$	Bad frequency	0.246
$\max_t T_{\text{null}}(f=0)$	Mean angle offset	0.192

TABLE II. The three null test PTE values used in the pass criteria #2.

Axion KS Test inputs	Description	Number of inputs	PTE
$\text{PTE}_{f,t}(T_{\text{null}})$	Overall	22380	0.128
$\text{PTE}_f(\sum_t T_{\text{null}})$	Per frequency	1492	0.122
$\text{PTE}_t(\sum_f T_{\text{null}})$	Per test	15	0.190