

Backgrounds from Frenkel defects in direct detection experiments

Wednesday 29 March 2023 19:13 (1 minute)

A promising strategy for direct detection of sub-MeV dark matter is to look for phonon excitations in crystals. The crystal targets used in such experiments are typically not completely pure, and have impurities or defects. Frenkel defect is an example of a point defect where an atom is dislodged from its position and occupies an interstitial position leaving behind a vacancy. These defects can diffuse and recombine to emit energy in the form of phonons, and can potentially create a background for direct detection experiments. We estimate the defect densities produced through thermal excitations as well as radiogenic nuclear recoils. For various defect configurations, we quantify the diffusion and recombination rates for both thermal and quantum tunneling mechanisms. We find that the thermally generated defects are effectively frozen at cryogenic temperatures and cannot diffuse to recombine with each other. The radiogenic defects produced on the surface can be annealed effectively at room temperature for typical defect configurations, but defects produced through radiogenic nuclear recoils in a shielded environment at cryogenic temperatures during the run-time of the experiment can recombine to produce eV-scale events. We estimate this recombination rate for different defect configurations.

Primary author: SHOLAPURKAR, Mukul

Presenter: SHOLAPURKAR, Mukul

Session Classification: Reception and Poster Session in the same room