

Gravitational Wave Pathway for Testable Leptogenesis

Arnab Dasgupta

PITT-PACC, Department of Physics and Astronomy,
University of Pittsburgh



International Conference on Neutrinos and Dark Matter (NuDM-2022)

25-28 September 2022
Sharm El-Sheikh, Egypt- Online

September 28, 2022


Outline

- ① Gravitational Wave from Strongly First-Order Phase transition
- ② Classically scale invariant paradigm
- ③ Matter-Antimatter Asymmetry
 - $U(1)_{B-L}$ conformal Model
- ④ Asymmetry via Mass-Gain mechanism.
- ⑤ Results

Sources for Stochastic GrW

- ① Emission of GrW from Cosmic strings
- ② From the collision of unstable domain walls
- ③ Production of GrW during or after inflation
 - ▶ Due to extra particle production.
 - ▶ Quantum fluctuations.
- ④ From Strongly First-Order Phase transitions.

First-Order Phase transition



$\langle \phi \rangle \neq 0$

● First-order phase transition proceeds through bubble nucleation



● One of the sources of Gravitational Wave is from Bubble Collision

[Kosowski et.al. 1992]

① The other two sources are

▣ Sound Waves of the plasma [Hindmarsh et.al. 2004]

▣ Turbulance of the plasma [Kamionkowski et.al. 1993]

Contribution to the GW

Sound wave contribution

$$\Omega_{\text{GW}} h^2 \approx \underbrace{\Omega_{\phi} h^2}_{\text{Bubble Collision}} + \overbrace{\Omega_{\text{sw}} h^2}^{\text{Sound wave contribution}} + \underbrace{\Omega_{\text{turb}} h^2}_{\text{Contribution from Magnetic Hydrodynamics}}$$

Bubble
Collision

Contribution

from Magnetic Hydrodynamics

Parameters

▣ α is proportional to the change in the trace of the energy-momentum tensor, ΔT_{μ}^{μ} , across the phase transition

$$\alpha = \frac{1}{\rho_{\gamma}^*} \left[\left(\rho_{\nu} \Big|_{\text{false}} - \rho_{\nu} \Big|_{\text{true}} \right) - \frac{I}{4} \left(\frac{\partial \rho_{\nu}}{\partial T} \Big|_{\text{false}} - \frac{\partial \rho_{\nu}}{\partial T} \Big|_{\text{true}} \right) \right]_{T=T_*}$$

↳ temperature-dependent effective potential
↳ energy density of relativistic radiation

T_* := Nucleation/Percolation temperature

▣ β/H_* : Inverse of the duration of the phase transition in units of the Hubble time H_*^{-1} at the time GW production.

$$\frac{\beta}{H_*} = T_* \frac{dS}{dT} \Big|_{T=T_*}$$

In case of strong supercooling (i.e. for $T_n < T_*$)

$$\frac{\beta}{H_*} = \frac{H_n}{H_*} T_n \frac{dS}{dT} \Big|_{T=T_n}$$

Parameters

▣ v_w : Bubble wall velocity

▣ K_i : Fraction of the released vacuum energy that are converted into energy of

$i=b$: Scalar-field gradients

$i=s$: Sound waves

$i=t$: turbulence

K_s & K_t are expressed in terms of K_{kin}

$$K_s = K_{kin} ; K_t = \epsilon K_{kin}$$

\downarrow
0.1

▣ These efficiency depends on type of phase transition

a) Non-runaway phase transitions in a plasma (NP) $\alpha \ll 1$

b) Runaway phase transitions in a plasma (RP) $\alpha \sim \mathcal{O}(10)$

c) Runaway phase transitions in vacuum (RV) $\alpha \rightarrow \infty$

Observables

III The three contribution can be parametrized in a model independent way

$$\Omega_b(f) h^2 = \Omega_b^{\text{peak}} h^2 \mathcal{S}_b(f, f_b) \quad \left| \quad \Omega_b^{\text{peak}} h^2 \approx 1.67 \times 10^{-5} \left(\frac{v_w}{\beta/H_*} \right)^2 \left(\frac{100}{g_*(T_*)} \right)^{1/3} \left(\frac{K_b \alpha}{1+\alpha} \right)^2 \left(\frac{0.11 v_w}{0.42 + v_w^2} \right)$$

$$\Omega_s(f) h^2 = \Omega_s^{\text{peak}} h^2 \mathcal{S}_s(f, f_s) \quad \left| \quad \Omega_s^{\text{peak}} h^2 \approx 2.65 \times 10^{-6} \left(\frac{v_w}{\beta/H_*} \right) \left(\frac{100}{g_*(T_*)} \right)^{1/3} \left(\frac{K_s \alpha}{1+\alpha} \right)^2$$


$$\Omega_t(f) h^2 = \Omega_t^{\text{peak}} h^2 \mathcal{S}_t(f, f_t) \quad \left| \quad \Omega_t^{\text{peak}} h^2 \approx 3.35 \times 10^{-4} \left(\frac{v_w}{\beta/H_*} \right) \left(\frac{100}{g_*(T_*)} \right)^{1/3} \left(\frac{K_t \alpha}{1+\alpha} \right)^{3/2}$$

$$f_b = 1.6 \times 10^{-2} \text{ mHz} \left(\frac{g_*(T_*)}{100} \right)^{1/6} \left(\frac{T_*}{100 \text{ GeV}} \right) \left(\frac{\beta/H_*}{v_w} \right) \left(\frac{0.62 v_w}{1.8 - 0.1 v_w + v_w^2} \right)$$


$$f_s = 1.9 \times 10^{-2} \text{ mHz} \left(\frac{g_*(T_*)}{100} \right)^{1/6} \left(\frac{T_*}{100 \text{ GeV}} \right) \left(\frac{\beta/H_*}{v_w} \right)$$

$$f_t = 2.7 \times 10^{-2} \text{ mHz} \left(\frac{g_*(T_*)}{100} \right)^{1/6} \left(\frac{T_*}{100 \text{ GeV}} \right) \left(\frac{\beta/H_*}{v_w} \right)$$

Amplitude Behavior



Characteristic Temperature



Classical scale Invariant Paradigm

■ The basic zero temperature potential is

$$V_{\text{tree}} = \lambda_H |H|^4 + \lambda |\Phi|^4 - \lambda' |\Phi|^2 |H|^2$$

⇒ The running of the quartic coupling λ breaks the gauge symmetry $\langle \Phi \rangle = \frac{M}{\sqrt{2}}$

■ The total effective potential can be schematically divided into the following form:

$$V_{\text{tot}} = V_{\text{tree}} + V_{\text{CW}} + V_{\text{th}} \quad \left. \vphantom{V_{\text{tot}}} \right\} \text{thermal effective potential}$$

└─┬─┘
one-loop Coleman-Weinberg potential

$$V_0 = V_{\text{tree}} + V_{\text{CW}} = \frac{1}{4} \lambda(t) G^4(t) \phi^4 \quad \circ \quad t = \log(\phi/\mu)$$

$$G = e^{-\int_0^t dt' \gamma(t')} \quad \gamma(t) = -\frac{a_2}{32\pi^2} g_D^2(t)$$

Classical scale Invariant Paradigm

▣ The running of the gauge coupling $g_D(t)$ and quartic coupling $\lambda(t)$ is given as

$$\frac{d\alpha_D(t)}{dt} = \frac{b}{2\pi} \alpha_D^2(t) \quad \alpha_D = \frac{g_D^2}{4\pi}; \quad \alpha_\lambda = \frac{\lambda}{4\pi}$$

$$\frac{d\alpha_\lambda(t)}{dt} = \frac{1}{2\pi} \left(a_1 \alpha_\lambda^2(t) + 8\pi \alpha_\lambda(t) \gamma(t) + a_3 \alpha_D^2(t) \right)$$

▣ Now, taking the renormalization scale μ to be M , the condition

$$\left. \frac{dV}{d\phi} \right|_{\phi=M} = 0 \quad \text{lead us to}$$

$$a_1 \alpha_\lambda(0)^2 + a_3 \alpha_D(0)^2 + 8\pi \alpha_\lambda(0) = 0$$

-①

Classical scale Invariant Paradigm

▣ The running can be solved analytically, the scalar potential can be given by

$$V_0(\phi, t) = \frac{\pi \alpha_\lambda(t)}{\left(1 - \frac{b}{2\pi} \alpha_D(0) t\right)^{a_2/b}} \phi^4$$

where

$$\alpha_D(t) = \frac{\alpha_D(0)}{1 - \frac{b}{2\pi} \alpha_D(0) t}; \quad \alpha_\lambda(t) = \frac{a_2 + b}{2a_1} \alpha_D(t) + \frac{A}{a_1} \alpha_D(t) \tan\left[\frac{A}{b} \ln[\alpha_D(t)/\pi] + C\right]$$

$$A \equiv [a_1 a_3 - (a_1 + b)^2 / 4]^{1/2}$$

Determined from eq (1)

$$V_{tot}(\phi, t) = V_0(\phi, u) + V_T(\phi, T)$$

↳ $u \equiv \log(\Lambda/M)$ where $\Lambda \equiv \text{Max}(\phi, T)$

$$V_T(\phi, T) = V_{th} + V_{daisy}(\phi, T)$$

$$\begin{cases} V_{th} = \sum_i \frac{n_{B_i}}{2\pi^2} T^4 \mathcal{J}_B \left[\frac{M_{B_i}}{T} \right] \\ V_{daisy}(\phi, T) = - \sum_i \frac{g_i T}{12\pi} [M_i^3(\phi, T) - M_i^3(\phi)] \end{cases}$$

Classical scale Invariant Paradigm

■ The action S_3 can be easily evaluated from the effective potential at sufficiently low temperature i.e. $T \ll M$

$$V_{\text{tot}} \approx \frac{g_D^2(t')}{2} T^2 \phi^2 + \frac{\lambda_{\text{eff}}(t')}{4} \phi^4$$

with $\lambda_{\text{eff}}(t') = \frac{4\pi\alpha_\lambda(t')}{\left(1 - \frac{b}{2\pi}\alpha_D(0)t'\right)^{a_2/b}}$ $t' = \ln(T/M)$

$$S = \frac{S_3}{T} - 4 \ln(T/M) \quad ; \quad \frac{S_3}{T} \approx -9.45 \frac{g_D(t')}{\lambda_{\text{eff}}(t')}$$

■ And the decay width is given as:

$$\Gamma(T) \approx M^4 \exp(-S(T)); \quad S(T) \equiv S_3(T)/T - 4 \log(T/M)$$

Classical scale Invariant Paradigm

U(1)_{B-L}

$$b = 12$$

$$a_1 = 10$$

$$a_3 = 48$$

$$a_2 = 24$$

	$SU(2)_L \times U(1)_Y \times U(1)_{B-L}$
N	$(1, 0, -1)$
ϕ	$(1, 0, 2)$

$$V = \lambda_H |H|^4 + \lambda |\phi|^4 - \lambda' |\phi|^2 |H|^2$$


$$\mathcal{L}_Y \supset -\gamma_D H^\dagger L_i N - \gamma_N \phi N N + h.c$$

One of the Problems in the SM

④ The Standard Model does not explain the present asymmetry.

1. The CP violation coming from Jarlskog invariant is of the order 10^{-20} .

2. The experimental lower bound on the Higgs mass implies the transition is not strongly first order.



Baryon Asymmetry of the Universe

- The observed Baryon Asymmetry of the Universe (BAU) is often quoted in terms of baryon to photon ratio

$$\eta_B = \frac{n_B - n_{\bar{B}}}{n_\gamma} = 6.04 \pm 0.08 \times 10^{-10}$$

- The prediction for this ratio from Big Bang Nucleosynthesis (BBN) agrees well with the observed value from Cosmic Microwave Background Radiation (CMBR) measurements (Planck, arXiv:1502.01589).

Kinds of Mechanism in generating Asymmetry

- ③ Baryogenesis from Decay/Scattering
- ④ Baryogenesis from Electroweak Phase Transitions
- ⑤ Spontaneous Baryogenesis
- ⑥ ... (Affleck-Dine, Gravitational Baryogenesis, etc.)

Sakharov's Conditions

① The three basic ingredients necessary to generate a net baryon asymmetry from an initially baryons symmetric Universe (Sakharov 1967):

1) Baryon Number (B) violation $X \rightarrow Y + B$

2) C and CP violation

$$\Gamma(X \rightarrow Y + B) \neq \Gamma(\bar{X} \rightarrow \bar{Y} + \bar{B})$$

3) Deperature from thermal Equilibrium.

Additional Subtleties on asymmetry




• Additional to the Sakharov's conditions one needs to take care of two more important issues.

1. In order to get asymmetry one needs to be sure to have at least 2 β^* coupling in the loop diagram.

2. The 2 β violating coupling should be to the right of the "cut" of the loop.

* this is necessary if the decaying (scattering) particle(s) does not have any other channel. If it does have a channel without the β coupling then one needs only 1 coupling in the loop.

Which Diagrams this Applies?




Leptogenesis in Conformal B-L

[Phys Rev D, 104, 115029]

■ The leptogenesis does not occur $T > T_n$

■ Schematically



■ Subject to sufficiently high Lorentz factor $\gamma_w > M_N / T_n$

Leptogenesis in Conformal B-L

[Phys Rev D, 104, 115029]

Basic Mechanism

In thermal equilibrium prior to the Phase transition (PT)

$$Y_N \equiv \frac{135}{8\pi^4} \zeta(3) \frac{g_N}{g_*}$$

- Now, provided the Lorentz boost of the wall $\gamma_w > M_N/T_n$, the Y_N is maintained across the bubble wall.
- The N 's are then out of equilibrium, massive, and can decay in a CP and B-L violating way

$$\frac{Y_B}{Y_B^{\text{obs}}} = \epsilon_N K_{\text{sph.}} \frac{Y_N}{Y_B^{\text{obs}}} \left(\frac{T_n}{T_{\text{RH}}} \right)^3$$

Leptogenesis in Conformal B-L

[Phys Rev D, 104, 115029]

Basic Mechanism

In thermal equilibrium prior to the Phase transition (PT)

$$Y_N \equiv \frac{135}{8\pi^4} \zeta(3) \frac{g_N}{g_*}$$

- Now, provided the Lorentz boost of the wall $\gamma_w > M_N/T_n$, the Y_N is maintained across the bubble wall.
- The N 's are then out of equilibrium, massive, and can decay in a CP and B-L violating way

$$\frac{Y_B}{Y_B^{\text{obs}}} = \epsilon_N K_{\text{sph.}} \frac{Y_N}{Y_B^{\text{obs}}} \left(\frac{T_n}{T_{\text{RH}}} \right)^3$$

takes care into account the entropy production from reheating following PT

Leptogenesis in Conformal B-L

[Phys Rev D, 104, 115029]

Basic Mechanism

▮ The wash-out from Inverse decay is given as

$$\Gamma_{ID} \approx \frac{3y^2}{16\pi} M_N \left(\frac{M_N}{T_{RH}} \right)^{3/2} \text{Exp} \left[-\frac{M_N}{T_{RH}} \right]$$

which is safely below H provided







$$\frac{M_N}{T_{RH}} \gtrsim \text{Log} \left[\frac{y^2}{8\pi} \frac{M_{Pl}}{T_{RH}} \left(\frac{M_N}{T_{RH}} \right)^{5/2} \right]$$

Results




▣ The plausibility for detecting the Gravitational Wave (GW) is by calculating the signal-to-noise ratio (SNR)

$$\rho = \left[t_{\text{obs}} \int_{f_{\text{min}}}^{f_{\text{max}}} df \left(\frac{\Omega_{\text{signal}}(f)}{\Omega_{\text{noise}}(f)} \right) \right]$$

Results




Results



Results

	α_{B-L}	ν_{B-L}
BP1	0.0072	10^5 GeV
BP2	0.012	10^5 GeV
BP3	0.019	10^5 GeV



Thank You !!