

Early Dark Energy meets massive neutrinos

Friday 9 September 2022 17:10 (10 minutes)

Early dark energy (EDE) alleviates the H_0 tension at the cost of increasing the clustering amplitude and worsening the S_8 discrepancy. Motivated by massive neutrinos' ability to suppress structure, we study their impact on EDE combining Planck and BOSS full-shape clustering data. A Bayesian analysis returns no evidence for a non-zero neutrino mass sum M_ν (< 0.15 , eV at 95% C.L.), with limits driven primarily by shifts in the BAO scale. A frequentist profile likelihood analysis reveals a correlation between M_ν and the EDE fraction f_{EDE} , which keeps H_0 fixed as M_ν increases. Compared to the best-fit baseline EDE model ($M_\nu = 0.06$, eV), a model with $M_\nu = 0.15$, eV maintains the same H_0 (km/s/Mpc) = (70.08, 70.12, respectively) whilst decreasing S_8 = (0.837, 0.831 respectively), whilst still representing a better fit ($\Delta\chi^2 = -3.1$) relative to Λ CDM. Our results indicate that an EDE+ M_ν model can keep the H_0 tension at the same level as baseline EDE while mitigating the enhanced clustering issue.

Presenter: Mr REEVES, Alexander (ETHZ)