



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 3692

Type: Oral not-in-competition (Graduate Student) / Orale non-compétitive (Étudiant(e) du 2e ou 3e cycle)

The importance of ion temperature profile in collisional sheath modelling

Tuesday 20 June 2023 16:15 (15 minutes)

Plasma discharges contains two distinct zones having different physical properties, namely the quasi-neutral bulk plasma and the sheath where the quasi-neutrality does not hold, separated by an intermediate transition zone called pre-sheath [1]. In particular, the sheath has a strong impact on the entire gas discharge since it is where the plasma interacts with the boundaries. The plasma-sheath transition is still a subject of active research today mainly due to its complex structure [2,3]. The modelling of an entire plasma discharge including the dynamics of sheaths is crucial to understand the behaviour of different plasmas such as nano-particle creation in sputtering magnetron discharge [4], also observed in the coldest region of tokamaks [5].

In this context a new and reliable numerical model for low-temperature plasma discharges including the sheaths is currently under development. Although kinetic models and Particle-In-Cell (PIC) methods [6, 7] are often preferred for their fidelity, they are limited by numerical constraints on the simulation time and memory requirements due to the high number of macro-particles necessary to accurately simulate high density plasmas and the sheaths.

Fluid approaches are limited by the model accuracy itself, but are less demanding in computational resources and is still capable of giving insights of the main physical phenomenon.

In this work, we focus on 1D plasma fluid model adapted for the simulation of medium to high pressure (10^{-1} - 10^2 Pa) direct-current (DC) argon discharges.

In particular, a non quasi-neutral drift-diffusion model of two charged species –ions and electrons was developed aiming at correctly modelling the sheaths. The results are compared with PIC simulation outputs. Our results emphasize for the first time the importance of the ion temperature profile when their collisionality in the sheaths is not negligible.

References

- [1] Langmuir I, Proceedings of the National Academy of Sciences 14 627–637 (1928)
- [2] Hershkowitz N, Physics of Plasmas 12 055502 (2005)
- [3] Riemann K, Plasma Sources Science and Technology 18 014006 (2008)
- [4] Arnas C et al., Phys. Plasmas 26, 053706 (2019)
- [5] Arnas C et al., Plasma Physics and Controlled Fusion 52 124007 (2010)
- [6] Hagelaar G et al., Journal of Applied Physics 93, 67 (2003)
- [7] Sahu R et al., Phys. Plasmas 27, 113505 (2020)

Keyword-1

plasma

Keyword-2

fluid model

Keyword-3

sheath

Primary author: Mr MUN, Jong Hern (University of Saskatchewan)

Co-authors: Mrs CECILE, Arnas (Aix-Marseille University); COUEDEL, Lenaic; Mrs MURAGLIA, Magali (Aix-Marseille University); Mr AGULLO, Olivier (Aix-Marseille University)

Presenter: Mr MUN, Jong Hern (University of Saskatchewan)

Session Classification: (DPP) T4-2 Plasma Physics Symposium VI | Symposium de physique des plasmas VI (DPP)

Track Classification: Symposia Day (Tues. June 20) / Journée de symposiums (mardi, le 20 juin): Symposia Day (DPP - DPP) - Plasma Physics | Physique des plasmas