



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 3830 Type: Oral Competition (Graduate Student) / Compétition orale (Étudiant(e) du 2e ou 3e cycle)

## (G\*) Laser Based Studies of Iron Monohydride (FeH): New Electronic States and Zeeman Spectroscopy in Support of Exoplanet Research

Monday 19 June 2023 16:45 (15 minutes)

We will report on the development of efforts to create new spectroscopic reference data to help astronomers to find exoplanets. Astronomers routinely observe spectra of the molecule iron hydride (FeH) in the atmosphere of M-class stars. By measuring the Doppler shifts of transitions in FeH, they can determine a star's radial velocity. If a star has an exoplanet, the star and the exoplanet will orbit their common center of mass inducing a periodic frequency wobble in the star's FeH spectrum. However, M-class stars often have strong magnetic fields that modify its FeH spectrum through Zeeman splitting. While the infrared transition of interest ( $E^4\Pi$ - $A^4\Pi$ ,  $\sim 1600$  nm) has been studied in detail, its response under magnetic field has not been studied in the lab, although it has been observed in sunspots (A. Asensio Ramos et al., 2004). Hence astronomers require an FeH spectrum to interpret their stellar observations when there is non-negligible magnetic field.

We are creating FeH in the lab to study the response of the  $E^4\Pi$  and  $A^4\Pi$  states to controlled magnetic fields. Transitions involving  $E^4\Pi$  and  $A^4\Pi$  are accessed through new higher-lying states that we have recently discovered. One of the new states is accessible from the ground state,  $X^4\Delta$ , via laser excitation spectroscopy in the green region ( $\sim 510$  nm) and fluoresces to both the  $E^4\Pi$  and  $A^4\Pi$  states. Thus, we can obtain information about the infrared transition indirectly through Zeeman spectroscopy of the visible transitions. The second new higher-lying state is assigned as the  $(2)^4\Phi$  state. This state is also accessed from the ground state ( $\sim 515$  nm) and fluoresces very strongly in the red ( $\sim 625$  nm) to the previously observed  $C^4\Phi$  state.

### Keyword-1

Laser spectroscopy

### Keyword-2

Molecular physics

### Keyword-3

Exoplanets

**Primary author:** BLACKMORE, Timothy

**Co-authors:** Dr ADAM, Allan (University of New Brunswick); Dr ROSS, Amanda (CNRS, Université de Lyon); TOKARYK, Dennis (University of New Brunswick); Dr CROZET, Patrick (CNRS, Université de Lyon)

**Presenter:** BLACKMORE, Timothy

**Session Classification:** (DAMOPEC) M3-2 Atomic- and molecular physics - laser spectroscopy | Physique atomique et moléculaire - spectroscopie laser (DPAMPC)

**Track Classification:** Technical Sessions / Sessions techniques: Atomic, Molecular and Optical Physics, Canada / Physique atomique, moléculaire et photonique, Canada (DAMOPEC-DPAMPC)