

Probing A New Regime Of Ultra-dense Gluonic Matter Using High-energy Photons With The CMS Experiment

JiaZhao Lin For The CMS Collaboration

Rice University



Supported by



U.S. DEPARTMENT OF
ENERGY

Office of
Science


The 38th Winter Workshop on Nuclear Dynamics

Feb 5 – 11, 2023

Marriott Puerto Vallarta Resort & Spa




Understand The Gluon That Binds Us All



- DIS results show a seemingly “indefinite rise” in gluon PDF with linear evolution (gluon splitting)



- What is the fate of gluons at extreme densities (small x)?



Ultra-dense Gluonic Matter


- QCD unitarity: Growth of gluon density cannot continue indefinitely!

- Gluons start to overlap and eventually recombine



Saturation

parton density probed (x)




Photon resolution power (Q)


Ultra-dense Gluonic Matter

- QCD unitarity: Growth of gluon density cannot continue indefinitely!

- Gluons start to overlap and eventually recombine



Saturation




Photon resolution power (Q) →

Vector Meson (VM) Photoproduction

- Photoproduced VM cross section at small x can test on the gluon density scaling behavior



- **VM Photoproduction** $\propto (xg(x, Q^2))^2$
at the **LO** in pQCD

Using Photon As The Probe



Ultra-Peripheral Collision (UPC)

- Nuclei “miss” each other ($b > R_A + R_B$)
- Boosted EM field of nuclei are source of photons
- Interactions via photon-photon or photon-nucleus




VM Photoproduction In UPC

- Emission of quasi-real photons by heavy nuclei

- Photons fluctuate to dipoles

- Dipoles scatter with nucleon/nucleus and form Vector Meson (VM)

VM Photoproduction $\propto (xg(x, Q^2))^2$



What we study


\mathbf{J}/ψ

→ Excellent probe of gluon PDFs

J/psi Photoproduction In γp

Eur.Phys.J.C 79 (2019) 5, 402


- Gluons inside a proton:
 - Investigated via γp interactions by HERA and LHC
 - Consistent results between HERA and LHC: power law
- **No clear signs of gluon saturation inside proton**



J/psi Photoproduction In γ Pb

Eur. Phys. J. C (2021) 81:712


- Measurements from ALICE
- The data trend **challenge** all existing theoretical models
 - Shadowing models
 - Saturation models
- Natural questions:
 - 1. Data in the gap region?
 - 2. If data are correct, what's the underlying physics?



J/psi Photoproduction In γ Pb


Eur. Phys. J. C (2021) 81:712

- Symmetric: both can serve as the photon source and the target
- Each data point: low energy photon + high energy photon contribution



$$\frac{d\sigma_{AA \rightarrow AA' J/\psi}}{dy} = N_{\gamma/A}(y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(y) + N_{\gamma/A}(-y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(-y)$$

No measurement for $\sigma(W)$ at $x \sim 10^{-5}$ in Pb



No easy access to $x \sim 10^{-5}$


Compact Muon Solenoid (CMS)

CMS DETECTOR


Total weight : 14,000 tonnes
Overall diameter : 15.0 m
Overall length : 28.7 m
Magnetic field : 3.8 T

- Located at Large Hadron Collider (LHC) in Geneva

- Equipped with state-of-the-art tracking detectors, calorimeters, and muon detectors



UPC J/psi Photoproduction Event



Clean!




- **UPC collisions produce few particles:**
 - **Require low energy measured in forward calorimeter to suppress hadronic collisions.**
 - **Select events with exactly two reconstructed tracks identified as muons.**

A Solution To The Two-way Ambiguity Puzzle


Guzey et al., EPJC 74 (2014) 2942

Control the impact parameter of UPCs via forward emitted neutrons



Nucleus excitation probability:

$$P_i(b) \propto 1/b^2$$






Spencer Klein & PAS, Ann Rev Nucl Part Sci Vol. 70:323-354

• Analogous to centrality:

$$\bullet \quad b_{XnXn} < b_{0nXn} < b_{0n0n}$$

Determining Neutron Multiplicity

CMS-PAS-HIN-22-002



• Neutron Classes \rightarrow Different impact parameters b

- XnXn \rightarrow smaller b
- 0n0n \rightarrow larger b

— Total fit
 - - - 1n
 - - - 2n
 - - - 3n

Entering A New Regime Of Small x Gluonic Matter

What is measured

Photon flux from theory

What we want

Dominant b ranges of different neutron classes:

- 0n0n: $b > 40$ fm
- 0nXn: $b \sim 20$ fm
- XnXn: $b < 15$ fm

$$\frac{d\sigma_{AA \rightarrow AAJ/\psi}^{0n0n}}{dy} = N_{\gamma/A}^{0n0n}(y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(y) + N_{\gamma/A}^{0n0n}(-y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(-y)$$

$$\frac{d\sigma_{AA \rightarrow AA'J/\psi}^{0nXn}}{dy} = N_{\gamma/A}^{0nXn}(y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(y) + N_{\gamma/A}^{0nXn}(-y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(-y)$$



$$\frac{d\sigma_{AA \rightarrow A'A'J/\psi}^{XnXn}}{dy} = N_{\gamma/A}^{XnXn}(y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(y) + N_{\gamma/A}^{XnXn}(-y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(-y)$$

→ Solve for $\sigma_{\gamma A \rightarrow J/\psi A'}(y)$ and $\sigma_{\gamma A \rightarrow J/\psi A'}(-y)$, and $x = \left(\frac{M_{VM}}{\sqrt{s_{NN}}}\right) e^{\mp y}$

Entering a new regime of small $x \sim 10^{-4} - 10^{-5}$ in nuclei!

Signal Extraction

CMS-PAS-HIN-22-002



Signal yields are extracted by fitting the mass and transverse momentum spectra


AnAn: All possible neutron emissions

Total Coh. J/psi Cross Section

CMS-PAS-HIN-22-002

ALICE, EPJC 81 (2021) 712
LHCb, arXiv:2206.08221

$$\frac{d\sigma_{J/\psi}^{coh}}{dy} = \frac{N(J/\psi)}{(1 + f_I + f_D) \cdot \epsilon(J/\psi) \cdot Acc(J/\psi) \cdot BR(J/\psi \rightarrow \mu\mu) \cdot L_{int} \cdot \Delta y}$$



AnAn: All possible neutron emissions


- A tension between ALICE forward and LHCb forward?

- LHCb new data seems to better connect to ALICE mid rapidity data.

Total Coh. J/psi Cross Section

CMS-PAS-HIN-22-002

ALICE, EPJC 81 (2021) 712
LHCb, arXiv:2206.08221







AnAn: All possible neutron emissions

$$\frac{d\sigma_{J/\psi}^{coh}}{dy} = \frac{N(J/\psi)}{(1 + f_I + f_D) \cdot \epsilon(J/\psi) \cdot Acc(J/\psi) \cdot BR(J/\psi \rightarrow \mu\mu) \cdot L_{int} \cdot \Delta y}$$

- A tension between ALICE forward and LHCb forward?
- LHCb new data seems to better connect to ALICE mid rapidity data.
- **CMS data cover a unique rapidity region and follows ALICE forward rapidity trend**

Coh. J/ψ In Neutron Configurations

CMS-PAS-HIN-22-002



- **First separation in different neutron classes**

- LTA models cannot well describe data in different neutron classes

→ A deeper look at J/Ψ production from γ +Pb at a given W without the “two-way ambiguity” may tell more.


$$\frac{d\sigma_{AA \rightarrow AA' J/\psi}}{dy} = N_{\gamma/A}(y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(y) + N_{\gamma/A}(-y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(-y)$$

High x

Low x

Result: $\sigma_{\gamma A \rightarrow J/\psi A'}(W)$


CMS-PAS-HIN-22-002



- ALICE, LHCb vs. IA:
 - Impulse Approx. (IA): neglects all nuclear effects
 - Data is close to IA at low W.
 - Data is significantly lower than IA at W~125 GeV.
- Larger suppression towards higher W.

Result: $\sigma_{\gamma A \rightarrow J/\psi A'}(W)$


CMS-PAS-HIN-22-002



- ALICE, LHCb vs. IA:
 - Impulse Approx. (IA): neglects all nuclear effects
 - Data is close to IA at low W .
 - Data is significantly lower than IA at $W \sim 125$ GeV.
- Larger suppression towards higher W .

Result: $\sigma_{\gamma A \rightarrow J/\psi A'}(W)$

CMS-PAS-HIN-22-002




- ALICE, LHCb vs. IA:
 - Impulse Approx. (IA): neglects all nuclear effects
 - Data is close to IA at low W.
 - Data is significantly lower than IA at $W \sim 125$ GeV.
- Larger suppression towards higher W.

- **First measurement with CMS**
 - $W < 40$ GeV: rapidly increasing
 - $40 < W < 400$ GeV: slowly raising --
Physics process changed

→ Models considering shadowing effects or gluon saturation effects all fail to describe data trend

Nuclear Suppression Factor


CMS-PAS-HIN-22-002



- $R_g^A = \frac{g_A(x, Q^2)}{A \cdot g_p(x, Q^2)} = \left(\frac{\sigma_{\gamma A \rightarrow J/\psi A}^{exp}}{\sigma_{\gamma A \rightarrow J/\psi A}^{IA}} \right)^{1/2}$
- represents nuclear gluon suppression factor at LO.

Nuclear Suppression Factor

CMS-PAS-HIN-22-002





- $$R_g^A = \frac{g_A(x, Q^2)}{A \cdot g_p(x, Q^2)} = \left(\frac{\sigma_{\gamma A \rightarrow J/\psi A}^{exp}}{\sigma_{\gamma A \rightarrow J/\psi A}^{IA}} \right)^{1/2}$$
 - represents nuclear gluon suppression factor at LO.

- $x \sim 10^{-2} - 10^{-3}$: Flat trend
- Quickly decrease towards lower x region

→ **Beyond model's expectation**

What's The Physics Behind?



CMS-PAS-HIN-22-002



- σ stops rapid rising trend → splitting and recombination of gluons become equal
- **Direct evidence for gluon saturation!!!**

What's The Physics Behind?

CMS-PAS-HIN-22-002



- σ stops rapid rising trend → splitting and recombination of gluons become equal
- **Direct evidence for gluon saturation!!!**

OR


- Nucleus target becomes totally absorptive to incoming photons
- **Black Disk Limit, internal structure invisible!!!**

Another Novel Regime Of QCD: Black Disk Limit

Physics Letters B 537 (2002) 51–61
Phys. Rev. Lett. 87 192301, 2001

In the strong absorption scenario, the interaction probability may reach the unitarity limit. The nucleus target becomes totally absorptive to incoming photons.


- Total cross section of dipole-nucleus interaction $\rightarrow 2\pi R_A^2$



$$\hat{\sigma}_{\text{PQCD}}^{\text{inel}} \leq \hat{\sigma}_{\text{black}} = \pi R_{\text{target}}^2$$

“Black Disk Limit (BDL)”

- opposite to the “color transparency”




The BDL represents a novel regime at small x when the LO QCD and the notion of the parton distributions becomes inapplicable for describing hard processes.

- New theoretical tools are needed in this regime!


Another Novel Regime Of QCD: Black Disk Limit

CMS-PAS-HIN-22-002





- Rapid growth reflects increased gluon density
 - Amplitude of interaction is proportional to gluon density

- Slow growth may suggest the periphery of the nucleus has not become fully “black”




Summary


- For the first time, **directly disentangled coh. $\sigma_{\gamma A \rightarrow J/\psi A'}(W)$** in UPC AA
- CMS measured coh. $\sigma_{\gamma A \rightarrow J/\psi A'}(W)$ to a **new unprecedentedly low-x gluon regime ($10^{-4} - 10^{-5}$)**.
- Flattening of coh. $\sigma_{\gamma A \rightarrow J/\psi A'}(W)$ not predicted by state-of-the-art models
 - **Gluon saturation?** or **black disk limit?** or other physic effects?



The Future



Last updated: January 2022




Exciting opportunities ahead

- Higher luminosities.
- A variety of ion species.
- Upgrades enabled by new technologies!

- **Various VM species in γ Pb with neutron tagging**
- **System size scan with different ion species**

When approaching the BDL

- Coh. cross section scales with $A^{2/3}$
- Incoh. cross section strongly suppressed; internal substructure becomes invisible




Thank You!

EXTRA


Advantages of J/psi in UPC γ Pb

- Higher density, easier to reach the saturation
- Nuclei as target, more nucleons, can lead to nuclear shadowing effects
- Photon flux $\propto Z^2$
- Possibility of multiphoton exchange: Neutron tagging



Problem of Mixing Contributions

A two-way ambiguity! →



Measured J/ψ at y

$$\frac{d\sigma_{AA \rightarrow AA' J/\psi}}{dy} = N_{\gamma/A}(y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(y) + N_{\gamma/A}(-y) \cdot \sigma_{\gamma A \rightarrow J/\psi A'}(-y)$$

What we measured

Photon flux from theory


What we want: VM cross section of one single γ +Pb at each y

Two unknowns but **one** equation!

VM Photoproduction in UPC

• A given $y \rightarrow$ Fixes ω, x, W

- $\omega = \frac{M_{VM}}{2} e^{\pm y}$
 - y : Rapidity of the VM
 - ω : Photon energy
 - M_{VM} : Mass of the VM
- $x = \left(\frac{M_{VM}}{\sqrt{s_{NN}}} \right) e^{\mp y}$
- $W^2 = M_{VM} \sqrt{s_{NN}} \cdot e^{\pm y}$
 - W : Centre-of-mass energy of the photon–target system



Differential Cross Section Calculation

$$\bullet \frac{d\sigma_{J/\psi}^{coh}}{dy} = \frac{N(J/\psi)}{(1+f_I+f_D) \cdot \epsilon(J/\psi) \cdot Acc(J/\psi) \cdot BR(J/\psi \rightarrow \mu\mu) \cdot L_{int} \cdot \Delta y}$$

Incoherent fraction

$$\bullet f_I = \frac{N(InCoh J/\psi)}{N(Coh J/\psi)}$$

Calculated from **pt fit**

Coherent J/ψ yields

- Raw yields within the mass window


Calculated from **mass fit** within $pt < 0.2$ GeV

Feed down ratio

$$\bullet f_D = \frac{N(feed-down J/\psi)}{N(primary J/\psi)}$$

EM Diss. Correction

- Pileup in EM dissociation (EMD): Multiple EMD within the same bunch crossing
- Leads to a decrease in $0n0n$ Events increase in $0nXn$ and $XnXn$ events



EM Diss. Correction

- The correction can be obtained by inverting migration matrix

$$\begin{pmatrix} N^{00} \\ N^{0X} \\ N^{X0} \\ N^{XX} \end{pmatrix}^{obs} = \begin{pmatrix} P_{00}^{00} & 0 & 0 & 0 \\ P_{00}^{0X} & P_{0X}^{0X} & 0 & 0 \\ P_{00}^{X0} & 0 & P_{X0}^{X0} & 0 \\ P_{00}^{XX} & P_{0X}^{XX} & P_{X0}^{XX} & P_{XX}^{XX} \end{pmatrix} \begin{pmatrix} N_{00} \\ N_{0X} \\ N_{X0} \\ N_{XX} \end{pmatrix}^{True}$$


- The matrix element can be obtained from ZB fraction

- $P_{00}^{00} = f_{00}$
- $P_{00}^{0X} = f_{0X}, P_{0X}^{0X} = f_{00} + f_{0X}$
- $P_{00}^{X0} = f_{X0}, P_{X0}^{X0} = f_{00} + f_{X0}$
- $P_{00}^{XX} = f_{XX}, P_{0X}^{XX} = f_{X0} + f_{XX}, P_{X0}^{XX} = f_{0X} + f_{XX}, P_{XX}^{XX} = f_{00} + f_{0X} + f_{X0} + f_{XX} = 1$

Flux From StarLight

arXiv:2111.11383


- The flux of a point-like source with additional cut-off at RA is widely used in phenomenological calculations for UPC processes, such as STARlight.
- This approach is well motivated in photon-nucleus interactions since the flux at impact parameters smaller than the nuclear radius is effectively suppressed by the requirement of no strong interactions between nuclei.




(Color online) Photon fluxes coming from a nucleus in the point-like source approximation and the realistic description as functions of impact parameter b calculated at different photon energies: 100 MeV (a), 100 GeV (b)

Saturation vs Shadowing

- Both relate to the same concept: density of gluons in nPDF at small-x is **reduced** wrt the simple addition of the gluon PDF
- Saturation: Dynamical description via gluon self-interactions that tame the growth of gluon \rightarrow CGC
- Nuclear shadowing: Gribov-Glauber model of multiple scatterings \rightarrow LTA



Gluon saturation



L. Frankfurt, V. Guzey, M. Strikman (Physics Reports 512 (2012) 255-393)

Nuclear shadowing

Theory Description

- Impulse approximation (IA): Photoproduction data from protons, does not include nuclear effects except coherence
- STARlight: Photoproduction data from protons + Vector Meson Dominance model, includes multiple scattering but no gluon shadowing
- EPS09 LO: parametrization of nuclear shadowing data
- LTA: Leading Twist Approximation of nuclear shadowing
- IIM BG, IPsat, BGK-I: Color dipole approach coupled to the Color Glass Condensate formalism with different assumptions on the dipole-proton scattering amplitude
- GG-HS: Color dipole model with hot spots nucleon structure
- b-BK: Color dipole approach coupled with impact-parameter dependent Balitsky-Kovchegov equation
- JMRT NLO: DGLAP formalism with main NLO contributions included
- CCT: Saturation in an energy dependent hot spot model
- CGC: Color dipole model
- NLO BFKL: BFKL evolution of HERA values