

Contribution ID: 61

Type: Talk

The critical endpoint at large N_c

Wednesday 26 July 2023 11:50 (25 minutes)

The phase diagram of QCD is investigated by varying number of colors N_c within a Polyakov loop quarkmeson chiral model. In particular, our attention is focused on the critical point(s); the critical point present for N_c =3 moves toward the mu_q-axis and disappears as soon as the number of color is increased. Yet, a distinct critical point emerges along the temperature axis for N_c =53 and moves toward finite density when increasing N_c further. Thus, the phase diagram at large N_c looks different than the N_c =3 results, since in this case the first-order transition is in the upper-left, while the crossover is in the lower-right regions of the (mu_B-T)plane. The N_c dependent pressure is also evaluated, which shows a scaling with N_c °0 in the confined and chirally broken phase and with N_c °2 in the deconfined one. Moreover, a chirally symmetric but confined "quarkyonic phase" can be seen at large density and moderate temperature with a pressure proportional to N_c .

Primary author: Dr KOVÁCS, Péter (Wigner Research Centre for Physics)
Co-authors: Prof. GIACOSA, Francesco (Kielce University); KOVÁCS, Győző (Wigner RCP)
Presenter: Dr KOVÁCS, Péter (Wigner Research Centre for Physics)
Session Classification: Plenary session