



# 19th International Conference on QCD in Extreme Conditions (XQCD 2023)

Contribution ID: 45

Type: **Poster**

## The QCD chiral phase transition for various numbers of flavors at imaginary baryon chemical potential

*Wednesday, 26 July 2023 19:35 (25 minutes)*

In order to constrain the QCD phase diagram with physical quark masses, the QCD chiral phase transition in the massless limit is investigated, although this is a challenging problem for lattice QCD. In 1984, Pisarski and Wilczek predicted a first-order transition for  $N_f \geq 3$ , based on RG investigations of a linear sigma model in three dimensions, which was supported by lattice QCD simulations on coarse lattices. However, recent lattice QCD results from our group provide strong evidence for a second order chiral phase transition for  $N_f = 2 - 6$  massless quark flavors. It was demonstrated that the first-order chiral transitions, observed on coarse lattices, terminate at a tricritical lattice spacing, and are thus not connected to the continuum chiral limit. As a consequence, the chiral transition in the continuum is of second order, as it is always approached from a crossover region. Adopting the same strategy, we investigate the nature of the chiral phase transition as a function of the number of quark flavors and the lattice spacing for a fixed imaginary baryon chemical potential. We find that first-order transitions, observed on coarse lattices, disappear towards the continuum limit, which coincides with the situation at zero density.

**Primary authors:** D'AMBROSIO, Alfredo; KAISER, Reinhold; PHILIPSEN, Owe

**Presenter:** KAISER, Reinhold

**Session Classification:** Poster session