

Thermal regularization of t -channel singularities of $2 \rightarrow 2$ scatterings in the early Universe

Michał Iglicki

University of Warsaw



based on

B. Grządkowski, M. Iglicki, S. Mrówczyński, [Nucl.Phys.B 984 \(2022\) 115967](#)
M. Iglicki, [JHEP 06 \(2023\) 006](#)

TAUP 2023
Vienna, 29 August 2023

Motivation: the Boltzmann equation

- How much dark matter (or anything else) is there?

$$\begin{aligned}\dot{n}_x + 3Hn_x = & - \sum_{2 \rightarrow F} C_{ij \rightarrow f_1 \dots f_F}^x \langle \sigma v \rangle_{ij \rightarrow f_1 \dots f_F} \left[n_i n_j - \bar{n}_i \bar{n}_j \frac{n_{f_1} \dots n_{f_F}}{\bar{n}_{f_1} \dots \bar{n}_{f_F}} \right] \\ & - \sum_{1 \rightarrow F} C_{i \rightarrow f_1 \dots f_F}^x \langle \Gamma \rangle_{i \rightarrow f_1 \dots f_F} \left[n_i - \bar{n}_i \frac{n_{f_1} \dots n_{f_F}}{\bar{n}_{f_1} \dots \bar{n}_{f_F}} \right]\end{aligned}$$

where

n, \bar{n}

— number density and equilibrium number density

H

— Hubble parameter

$C_{i(j) \rightarrow f_1 \dots f_F}^x$


— combinatoric factors

$\langle \sigma v \rangle$

— thermally averaged cross section

$\langle \Gamma \rangle$

— thermally averaged decay width



} (see the backup slides for details)


Motivation: the Boltzmann equation

- How much dark matter (or anything else) is there?


$$\dot{n}_x + 3H n_x = - \sum_{2 \rightarrow F}^{\infty} C_{ij \rightarrow f_1 \dots f_F}^x \cancel{\langle \sigma v \rangle_{ij \rightarrow f_1 \dots f_F}} \left[n_i n_j - \bar{n}_i \bar{n}_j \frac{n_{f_1} \dots n_{f_F}}{\bar{n}_{f_1} \dots \bar{n}_{f_F}} \right]$$
$$- \sum_{1 \rightarrow F} C_{i \rightarrow f_1 \dots f_F}^x \langle \Gamma \rangle_{i \rightarrow f_1 \dots f_F} \left[n_i - \bar{n}_i \frac{n_{f_1} \dots n_{f_F}}{\bar{n}_{f_1} \dots \bar{n}_{f_F}} \right]$$

- problem: in multi-component DM scenarios
 $\langle \sigma v \rangle$ is sometimes singular!

example:




[GeV]				
m_X	m_{ψ_+}	m_{ψ_-}	m_{h_2}	$\langle \sigma v \rangle$
90	130	30	160	finite
70	130	30	160	singular
70	130	80	160	finite



numerical errors?
physics, actually!

(VFDM model: A. Ahmed et al., Eur.Phys.J.C 78 (2018) 11, 905)

t -channel singularity: definition




$$\mathcal{M} \sim \frac{1}{\textcolor{red}{t} - M^2}, \quad \textcolor{red}{t} \equiv p^2$$

$\textcolor{red}{t} = M^2 \Rightarrow$ **singular** matrix element
 \Rightarrow **infinite** cross section


- IR regularization not applicable if $M > 0$
 - Dyson resummation not helpful if $\Gamma = 0$
- $\left. \begin{array}{l} \\ \end{array} \right\} \Rightarrow$ **massive, stable mediator required**

t -channel singularity: examples


SM: muon colliders




SM: weak Compton scattering



BSM: dark matter in the early Universe



$2 \leftrightarrow 2$ process: when does the t -channel singularity occur?



$$\begin{aligned}\mathcal{M} &\sim \frac{1}{t - M^2} \\ t &\equiv p^2 = (p_1 - p_3)^2 \\ s &\equiv (p_1 + p_2)^2 = (p_3 + p_4)^2\end{aligned}$$

- singularity: $t = M^2$ (massive, stable mediator required)
- cross section
- thermally averaged cross section

$$\sigma(s) \supset \int_{t_{\min}(s)}^{t_{\max}(s)} \frac{dt}{(t - M^2)^2}$$

$$\langle \sigma v \rangle(T) \supset \int \sigma(s) f(E_1, E_2, T) ds$$

- singularity condition

$$t_{\min}(s) < M^2 < t_{\max}(s)$$

$$t_{\min} = m_1^2 + m_3^2 - 2E_1E_3 - 2|\mathbf{p}_1||\mathbf{p}_3| \quad t_{\max} = m_1^2 + m_3^2 - 2E_1E_3 + 2|\mathbf{p}_1||\mathbf{p}_3|$$

$2 \leftrightarrow 2$ process: when does the t -channel singularity occur?

- singularity condition


$$t_{\min}(s) < M^2 < t_{\max}(s)$$

$$t_{\min} = m_1^2 + m_3^2 - 2E_1 E_3 - 2|\mathbf{p}_1||\mathbf{p}_3| \quad t_{\max} = m_1^2 + m_3^2 - 2E_1 E_3 + 2|\mathbf{p}_1||\mathbf{p}_3|$$




- in terms of the CM energy (\sqrt{s})

$$t_{\min}(s) < M^2 < t_{\max}(s)$$

$$\Leftrightarrow s_1 < s < s_2$$



example: weak Compton scattering



$2 \leftrightarrow 2$ process: when does the t -channel singularity occur?

- singularity condition:


$$t_{\min}(s) < M^2 < t_{\max}(s)$$

$$t_{\min} = m_1^2 + m_3^2 - 2E_1 E_3 - 2|\mathbf{p}_1||\mathbf{p}_3| \quad t_{\max} = m_1^2 + m_3^2 - 2E_1 E_3 + 2|\mathbf{p}_1||\mathbf{p}_3|$$

- in terms of the CM energy (\sqrt{s}):

$$t_{\min}(s) < M^2 < t_{\max}(s)$$

$$\Leftrightarrow s_1 < s < s_2$$



- thermally averaged cross section \leftarrow integration over $\sqrt{s} \in [\sqrt{s_{\min}}, \infty)$
(weighted by thermal distribution functions)

- conclusion for the cosmological case:

if $s_2 > s_{\min} \equiv \max\{(m_1 + m_2)^2, (m_3 + m_4)^2\}$,

singularity in the allowed range

$\Rightarrow \langle \sigma v \rangle = \infty$

$$s_2 \equiv \frac{-\beta + \sqrt{\beta^2 - 4\alpha\gamma}}{2\alpha}$$



α, β, γ – functions of m_1, m_2, m_3, m_4 and M

if $s_2 > s_{\min} \equiv \max\{(m_1 + m_2)^2, (m_3 + m_4)^2\}$, singularity in the allowed range

$m_1 > M + m_3$ and $m_4 > M + m_2$

or

$$m_2 > M + m_4 \text{ and } m_3 > M + m_1$$



◊ Coleman-Norton theorem

S. Coleman & R. E. Norton, Nuovo Cim 38, 438-442 (1965)

"It is shown that a Feynman amplitude has singularities on the physical boundary if and only if the relevant Feynman diagram can be interpreted as a picture of an energy- and momentum-conserving process occurring in space-time, with all internal particles real, on the mass shell, and moving forward in time"

note: one of the external states decays,
so it **cannot be an asymptotic state**


Coming back to our example. . .

- condition for the singularity to occur:

$$m_{\psi_+} > m_X + m_{\psi_-}$$

and

$$m_{h_2} > 2m_X$$



$[{\rm GeV}]$						
m_X	m_{ψ_+}	m_{ψ_-}	m_{h_2}	$\langle \sigma v \rangle$	$\psi_+ \rightarrow \psi_- X$	$h_2 \rightarrow XX$
90	130	30	160	finite	allowed	forbidden
70	130	30	160	singular	allowed	allowed
70	130	80	160	finite	forbidden	allowed

Known approaches to the problem

- complex mass of unstable particles



I. Ginzburg, Nucl.Phys.B Proc.Supp. 51 (1996) 85-89

- finite lifetime of the particle should affect the wavefunction
- $p_k \rightarrow \tilde{p}_k \equiv p_k + i p'_k(\Gamma_k)$
- problem: $(\tilde{p}_1 - \tilde{p}_3)^2 \neq (\tilde{p}_4 - \tilde{p}_2)^2 \Rightarrow$ lack of symmetry

- finite beam width

- $n(x, y) \sim e^{-\frac{x^2+y^2}{2a^2}} \neq 1$
⇒ momentum uncertainty
- final results proportional to the width:
 $\int dt |\mathcal{M}|^2 \sim a$
- works for colliders, but inapplicable in the cosmological context

G. L. Kotkin et al., Yad. Fiz. 42 (1982) 692
G. L. Kotkin et al., Int. Journ. Mod. Phys. A 7 (1992) 4707
K. Melnikov & V. G. Serbo, Nucl.Phys. B483 (1997) 67
C. Dams & R. Kleiss, Eur.Phys.J.C29 (2003) 11
C. Dams & R. Kleiss, Eur.Phys.J. C36 (2004) 177



Let us reconsider Dyson resummation...

$$\frac{i\Delta}{p} = \frac{i\Delta^{(0)}}{p} + \frac{i\Delta^{(0)}}{p} \text{---} \textcolor{teal}{i\Pi} \text{---} \frac{i\Delta^{(0)}}{p} + \frac{i\Delta^{(0)}}{p} \text{---} \textcolor{teal}{i\Pi} \text{---} \frac{i\Delta^{(0)}}{p} + \dots$$

assumptions: $|\Pi|$ small, $p^2 \simeq M^2$

scalar: $\frac{1}{p^2 - M^2} \rightarrow \frac{1}{p^2 - M^2 + \Pi}$

fermion: $\frac{\not{p} + M}{p^2 - M^2} \rightarrow \frac{\not{p} + M}{p^2 - M^2 + \text{Tr} \left[\frac{\not{p} + M}{2} \Pi \right]}$

vector: $\frac{-g_{\mu\nu} + \frac{p_\mu p_\nu}{M^2}}{p^2 - M^2} \rightarrow \frac{-g_{\mu\nu} + \frac{p_\mu p_\nu}{M^2}}{p^2 - M^2 + \frac{1}{3} \left(-g^{\mu\nu} + \frac{p^\mu p^\nu}{p^2} \right) \Pi_{\mu\nu}}$

\Rightarrow regulator: $\Sigma \equiv \begin{cases} \Im \Pi \\ \Im \left(\text{Tr} \left[\frac{\not{p} + M}{2} \Pi \right] \right) \\ \Im \left[\frac{1}{3} \left(-g^{\mu\nu} + \frac{p^\mu p^\nu}{p^2} \right) \Pi_{\mu\nu} \right] \end{cases}$

problem: $\Sigma \xrightarrow[\text{opt. th.}]{p^2 \rightarrow M^2}$ decay width

\Rightarrow no regularization for a stable mediator, but...

- early Universe = hot gas
- every particle interacts with a thermal medium
- the mean life time cannot be infinite \Rightarrow effective width
- QFT in a thermal medium: Keldysh-Schwinger formalism


similar considerations (and results) by
H.A. Weldon, Phys. Rev. D 28 (1983) 2007



One-loop self-energy

warning: hereafter, m_1 and m_2 are masses of the loop states

- one-loop contribution to mediator's **self-energy**




$$i\Pi(x, y) = i\Delta_1(x, y) \mathcal{A}(y) i\Delta_2(y, x) \mathcal{A}(x)$$

- non-zero imaginary part of the self-energy appears as a result of interactions with the thermal medium of particles

mediator + thermal bath
particle




unstable intermediate state



Calculation of the one-loop self-energy

- one-loop contribution to the self-energy

$$i\Pi(x, y) = i\Delta_1(x, y) \mathcal{A}(y) i\Delta_2(y, x) \mathcal{A}(x)$$



- non-zero imaginary part of the self-energy appears as a result of interactions with the thermal medium of particles

$$\Pi^+(p, T) = \frac{i}{2} \int \frac{d^4 k}{(2\pi)^4} \left[\Delta_1^+(k+p) \mathcal{A} \Delta_2^{\text{sym}}(k, T) \mathcal{A} + \Delta_1^{\text{sym}}(k, T) \mathcal{A} \Delta_2^-(k-p) \mathcal{A} \right]$$

$$\Delta_i^{\text{sym}}(k, T) \equiv \frac{i\pi}{E_i} \left(\delta(E_i - k_0) + \delta(E_i + k_0) \right) \times [2\eta_i f(E_i, T) - 1] \times (\text{numerator})$$

$$\Delta_i^\pm(p) \equiv \frac{(\text{numerator})}{p^2 - m_i^2 \pm i \operatorname{sgn}(p_0) \varepsilon}, \quad E_i \equiv \sqrt{\mathbf{k}^2 + m_i^2},$$

$$f(E_i, T) = (e^{E_i/T} + \eta_i)^{-1}, \quad \eta_i \equiv +1 \text{ for fermions, } -1 \text{ for bosons}$$

after tedious calculations... (**assumption**: $m_1 > m_2 + M$)

$$\Sigma(|\mathbf{p}|, T) \equiv \Im \Pi^+ (|\mathbf{p}|, T)$$

$$= \frac{1}{16\pi} \frac{X_0}{\beta |\mathbf{p}|} \ln \left[1 + \eta_2 \frac{e^{-\beta(b-a)} e^{\beta E_p} (1 - e^{-2\beta a}) (1 - \eta_1 \eta_2 e^{-\beta E_p})}{(1 + \eta_1 e^{-\beta(b-a)}) (1 + \eta_2 e^{-\beta(b+a)} e^{\beta E_p})} \right]$$

$$a \equiv \frac{\lambda(m_1^2, m_2^2, M^2)^{1/2}}{2M^2} |\mathbf{p}|, \quad b \equiv \frac{m_1^2 - m_2^2 + M^2}{2M^2} E_p, \quad E_p \equiv \sqrt{\mathbf{p}^2 + M^2}$$

$$\lambda(m_1^2, m_2^2, M^2) \equiv [m_1^2 - (m_2 + M)^2] [m_1^2 - (m_2 - M)^2]$$

$$\eta_i \equiv +1 \text{ for fermions, } -1 \text{ for bosons, } \quad X_0 = \eta_2 |\mathcal{M}_{\text{dec}}|^2 \times \begin{cases} 1 & \text{scalar} \\ 1/2 & \text{fermion} \\ 1/3 & \text{vector} \end{cases}$$

effective width:

$$\Gamma_{\text{eff}}(|\mathbf{p}|, T) \equiv M^{-1} \Sigma(|\mathbf{p}|, T)$$



Breit-Wigner propagator:

$$\frac{1}{(t - M^2)^2} \rightarrow \frac{1}{(t - M^2)^2 + M^2 \Gamma_{\text{eff}}(|\mathbf{p}|, T)^2}$$

Result discussion: general properties

$$\Sigma(|\mathbf{p}|, T) = \frac{1}{16\pi} \frac{X_0}{\beta|\mathbf{p}|} \ln \left[1 + \eta_2 \frac{e^{-\beta(b-a)} e^{\beta E_p} (1 - e^{-2\beta a}) (1 - \eta_1 \eta_2 e^{-\beta E_p})}{(1 + \eta_1 e^{-\beta(b-a)}) (1 + \eta_2 e^{-\beta(b+a)} e^{\beta E_p})} \right]$$

$$a \equiv \frac{\lambda(m_1^2, m_2^2, M^2)^{1/2}}{2M^2} |\mathbf{p}|, \quad b \equiv \frac{m_1^2 - m_2^2 + M^2}{2M^2} E_p, \quad E_p \equiv \sqrt{\mathbf{p}^2 + M^2}$$

$$\lambda(m_1^2, m_2^2, M^2) \equiv [m_1^2 - (m_2 + M)^2] [m_1^2 - (m_2 - M)^2]$$

$$\eta_i \equiv +1 \text{ for fermions, } -1 \text{ for bosons}, \quad X_0 = \eta_2 |\mathcal{M}|_{\text{dec}}^2 \times \begin{cases} 1 & \text{scalar} \\ 1/2 & \text{fermion} \\ 1/3 & \text{vector} \end{cases}$$

observations:

- $b > a + E_p$, since $E_p > |\mathbf{p}|$ and $m_1^2 - m_2^2 - M^2 > \lambda^{1/2}$
- $a > 0$ and $E_p > 0$
- a , b and E_p do not depend on T
- $\text{sgn}(\text{logarithmic part}) = \eta_2 = \text{sgn}(X_0) \Rightarrow \Sigma > 0$

Result discussion: limiting cases

$$\Sigma(|\mathbf{p}|, T) = \frac{1}{16\pi} \frac{X_0}{\beta|\mathbf{p}|} \ln \left[1 + \eta_2 \frac{e^{-\beta(\mathbf{b}-\mathbf{a})} e^{\beta E_p} (1 - e^{-2\beta a}) (1 - \eta_1 \eta_2 e^{-\beta E_p})}{(1 + \eta_1 e^{-\beta(\mathbf{b}-\mathbf{a})}) (1 + \eta_2 e^{-\beta(\mathbf{b}+\mathbf{a})} e^{\beta E_p})} \right]$$


- $m_1 = m_2 + M$ (no decay)

$$\textcolor{blue}{a} \equiv \frac{\lambda(m_1^2, m_2^2, M^2)^{1/2}}{2M^2} |\mathbf{p}| \rightarrow 0 \quad \Rightarrow \quad \Sigma \rightarrow 0$$

- $\beta \rightarrow \infty$ (zero temperature) or $p \rightarrow \infty$

$$\beta \textcolor{blue}{a}, \beta \textcolor{blue}{b}, \beta \textcolor{blue}{E_p} \rightarrow \infty \quad \Rightarrow \quad \ln[1+\dots] \rightarrow 0 \quad \Rightarrow \quad \Sigma \rightarrow 0$$

- ★ minimal energy E_2 needed to produce particle 1 on-shell increases with $|\mathbf{p}|$
 \Rightarrow **statistical suppression**
 - ★ zero temperature \leftrightarrow no medium: $f(E, T) \rightarrow 0$




- $p \rightarrow 0$ (mediator at rest)

$$\Sigma \rightarrow \frac{X_0}{16\pi M} \frac{\lambda(m_1^2, m_2^2, M^2)^{1/2}}{M^2} \frac{\eta_2 e^{-\beta(b_0 - M)} (1 - \eta_1 \eta_2 e^{-\beta M})}{(1 + \eta_1 e^{-\beta b_0})(1 + \eta_2 e^{-\beta(b_0 - M)})}$$

finite result


$$b_0 \equiv (m_1^2 - m_2^2 + M^2)/(2M) > M$$

Numerical example: effective width




VFDM model: A. Ahmed et al., Eur.Phys.J.C 78 (2018) 11, 905


$$\begin{aligned} m_X &= 70 \text{ GeV} & (\text{upper plot}) \\ m_{\psi_+} &= 130 \text{ GeV} & m_{\psi_-} = 30 \text{ GeV} \\ m_{h_1} &= 125 \text{ GeV} & m_{h_2} = 160 \text{ GeV} \\ g_x &= 0.1 & \sin \alpha = 0.1 \end{aligned}$$



Numerical example: thermally averaged cross section




$$\langle\sigma v\rangle_{12\rightarrow34}(T) = \int d\Phi_1 d\Phi_2 f(E_1, E_2, T) \times \int d\Phi_3 d\Phi_4 |\mathcal{M}|_{\text{dec}}^2 \frac{(2\pi)^4 \delta^{(4)}(p_1 + p_2 - p_3 - p_4)}{(t - M^2)^2 + M^2 \Gamma_{\text{eff}}(|\mathbf{p}|, T)^2}$$
$$d\Phi_i \equiv \frac{d^3 p_i}{(2\pi)^3 2E_i} \quad - \text{phase-space element}$$




Summary

- *t*-channel singularity of $\langle\sigma v\rangle$ occurs if
 - the process can be seen as a sequence of decay and fusion processes



- the mediator is massive and stable
- the singularity is present both in SM and BSM physics
- known approaches are either unsatisfactory or inapplicable




- interactions with the medium result in a non-zero effective width that regularizes the singularity (but the result is still huge)
- the effective width depends on temperature and mediator's momentum (momentum transfer) and behaves in an expected, natural way

$$\Gamma_{\text{eff}} = \Gamma_{\text{eff}}(T, |\mathbf{p}|)$$

Summary



- *t*-channel singularity of $\langle\sigma v\rangle$ occurs if
 - the process can be seen as a sequence of decay and fusion processes

thank you!



- the mediator is massive and stable
- the singularity is present both in SM and BSM physics
- known approaches are either unsatisfactory or inapplicable
- interactions with the medium result in a non-zero effective width that regularizes the singularity (but the result is still huge)
- the effective width depends on temperature and mediator's momentum (momentum transfer) and behaves in an expected, natural way

$$\Gamma_{\text{eff}} = \Gamma_{\text{eff}}(T, |\mathbf{p}|)$$



BACKUP SLIDES

Boltzmann equation: details

Kolb & Turner, The Early Universe
 Gondolo & Gelmini, Nucl.Phys.B 360 (1991) 145-179

$$\dot{n}_x + 3H n_x = - \sum_{2 \rightarrow F} C_{ij \rightarrow f_1 \dots f_F}^x \langle \sigma v \rangle_{ij \rightarrow f_1 \dots f_F} \left[n_i n_j - \bar{n}_i \bar{n}_j \frac{n_{f_1} \dots n_{f_F}}{\bar{n}_{f_1} \dots \bar{n}_{f_F}} \right] \\ - \sum_{1 \rightarrow F} C_{i \rightarrow f_1 \dots f_F}^x \langle \Gamma \rangle_{i \rightarrow f_1 \dots f_F} \left[n_i - \bar{n}_i \frac{n_{f_1} \dots n_{f_F}}{\bar{n}_{f_1} \dots \bar{n}_{f_F}} \right]$$

n, \bar{n} — number density and equilibrium number density

H — Hubble parameter

combinatorial factors:

$$C_{ij \rightarrow f_1 \dots f_F}^x \equiv \frac{\delta_{ix} + \delta_{jx} - \delta_{f_1 x} - \dots - \delta_{f_F x}}{1 + \delta_{ij}}, \quad C_{i \rightarrow f_1 \dots f_F}^x \equiv \delta_{ix} - \delta_{f_1 x} - \dots - \delta_{f_F x}$$

thermally averaged cross section and decay width:

$$\langle \sigma v \rangle_{ij \rightarrow f_1 \dots f_F} = \frac{g_i g_j}{\bar{n}_i \bar{n}_j} \int \frac{d^3 p_i}{(2\pi)^3} \frac{d^3 p_j}{(2\pi)^3} \bar{f}_i(p_i) \bar{f}_j(p_j) v_{ij} \sigma_{ij \rightarrow f_1, f_2, \dots, f_F}$$

$$\langle \Gamma \rangle_{i \rightarrow f_1, f_2, \dots, f_F} = \frac{g_i}{\bar{n}_i} \int \frac{d^3 p_i}{(2\pi)^3} \bar{f}_i(p_i) \frac{m_1}{E_1} \Gamma_{i \rightarrow f_1, f_2, \dots, f_F}$$

$\bar{f}(p)$ — equilibrium distribution function (Bose-Einstein or Fermi-Dirac)

g — internal degrees of freedom

Møller velocity: $v_{ij} \equiv \sqrt{(\mathbf{v}_i - \mathbf{v}_j)^2 - (\mathbf{v}_i \times \mathbf{v}_j)^2} = \frac{[(p_i p_j)^2 - m_i^2 m_j^2]^{1/2}}{E_i E_j}$

Values of s_1, s_2 in terms of masses

- in terms of the CM energy (\sqrt{s}):

$$t_{\min}(\textcolor{red}{s}) < M^2 < t_{\max}(\textcolor{red}{s})$$


$$\Leftrightarrow \quad s_1 < \textcolor{red}{s} < s_2$$

$$s_{1,2} \equiv \frac{-\beta \mp \sqrt{\beta^2 - 4\alpha\gamma}}{2\alpha}$$

$$\alpha \equiv M^2$$

$$\beta \equiv M^4 - M^2(m_1^2 + m_2^2 + m_3^2 + m_4^2) + (m_1^2 - m_3^2)(m_2^2 - m_4^2)$$


$$\gamma \equiv M^2(m_1^2 - m_2^2)(m_3^2 - m_4^2) + (m_1^2 m_4^2 - m_2^2 m_3^2)(m_1^2 - m_2^2 - m_3^2 + m_4^2)$$



Known approaches to the problem

→ complex mass of unstable particles

I. Ginzburg, Nucl.Phys.B Proc.Suppl. 51 (1996) 85-89



idea: finite lifetime should affect the wavefunction

- at rest: $e^{im_1 t} \rightarrow e^{im_1 t} e^{-\Gamma_1 t}$
 $= e^{i\tilde{m}_1 t}, \quad \tilde{m}_1 \equiv m_1 \left(1 + i \frac{\Gamma_1}{m_1}\right)$
- after Lorentz boost: $p_1 \rightarrow \tilde{p}_1 \equiv p_1 \left(1 + i \frac{\Gamma_1}{m_1}\right)$

→ problem: $(\tilde{p}_1 - \tilde{p}_3)^2 \neq (\tilde{p}_4 - \tilde{p}_2)^2 \Rightarrow$ lack of symmetry
(momentum conservation...)

Known approaches to the problem

→ finite beam width

G. L. Kotkin et al., Yad. Fiz. 42 (1982) 692

G. L. Kotkin et al., Int. Journ. Mod. Phys. A 7 (1992) 4707

K. Melnikov & V. G. Serbo, Nucl.Phys. B483 (1997) 67

C. Dams & R. Kleiss, Eur.Phys.J.C29 (2003) 11

C. Dams & R. Kleiss, Eur.Phys.J. C36 (2004) 177

idea: at colliders, the beams have finite size




they should not be treated as plain waves

example:

Gaussian beam moving along z axis

$$n(x, y) \sim e^{-\frac{x^2+y^2}{2a^2}} \quad a - \text{beam width}$$




$$\begin{aligned} \int \frac{dt}{|t - M^2 + i\epsilon|^2} &\rightarrow \int \frac{a^3 e^{-\frac{a^2 \kappa^2}{2}}}{(2\pi)^{3/2}} \frac{d^3 \kappa dt}{(t - M^2 + i\epsilon - \kappa \cdot \mathbf{q})(t - M^2 - i\epsilon + \kappa \cdot \mathbf{q})} \\ &\sim \frac{\pi a}{|\mathbf{q}|}, \quad \mathbf{q} \equiv \left[\frac{E_3}{E_1} \mathbf{p}_1 - \mathbf{p}_3 \right]_{t=M^2} \end{aligned}$$

→ problem: inapplicable in cosmological context

Process of interest in relation to other diagrams

- self-energy cut



- part of a larger diagram

