


# The physics-driven surface background model for XENONnT




Cecilia Ferrari on behalf of the XENON Collaboration

MOTIVATIONS


The XENONnT experiment, searching for WIMP Dark Matter, has already excluded a vast region of its parameters space.

To probe lower WIMP cross sections an increased statistics is demanded.


1. Wait more time (detector now in data-taking)
2. Improve the data selection (extend the fiducial volume FV)



THE PHYSICS INPUTS




Air-born  $^{222}\text{Rn}$  daughters may plate out on Teflon panels used to optically define the TPC volume.




$^{210}\text{Pb}$ , due to its long half life (22y), constitutes a stable background for WIMP searches. To model it we need:

- $^{210}\text{Pb}$  chain energy spectrum (obtained with GEANT4)
- Electron survival probability (obtained with PyCOMes)

Indeed, electrons travelling towards the anode, may be trapped by the Teflon panels, due to the Teflon charge up phenomenon.




THE SPATIAL INPUTS



Thanks to the LXe high stopping-power, the surface background events are clustered in the neighborhood of the TPC borders. However, in terms of contamination level, the Teflon panels differ from one another:


they have been differently exposed to air. This is evident in the events Theta distribution.

By using the data-driven R and Theta distributions and assuming an initial uniform z distribution, it is possible to build a good model for this background. This is fundamental to design a new (R,Theta,Z) FV cut and gain statistics.



THE OBSERVABLE SPACE

DOI:10.1103/PhysRevD.102.072010





The XENONnT experiment features a double-phase Xenon cylindrical TPC. Whenever an event occurs in LXe volume, the deposited energy is split into three different channels:

- Excitation photons produce the S1 signal.
- Ionization electrons extracted in the GXe give S2 signal.
- Heat is not detected.

On the s1-s2 plane, it is possible to discriminate the WIMP signal-like events from the backgrounds. The surface background (green), is characterized by lower S2 signals due to the electrons collection at Teflon panels.


Traditionally, the analysis is performed in the signal corrected space and, to reduce the computational time, only the r spatial information is used.



With flamedisx modelling and inference toolkit, it is instead possible to build a full dimensional model for this background, by exploiting blocks representing the conditional probabilities of the physics processes that build up an event.



By using the explicit likelihood fit it is possible to heavily reduce the computational costs.

THE SURFACE BACKGROUND MODEL



To reproduce the reduced S2 values, the block modeling the electron detection efficiency has to be modified to include the survival probability map and a nuisance parameter p0 that reduces the nominal electron lifetime.

$$\text{electron\_detection\_eff} = \text{extraction\_eff} \exp(-\text{drift\_time} \cdot p0 / e\_lifetime) \cdot \text{survival\_probability}(r,z)$$








By optimizing the model on a data-set of randomly selected 150 events, we get the following best fit values.

- p0:  $20.2 \pm 0.2$
- Single electron gain:  $(4.25 \pm 0.07)$  PE/electrons
- Single electron width:  $(29.1 \pm 1.9)$  PE/electrons
- Photoelectron std:  $(2.3 \pm 1.2)$  PE/PE

From both Goodness of Fit tests and visual matching, we can say the model agrees with the data.

Notice that s1, s2 and z trends are determined by the input Pb210 energy spectrum and the s2 reduction effect, modeled through the electron detection efficiency block.



The lowER searches fiducial volume (FV)

CONCLUSIONS AND OUTLOOK

Thanks to this study, we measured the surface background effective activity inside the XENONnT TPC:

$$A_{\text{Pb210chain}}^{\text{effective}} = (1.97 \pm 0.11) \text{ mBq/m}^2$$

In the near future, we will design an optimized (R,Theta,Z) fiducial volume and refit, with the flamedisx toolkit, the XENONnT SRO data for WIMP searches.

XENON Preliminary

For further information on this study, please contact [cecilia.ferrari@gssi.it](mailto:cecilia.ferrari@gssi.it)



Teflon panels radial position