



Contribution ID: 409

Type: **Parallel talk**

Muon ($g-2$) and Thermal WIMP DM in $U(1)_{L_\mu-L_\tau}$ Models

Tuesday 29 August 2023 16:30 (15 minutes)

$U(1)_{L_\mu-L_\tau} \equiv U(1)_X$ model is anomaly free within the Standard Model (SM) fermion content, and can accommodate the muon ($g-2$) data for $M_Z \sim O(10-100)$ MeV and $g_X \sim (4-8) \times 10^{-4}$. WIMP type thermal dark matter (DM) can be also introduced for $M_Z \sim 2M_{DM}$, if DM pair annihilations into the SM particles occur only through the s-channel Z' exchange. In this work, we show that this tight correlation between M_Z and M_{DM} can be completely evaded both for scalar and fermionic DM, if we include the contributions from dark Higgs boson (H_1). Dark Higgs boson plays a crucial role in DM phenomenology, not only for generation of dark photon mass, but also opening new channels for DM pair annihilations into the final states involving dark Higgs boson, such as dark Higgs pair as well as $Z Z$ through dark Higgs exchange in the s-channel, and co-annihilation into $Z H_1$ in case of inelastic DM. Thus dark Higgs boson will dissect the strong correlation $M_Z \sim 2M_{DM}$, and much wider mass range is allowed for $U(1)_X$ -charged complex scalar and Dirac fermion DM, still explaining the muon ($g-2$). We consider both generic $U(1)_X$ breaking as well as $U(1)_X \rightarrow Z_2$ (and also into Z_3 only for scalar DM case).

Reference : <https://arxiv.org/abs/2204.04889>

Submitted on behalf of a Collaboration?

No

Authors: Dr KIM, Jongkuk (KIAS); Prof. KO, Pyungwon (KIAS (Korea Institute for Advanced Study)); BAEK, Seungwon (Korea Univ.)

Presenter: Dr KIM, Jongkuk (KIAS)

Session Classification: Dark matter and its detection

Track Classification: Dark matter and its detection