



Contribution ID: 559

Type: **Parallel talk**

## Measurement of low-energy Compton and neutron scattering in Si CCDs with single-electron resolution

*Monday 28 August 2023 15:30 (15 minutes)*

DAMIC-M employs skipper charged-coupled devices (CCDs) with detection threshold of just a few ionization charges to search for low-mass dark matter candidates. An important component of the background comes from small-angle Compton scatters of environmental gamma-rays which must thus be characterized down to  $O(10\text{ eV})$  energy. We used an Am-241 source to measure gamma-ray scattering on silicon atomic shell electrons in a skipper CCD with single-electron resolution. The measurement found notable differences between data and theoretical expectations in the L-shell energy region ( $<150\text{ eV}$ ). We also present preliminary data on a nuclear recoil ionization efficiency measurement in Si down to few ionization charges, obtained with a skipper CCD exposed to low-energy neutrons ( $<24\text{ keV}$ ) from a SbBe photoneutron source. Lastly, we report on a novel method under exploration to identify nuclear recoils through the associated production of lattice defects in the silicon.

### Submitted on behalf of a Collaboration?

Yes

**Author:** SMIDA, Radomir (University of Chicago)

**Presenter:** SMIDA, Radomir (University of Chicago)

**Session Classification:** Dark matter and its detection

**Track Classification:** Dark matter and its detection