Spin Transfer to Λ Hyperons at CLAS12

29/Mar./23, Matthew McEneaney, Duke University

matthew.mceneaney@duke.edu

Research supported by:

U.S. DEPARTMENT OF ENERGY | Office of Science

Λ Hyperons

- Λ polarization is easily accessible from the $\Lambda \rightarrow p\pi^-$ channel:

$$\frac{dN}{d\Omega_p} \propto 1 + \alpha P_b D(y) D^{\Lambda}_{LL'} \cos\theta_{pL'}$$

- $D^{\Lambda}_{LL'}$ describes probability for a quark to transfer its polarization to the Λ
- Related to helicity FF G_1^{Λ}

$$G_1^{\Lambda} \propto (\; S_q \rightarrow \; S_\Lambda \; - \; S_\Lambda \leftarrow \; S_q \;)$$





V. Burkert, et al. NIM 2020.

Choice of Λ Polarization Axis

- $\cos\theta_{pL'}$ is the angle between the proton momentum and the Λ polarization axis L'
- Two choices used for L'
  - Along P_Λ
  - Along P_{γ^*}



Signal Decomposition in MC



Event selection:

$p\pi^-$ and scattered e^-

Kinematic cuts:

$Q^2 > 1$ & $W > 2$ & $y < 0.8$

Λ cuts:

$x_F > 0$ & $z_{p\pi^-} < 1$

Signal Fit on MC



Integrate from $1.10 - 1.13\ GeV$

Event selection:
$p\pi^-$ and scattered e^-
Kinematic cuts:
$Q^2 > 1\ \&\ W > 2\ \&\ y < 0.8$
Λ cuts:
$x_F > 0\ \&\ z_{p\pi^-} < 1$

$$f(x;\alpha,n,\mu,\sigma) = N \begin{cases} \exp -\frac{(x-\mu)^2}{2\sigma^2}, & \frac{x-\mu}{\sigma} > \alpha \\ A(B - \frac{x-\mu}{\sigma})^{-n}, & \frac{x-\mu}{\sigma} \leq \alpha \end{cases}$$

Fit function: Gaussian Peak with power law tail (Crystal Ball)

Signal Fit on Data



Integrate from $1.10 - 1.13\ GeV$

Event selection:

$p\pi^-$ and scattered e^-

Kinematic cuts:

$Q^2 > 1\ \&\ W > 2\ \&\ y < 0.8$

Λ cuts:

$x_F > 0\ \&\ z_{p\pi^-} < 1$

$$f(x;\alpha,n,\mu,\sigma) = N \begin{cases} \exp -\frac{(x-\mu)^2}{2\sigma^2}, & \frac{x-\mu}{\sigma} > \alpha \\ A(B - \frac{x-\mu}{\sigma})^{-n}, & \frac{x-\mu}{\sigma} \leq \alpha \end{cases}$$

Fit function: Gaussian Peak with power law tail (Crystal Ball)

$D_{LL'}$ Extraction

- Linear fit to cross-section requires acceptance correction
- Maximum Likelihood (ML) method requires **equal** amounts of positive and negative helicity events

$$D^{\Lambda}_{LL'} = \frac{1}{\alpha \overline{P_b^2}} \cdot \frac{\Sigma_{i=1}^{N_\Lambda} P_{b,i} D(y_i) \cos\theta^i_{pL'}}{\Sigma_{i=1}^{N_\Lambda} D^2(y_i) \cos^2\theta^i_{pL'}}$$

- No acceptance correction needed for ML



Maximum Likelihood Results

$$D_{LL'}^{\Lambda} = \frac{D_{LL'_{sig}} - \epsilon D_{LL'_{bg}}}{1-\epsilon}$$

vs. z





vs. x_F





Note: errors are solely statistical

HERMES Results from:
A. Airapetian, et al. Physical Review D, 74(7), Oct 2006.

Graph Neural Networks (GNNs)

- **Idea**: use GNN to reduce background in invariant mass spectrum on event-by-event basis
- Pass each event as fully-connected, bidirectional graph
- Each particle is a node with its own data: p_T, θ, ϕ, etc.



Graph Neural Networks (GNNs)

- At basic level, function as generalized form of CNNs



Graph Isomorphism Network (GIN)

- Similar to Weisfeiler-Lehman (WL) Test, essentially ensures aggregation is injective
- Compare with basic GNN convolution:

$$h_v^{(k)} = \mathrm{MLP}^{(k)}\left(\left(1+\epsilon^{(k)}\right)\cdot h_v^{(k-1)} + \sum_{u\in\mathcal{N}(v)} h_u^{(k-1)}\right)$$



Graph Isomorphism Network (GIN)

- Aggregation in final layer is across all previous layers/iterations



GIN Evaluation on MC

Signal shape is preserved



Before NN



After NN

83.7% Test accuracy and background is significantly reduced!

GIN Evaluation on Data

- Evaluate on 24k events
- FOM=$N_{sig}/\sqrt{N_{tot}}$ is 65.74 compared to 34.11 without the GIN



Domain Adaptation

- **Problem**: target domain does not match source domain



Data



MC

Domain Adversarial NNs

- Minimizes distinction between real and training data
- Two objectives: classification task and domain discrimination



Domain Adversarial NNs

- Reverse gradient from discriminator loss during backpropagation



arXiv:1505.07818v4

DAGIN Evaluation on MC

Signal shape is preserved



Before NN



After NN

82.9% Test accuracy and background is significantly reduced!

DAGIN Evaluation on Data

- Evaluate on 24k events
- FOM=$N_{sig}/\sqrt{N_{tot}}$ is 42.09 compared to 34.11 without the GIN



NN Output on Data and MC

- Kolmogorov-Smirnov statistic is 0.035 for GIN and 0.261 for DAGIN





Summary and Next Steps

- Preliminary extraction of $\langle D_{LL'}^{\Lambda} \rangle = 0.0618 \pm 0.0963(stat)$ is consistent with previous measurements from HERMES and NOMAD
- Machine learning methods need further validation
- Next steps:
  - Secondary vertex reconstruction method recently implemented
  - Measurement of Λ spontaneous transverse polarization
  - ΛK spin orbit correlations

Thank you!

Longitudinal Spin Transfer

- Polarized electron selects a quark with the opposite spin
- Polarization of quark is transferred to hadrons after fragmentation



Interpretation of $D_{LL'}^{\Lambda}$

- In the naïve quark model, the strange quark carries the Λ spin
- By measuring the light quark spin contribution, we can better understand the spin structure of the Λ



Optimization of NN Cut

- Scan FOM and purity as a function of NN cut