DIS2023: XXX International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 267

Type: Parallel talk

Evolution of initial state fluctuations in the hotspot model

The hotspot model has proven to be an efficient tool to study coherent and incoherent diffraction HERA data by modeling the initial state fluctuations of the gluon density of the proton. The hotspot model in its original form is a non-perturbative model applicable for low momentum transfer and underestimates the incoherent cross section in orders of magnitude when extended for large momentum transfer studies for J/ψ photoproduction at HERA. In lieu of a perturbative approach to study the spatial gluon density fluctuations at large momentum transfers, we present here a model of the evolution of initial state fluctuations in the hotspot model inspired by the DGLAP Parton shower approach. This is reliable as both the Bjorken limit and the incoherent diffraction at large momentum transfer probes the gluon wave function at smaller length scales as we increase the resolution. In addition to the geometrical fluctuations, we have additional sources of fluctuations in our model namely the hotspot width, number, and normalisation fluctuations which leads to a good agreement of our model's prediction with data. In the framework developed, we obtain a good description of both the normalization and shape of coherent and incoherent differential cross sections (t-spectrum) simultaneously.

Submitted on behalf of a Collaboration?

No

Participate in poster competition?

Yes

Primary author: KUMAR, Arjun (Indian Institute of Technology Delhi)
Co-author: TOLL, Tobias (Indian Institute of Technology Delhi)
Presenter: KUMAR, Arjun (Indian Institute of Technology Delhi)
Session Classification: WG2

Track Classification: WG2: Small-x, Diffraction and Vector Mesons