


# Tagged DIS and the EMC Effect at the EIC

Alex Jentsch (BNL), Zhoudunming Tu (BNL),  
Christian Weiss (JLAB)

Deep-Inelastic Scattering Conference


March 27-31, 2023

East Lansing, Michigan




# The EMC Effect

- Discovered by the European Muon Collaboration ~40 years ago.
  - Puzzle: why the dip?
- Still an unanswered question, and one we hope the EIC can aid in answering.
- Established via measurements with **different nuclear targets!**




deuteron

Nuclear effects  
modify nucleon  
structure? How?



Heavier nucleus (A > 2)


$$R_{EMC} = (\sigma_A/A)/(\sigma_D/2)$$



Understanding the origin of the EMC effect and nuclear modifications of prime interest in nuclear physics!

# Tagged DIS as a tool at the EIC

- **Tagged DIS** measurements → “tag” (generally) far-forward **spectators** in final state.
  - Provides more information than inclusive cross sections → information on nucleon configuration.
- **Lots of topics!**
  - Short-range correlations<sup>1</sup>.
  - Gluon distributions in nuclei.
  - Free nucleon structure functions.
  - Nuclear modifications of nucleons in light nuclei.
    - EMC effect, anti-shadowing, etc.




Tagged spectator nucleon momentum → experimental handle on nuclear configurations with free and modified nucleons.

[1] Z. Tu, A. Jentsch, M. Baker, *et al.*, Phys. Lett. B **811**, 135877 (2020)



# Tagged DIS with deuterons



- Spectator kinematics  $\rightarrow$  determines nuclear configuration.
  - Loosely bound configuration – enables extraction of free nucleon structure via pole extrapolation (previous study<sup>2</sup>).
  - Configuration with strongly-interacting nucleons – opens up study of nuclear modifications.
    - Differential study of transition region where nuclear effects manifest!


Tagged DIS on the deuteron enables study of free and modified nuclear structure in a single nucleus!

# The Deuteron – a stand-alone lab for nuclear physics

- Off-shellness in deuterons as a probe of nuclear effects.


Tagged DIS Process:  $e + d \rightarrow e' + X + p' \text{ or } n'$

Low off-shellness




average-size  
more-likely

High off-shellness



small-size  
less-likely

Deuteron: nucleon internal momentum



$$-t'^2 = M_N^2 - (p_d - p_p)^2$$


Virtuality/off-shellness in the deuteron

**Question:** can the EMC effect be controlled via the off-shellness **without altering the nuclear species?**

# Monte Carlo sample for study


General-purpose eA DIS MC generator

<https://eic.github.io/software/beagle.html>




## For e+d collisions:

- BeAGLE<sup>3</sup> → hard nucleon scattering (DIS process)
- Spectator momentum calculated via deuteron spectral function, using parametrization of Ciofi and Simula.
  - ✓ C. Ciofi degli Atti and S. Simula, Phys. Rev. C 53, 1689 (1996)
  - ✓ Same process as in [1].
- BeAGLE MC samples passed through **full detector simulations**, including beam effects!




# Simulating the EMC Effect in BeAGLE



Use EMC effect slope measurements from data with different nuclear targets.

\*Data from J. Seely *et al.* Phys. Rev. Lett. **103**, 202301 (2009)



Linear fit to virtuality dependence → Minimal parametrization:


Frankfurt and Strikman, Nuc. Phys. B **250** (1985)

C. Ciofi *et al.*, Phys. Rev. C **76**, 055206 (2007)

And others...

# Simulating the EMC Effect in BeAGLE


## BeAGLE



Add EMC effect according to the linear parametrization



- Only apply to  $0.3 < x_{bj} < 0.7$
- $Q^2$  independent
- Weight =  $F_2(\text{bound}) / F_2(\text{free})$



Linear fit to virtuality dependence → Minimal parametrization:

Frankfurt and Strikman, Nuc. Phys. B **250** (1985)


C. Ciofi *et al.*, Phys. Rev. C **76**, 055206 (2007)

And others...



# Simulating the EMC Effect in BeAGLE


EMC Weight Distribution,  $0.45 < x_n < 0.55$



## Result → EMC Weight in BeaGLE

- Weight factor simulates the EMC effect from the *virtuality* in the deuteron.
- Applied event-by-event to compare **with and without weight** → enables study of sensitivity to EMC effect in various observables.

# Full Detector Simulations – Tagged Spectators



**Sample of MC events run through GEANT4 to extract acceptances + momentum smearing.**

## Tagged deuteron spectators

- **Neutrons:** reconstructed in ZDC ( $\theta < 5$  mrad acceptance).
- **Protons:** reconstructed in B0 tracker ( $6 < \theta < 20$  mrad) and off-momentum detectors ( $\theta < 5$  mrad).

# The EMC Effect @ the EIC

- Approach:

- Measure deuteron reduced cross-section  $\sigma_D$ , with and without the off-shell effects included.
  - No FSI included.
- Ratio of  $\sigma_D$  **inside and outside the EMC region** (e.g.  $x \sim 0.5$  and  $x \sim 0.2$ )

➤ Quantity allows direct comparison of cross section with and without EMC weight ( $x \sim 0.2$  chosen to avoid anti-shadowing region).


$$\frac{\sigma_D(\alpha_p, p_{T,p}, x_n = 0.5)}{\sigma_D(\alpha_p, p_{T,p}, x_n = 0.2)}$$

# The EMC Effect @ the EIC

5x41 GeV/n Integrated Luminosity  $\sim 25 \text{ fb}^{-1}$

## • Approach:

- Measure deuteron reduced cross-section  $\sigma_D$ , with and without the off-shell effects included.
  - No FSI included.
- Ratio of  $\sigma_D$  **inside and outside the EMC region** (e.g.  $x \sim 0.5$  and  $x \sim 0.2$ )
- Establish required integrated luminosity.
  - Challenging measurement  $\rightarrow$  high- $x$  + low probability nuclear configuration + lower beam energies.
- **Neutron spectator not possible in 5x41 GeV/n due to detector acceptance.**







# The EMC Effect @ the EIC

5x110 GeV/n Integrated Luminosity  $\sim 16 \text{ fb}^{-1}$

- EIC versatility  $\rightarrow$  different beam energy configurations!




- Higher energy configuration (5x110 GeV/n).
- **More favorable detector acceptance  $\rightarrow$  study of proton *and* neutron spectators with same beam configuration.**
- Measurement of same observable with different beam energies/spectator reconstruction enables better understanding of experimental systematics.


# Different nuclear configurations

- EIC kinematic coverage enables broad, differential study of effects.

- Spectator kinematic coverage  $\rightarrow$  varied deuteron nuclear configurations.



Integrate cross section  
over  $p_{T,p}^2$  in each  $\alpha$  bin.



# Summary and Takeaways

- Deuteron can be used as a general tool to study nucleon structure and the onset of modifications via nuclear effects, including the EMC effect.
- EMC effect can be parametrized using the virtuality/off-shellness in lieu of using multiple nuclear species → allows EMC effect to be studied in one collision system.
- EIC far-forward detection capabilities enable broad coverage of spectator kinematics and differential study of various nuclear configurations.
- Large luminosity ( $\sim 10 \text{ fb}^{-1}$ ) needed to acquire necessary statistics at high- $x$  and  $\alpha \neq 1$ .
- Final-state interactions to be included in the coming days.
- Look for the final results of the study on the arXiv very soon!

