

Exotic tetraquarks with some heavy quarks in lattice QCD

talk at *Exotic hadrons in LHCb*, CERN, Genève

Pedro Bicudo

CeFEMA, Departamento de Física,
Instituto Superior Técnico
Universidade de Lisboa, Portugal

bicudo@tecnico.ulisboa.pt


October 18, 2022

Abstract

We review [64] all the different direct and indirect approaches that lattice QCD has been employing to study multiquarks, focusing in the tetraquarks with heavy quarks.

In the new millennium, the interest in tetraquarks exploded with several experimental discoveries of tetraquark resonances with heavy quarks, starting with the Z_c and Z_b and continuing with many more tetraquarks such as the the T_{cc} .

Lattice QCD predicted the T class of tetraquark, but so far it has not yet been able to comprehend the Z class of tetraquarkss.



Introduction: many exotics with some heavy quarks

Static potentials and colour field densities

String flip-flop potentials with static quarks

Potentials with static heavy quarks for the T_{QQ} family

Diquark-antidiquark and meson-meson in a boundstate

Potentials with static heavy quarks $Q\bar{Q}$ for Z_b family

HAL QCD non-static potentials

Search for tetraquark resonances high in the spectrum

Boundstate search with non-relativistic bottom quarks

Scattering study with the Lüscher method for phase shifts

Outlook for tetraquarks with lattice QCD

Introduction: many exotics with some heavy quarks





Figure: New hadrons discovered at the LHC, most of them at LHCb, including baryons, tetraquarks and pentaquarks, plotted as mass versus preprint submission date [48].

Tetraquarks were first proposed by Jaffe within the bag model in 1977 [45, 46]. In the early eighties, Richard [1, 8, 80, 35] and colleagues proposed tetraquarks with some heavy quarks would most likely form boundstates. After many decades, finally multiquark resonances have been confirmed in more than forty different experiments and lattice QCD computations, Fig. 1 shows the discoveries at LHC only [34, 48].



Concerning lattice QCD, we all know it is a discretization of QCD, needing

- a large ensemble of configurations,
- a large volume V ,
- a small lattice spacing a
- and a physical pion mass m_π to approach nature.

Notice it uses the Euclidean time evolution of correlations to compute the energies of observables. For instance, denoting $|O_1(t)\rangle$ the physical quantum state corresponding to the operator $O_1(t)$ at time, it can be decomposed in eigenvectors of the hamiltonian,

$$|O_1(0)\rangle = \sum_i c_{1i} |v_i\rangle \Rightarrow |O_1(t)\rangle = \sum_i c_{1i} e^{-\lambda_i t} |v_i\rangle \quad (1)$$

and the matrix element between two operators at different times is

$$\langle O_2(t) | O_1(0) \rangle = \sum_i c_{2i}^* c_{1i} e^{-\lambda_i t} , \quad (2)$$

clearly for a time long enough the groundstate dominates this matrix element.

We will review the different lattice QCD results on tetraquarks with some heavy quarks, obtained with dynamical quarks for the u , d , s , c quarks, non-relativistic quarks for b quarks and static quarks as an approximation of b quarks.

Static potentials and colour field densities




Figure: The tetraquark (4Q) Wilson loop for the calculation of the 4Q potential V_{4Q} , utilized in Ref. [62].




Figure: Lagrangian density 3D plot for a tetraquark, from Ref. [25], showing a clear tetraquark double-Y flux tube.

The first computations for tetraquarks in lattice QCD used static quarks only, less expensive than dynamical quarks, and can be computed in pure gauge QCD [63, 62] and Alexandrou et al. [3].

The authors find that the potential V_{4Q} is described by the OGE Coulomb plus a four-body potential V_{c4Q} , with a double-Y extension of the three-body linear potential,

$$V_{4Q} = -\alpha_{4Q} \left\{ \left(\frac{1}{r_{12}} + \frac{1}{r_{34}} \right) + \frac{1}{2} \left(\frac{1}{r_{13}} + \frac{1}{r_{14}} + \frac{1}{r_{23}} + \frac{1}{r_{24}} \right) \right\} + \sigma_{4Q} L_{\min} + C_{4Q} \quad (3)$$

The colour electric and magnetic square field densities were studied with four static quarks, by Cardoso et al. [25, 26]. Fig. 3 shows a double-y with two Steiner junctions.

String flip-flop potentials with static quarks




Figure: Triple string flip-flop potential [9].




Figure: Evidence of flip-flop of the flux tube in the colour field densities for the $QQ\bar{Q}\bar{Q}$ system [24].

Depending on the geometry of our system, we can have three different groundstate colour singlets two different $1\bar{1}$ and one $\bar{3}3$. Moreover, the orthogonal colour singlet to the $\bar{3}3$ is the $6\bar{6}$ and the orthogonal colour singlet to the $1\bar{1}$ is the $8\bar{8}$.

In lattice QCD, the dependence of the potential on the positions of four quark systems has been studied. The string flip-flop was already observed in the first study of tetraquark potentials [62]. Then Cardoso et al [24] confirmed this flip-flop studying the flip-flop of the colour field densities, see Fig. 5. Recently the computation, not only of the groundstate potential, but also of the first excited potential was performed [10]. Indeed, the flip-flop was observed both in the groundstate and in the excited potential. The mixing of both potentials in the transition region was also observed.

However, the spin-dependence of the flip-flop potential has not yet been computed, this leads to more bound state tetraquarks of the type T_{bb} than the ones obtained with lattice QCD [9].

Potentials with static heavy quarks for the T_{QQ} family




Figure: The lattice QCD potentials fitted in Ref. [78, 79, 16], left scalar-isoscalar and right vector-isovector.

The first lattice computations for T_{bb} and related states, proposed by Richard many years ago, utilized static potentials computed with static heavy quarks and dynamical light quarks, coupled to two heavy-light mesons, first with quenched light quarks [60, 59, 31, 30] and then for dynamical light quarks [78, 79, 6, 23, 16]. In the operators,

$$(C\Gamma)_{AB} \left(\bar{Q}_C(\mathbf{r}_1) \psi_A^{(1)}(\mathbf{r}_1) \right) \left(\bar{Q}_C(\mathbf{r}_2) \psi_B^{(2)}(\mathbf{r}_2) \right), \quad (4)$$

\bar{Q} denotes a static quark operator, ψ a light antiquark operator, A, B and C are spin indices and $C = \gamma_0 \gamma_2$ is the charge conjugation matrix [78, 79]. More recently heavy quark spin effects were partly included [15], the extrapolation to the chiral limit was performed [16, 12, 13, 15, 11, 14] and the light quarks were extended from the u, d flavours to the s and c flavours [12]. Using an educated ansatz for the potential, the dynamics provided to the heavy quarks in the Schrödinger equation with the Born-Oppenheimer approximation [18] led to the prediction of boundstates.




Figure: The picture of perturbative one-gluon exchange at short distances and of meson wavefunction screening at large distances, shown in Ref. [16].

channel	α	d/a	p	χ^2/dof
scalar isosinglet	0.293(33)	4.51(54)	2.74(1.20)	0.35
vector isotriplet	0.201(77)	2.48(69)	2.0 (fixed)	0.06

Table: χ^2 minimizing fit results of the ansatz (5) to the lattice static antiquark-antiquark potential; fitting range $2 \leq r/a \leq 6$; lattice spacing $a \approx 0.079$ fm

We expect a Coulomb potential at short distances, and the two heavy antiquarks are in the groundstate if they are in a triplet colour state 3, s-wave and thus spin 1. At large distances we expect a screening typical of the static-light wavefunction, sketched in Fig. 7. Our best ansatz is,

$$V(r) = -\frac{\alpha}{r} \exp\left(-\left(\frac{r}{d}\right)^p\right), \quad (5)$$

A binding energy of -90 ± 43 MeV is predicted. Other tetraquarks such as $ls\bar{b}\bar{b}$, $lc\bar{b}\bar{b}$, $sc\bar{b}\bar{b}$, $ll\bar{c}\bar{b}$, $ll\bar{c}\bar{c}$ do not bind in this approach, but a p-wave resonance with the same flavour was also predicted. Partly including the spin of the heavy antiquarks reduces binding slightly to -60 ± 45 MeV, in the binding channel of a BB^* . Importantly, the quantum numbers predicted for the T_{bb} are $I(J^P) = 0(1^+)$. At small distance r , the heavy quarks have spin 1, and the light quarks are in a scalar-isoscalar state.

Potentials with static heavy quarks QQ for Z_b family





Figure: Eigen-energies of $\bar{b}b\bar{d}u$ system, computed in Ref. [71], for various separations r between static quarks b and \bar{b} are shown by points. The eigenstate dominated by $B\bar{B}^*$ (red circles) has energy significantly below $m_B + m_{B^*}$ and shows sizable attraction. Lattice spacing is $a \simeq 0.124$ fm.

The approach of computing, in a first step, potentials in lattice QCD using static quarks for the heavy quarks, has also been applied to the case where the heavy pair is a $Q\bar{Q}$. In the Z_b family of tetraquarks, with flavour $u\bar{d}b\bar{b}$, we have two possible flavours for the meson-meson pairs, either the pair $u\bar{b} = B^{+(*)}$ and $\bar{d}b = \bar{B}^{0(*)}$ or the pair $u\bar{d}$ and $b\bar{b}$, say corresponding to a meson π^+ and a quarkonium meson such as an Υ . Potentials possibly leading to a boundstate in the $B^{+(*)} - \bar{B}^{0(*)}$ channel was found in Refs. [68, 69, 72, 71, 75, 2]. Nevertheless a difficulty resides in identifying the different open coupled channels, because the different channels are not orthogonal.

HAL QCD non-static potentials



The HAL QCD method computes potentials with dynamical quarks [43, 5, 44], extracting them from the wavefunction / Nambu-Bethe-Salpeter amplitude,

$$V(r) = \frac{\Delta\phi(r)}{2m\phi(r)} + E. \quad (6)$$

from operators of two hadrons, each one composed of quarks.

Figure: Coupled channel potentials obtained in Ref [4] with the HAL QCD method using dynamical light quarks and non-relativistic heavy quarks.

Very recently, in Ref. [4] for the T_{bb} study, a 2×2 time dependent coupled channel was studied with pairs of B and B^* mesons. The potentials are shown in Fig. 11, for the 2×2 coupled channel system of a BB^* pair and a B^*B^* pair. They are in general comparable to the potentials obtained with static quarks in the same 2×2 coupled channel system of Ref. [15], where they are denominated V_j and V_5 , with evidence for OGE attraction in the BB^* . Moreover, observing in detail the HAL QCD potentials, there is a new evidence for some OPEP attraction in the B^*B^* pair.

Search for tetraquark resonances high in the spectrum




Figure: The spectrum for quantum numbers $J^G(J^{PC}) = 1^+(1^{+-})$ from Ref. [73]. (a) Position of the experimental Z_c^+ candidates [19]. (b,c) The discrete energy spectrum from our lattice simulation: (b) shows energies based on complete 22×22 matrix of interpolators, (c) is based on the 18×18 correlator matrix without diquark-antidiquark interpolating fields \mathcal{O}_{1-4}^{4q} .

After Z_c was discovered in three different experiments, full dynamical lattice QCD calculations were performed [74, 73, 36, 27]. The authors used the technique of comparing the spectrum just with several meson-meson operators, and the spectrum after adding diquark-antidiquark operators. Representative examples are

$$\mathcal{O}_1^{\psi(0)\pi(0)} = \bar{c}\gamma_i c(0) \bar{d}\gamma_5 u(0) \quad (7)$$

for meson meson operators, and for the diquark-antidiquark operators,

$$\mathcal{O}_1^{4q} \propto \epsilon_{abc}\epsilon_{ab'c'} (\bar{c}_b C \gamma_5 \bar{d}_c c_{b'} \gamma_i C u_{c'} - \bar{c}_b C \gamma_i \bar{d}_c c_{b'} \gamma_5 C u_{c'}) .$$

While this same technique succeeded in identifying the $X(3872)$ state in the spectrum [66], no evidence for the Z_c was found in the spectrum, see for instance Fig. 12

Boundstate search with non-relativistic bottom quarks




Figure: Results of Ref. [47] for the $ud\bar{c}\bar{c}$ and $us\bar{c}\bar{c}$ doubly charm tetraquark states, colour coded. Left panel: chiral extrapolation for several pion masses at $a = 0.1207$ fm for each of the states. Right panel: Continuum extrapolation at the chiral extrapolation to the physical pion mass.

With no static quarks, the T_{bb} fully exotic tetraquark has been studied by Francis et al. [33], Leskovec et al. [51] and Junnarkar et al. [47], using the NRQCD lattice action [76, 50, 58, 7, 22, 29, 52, 53] to calculate bottom quark propagators. All the lattice QCD computations agree that there is a bound T_{bb} tetraquark, and as well in T_{bbs} . There is still some tension in the other quantum numbers of this family of tetraquarks; in general Ref. [47] tends to find binding in T_{cc} , T_{ccs} and T_{bbc} where the other groups don't. The later group employed the chiral extrapolation to the physical m_π , and the continuum limit to vanishing a , only the thermodynamic limit with large volumes remain to be checked. This is illustrated, as in Fig. 13 for the tetraquarks T_{cc} and T_{ccs} . Also notice Hughes et al found a null evidence for a full bottom tetraquark boundstate in Ref. [42].

Scattering study with the Lüscher method for phase shifts




Figure: The center-of-momentum energy $E_{cm} = (E^2 - \vec{P}^2)^{1/2}$ of the $cc\bar{u}\bar{d}$ system [67] in various lattices. The lattice energies are shown by large circles and squares. The free DD^* energies are shown by lines.




Figure: $p \cot \delta_{l=0}^{(J=1)}$ for DD^* scattering (red line) and $ip = +|p|$ (cyan line) versus p^2 . The virtual bound state occurs [67] at the momenta indicated by the magenta octagon, where two curves intersect.

Lüscher et al and others developed a technique to compute phase shifts from the momenta and energy of open channels [56, 57, 56, 55, 54, 20, 65]. For instance in a single channel with two particles, using the effective range approximation [28, 61],

$$k^{2l+1} \cot \delta_l(k) = \frac{-1}{a_l} + \frac{r_l}{2} k^2 + o(k^4) \quad (8)$$

with scattering length a_l and effective range r_l , a pole is equivalent to $\cot \delta_l + i = 0$. The scattering of a pair of pairs of B and B^* mesons was also studied [4, 77], and the T_{bb} boundstate was obtained. However, very recently the T_{cc} was studied in the channel DD^* by Padmanath and Prelovsek [67]. However only a virtual boundstate pole was found, with $\text{Im}(k) < 0$, whereas for a boundstate we should have $\text{Im}(k) > 0$.

Outlook for tetraquarks with lattice QCD

state	qnumber	δ mass	width	decay mode	significance	experiment
$T_{cc}(3874)$	$ud\bar{c}\bar{c} ?$	-360 ± 44 KeV	48 ± 16 KeV	virtual $D^0 D^{*+}$	$15.5 \pm 6.5\sigma$	LHCb [20, 21]
	$ud\bar{c}\bar{c} 1^+$	-23 ± 11 MeV	0	-	-	dynamical lattice QCD [56]
		vBS ~ -9 MeV	0	-	-	dynamical lattice QCD [57]
T_{ccs}	$us\bar{c}\bar{c} 1^+$	-8 ± 8 MeV	0	-	-	dynamical lattice QCD [56]
T_{bc}	$us\bar{b}\bar{c} 1^+, 0^*$	$\sim -40 \pm 50$ MeV	0	-	-	heavy quark lattice QCD [58]
T_{bb}	$ud\bar{b}\bar{b} 1^+$	-90 ± 43 MeV	0	-	-	static lattice QCD [59–63]
		-59 ± 38 MeV	0	-	-	2×2 static lattice QCD [64]
		-189 ± 13 MeV	0	-	-	heavy quark lattice QCD [65]
		~ -113 MeV	0	-	-	heavy quark lattice QCD [66, 67]
		-143 ± 34 MeV	0	-	-	heavy quark lattice QCD [56]
		-128 ± 34 MeV	0	-	-	heavy quark lattice QCD [68]
	$ud\bar{b}\bar{b} 0^*$	~ -120 MeV	0	-	-	heavy quark lattice QCD [69]
		-154.8 ± 37.2 MeV	0	-	-	scattering lattice QCD [70]
		-83.0 ± 30.2 MeV	0	-	-	scattering lattice QCD [70]
		-50.0 ± 5.1 MeV	0	-	-	static lattice QCD [71]
	-5 ± 18 MeV	0	-	-	heavy quark lattice QCD [56]	
T_{bbs}	$us\bar{b}\bar{b}, bs\bar{b}\bar{b} 1^+$	-98 ± 10 MeV	0	-	-	heavy quark lattice QCD [65]
		~ -36 MeV	0	-	-	heavy quark lattice QCD [66, 67]
		-87 ± 32 MeV	0	-	-	heavy quark lattice QCD [56]
		~ -80 MeV	0	-	-	heavy quark lattice QCD [72]
		-86 ± 32 MeV	0	-	-	heavy quark lattice QCD [58]
T_{bbc}	$uc\bar{b}\bar{b} 1^+$	-6 ± 11 MeV	0	-	-	heavy quark lattice QCD [56]
T_{bbcs}	$sc\bar{b}\bar{b} 1^+$	-8 ± 3 MeV	0	-	-	heavy quark lattice QCD [56]

Figure: The tetraquark boundstates, or very narrow resonances [64]. Experimentally only the T_{cc} has been observed, thus we include the lattice QCD predictions as well, where we detail the approach to address the heavy quarks (static, heavy quark effective theory, or dynamical) [47, 67, 77, 16, 12, 13, 11, 14, 15, 32, 33, 41, 51, 40, 4, 23, 70, 77].

The case where lattice has most success is in the boundstates of the T family, such as the T_{bb} which has not yet been observed but is presently used as a benchmark for different lattice QCD studies.




Figure: The pole in the scattering amplitude related to T_{cc} in the complex energy plane: the lattice result [67] (magenta) and the LHCb result (orange).

The T_{cc} recently found at LHCb is an excellent case study for future lattice QCD efforts. The extension of the Lüscher technique to several channels and to three particle resonances is under development by Hansen et al, [37, 38, 39], this should enable the study of the T_{cc} . Another new development is the Master field approach, where a very large lattice is used. The statistical average over several configurations is no longer necessary. In this case the One Pion Exchange Potential should be able to use its full extent, and this is important for deuteron-like molecules. T_{cc} is expected to have both a Dd -like attraction at short distances and a π exchange at large distances.

However, for the Z_b , $Z_c \dots$ family, high in the spectrum, with many decay coupled channels, other new technical advances are still necessary to study them with lattice QCD.

We expect Tetraquarks will become a priority for the lattice QCD community, with increased precision, new techniques, and computations to directly address theoretical questions on the properties of tetraquarks.

MERCI !!!

- [1] J. P. Ader, J. M. Richard, and P. Taxil. DO NARROW HEAVY MULTI - QUARK STATES EXIST? *Phys. Rev.*, D25:2370, 1982.
- [2] Maurizio Alberti, Gunnar S. Bali, Sara Collins, Francesco Knechtli, Graham Moir, and Wolfgang Söldner. Hadroquarkonium from lattice QCD. *Phys. Rev. D*, 95(7):074501, 2017.
- [3] C. Alexandrou and G. Koutsou. The Static tetraquark and pentaquark potentials. *Phys.Rev.*, D71:014504, 2005.
- [4] S. Aoki and T. Aoki. Lattice study on a tetra-quark state Tbb in the HALQCD method. *PoS*, LATTICE2022, 2023.
- [5] Sinya Aoki, Noriyoshi Ishii, Takumi Doi, Tetsuo Hatsuda, Yoichi Ikeda, Takashi Inoue, Keiko Murano, Hidekatsu Nemura, and Kenji Sasaki. Extraction of Hadron Interactions above Inelastic Threshold in Lattice QCD. *Proc. Japan Acad. B*, 87:509–517, 2011.
- [6] Gunnar Bali and Martin Hetzenegger. Static-light meson-meson potentials. *PoS*, LATTICE2010:142, 2010.
- [7] Gunnar S. Bali. QCD forces and heavy quark bound states. *Phys. Rept.*, 343:1–136, 2001.
- [8] J. I. Ballot and J. M. Richard. FOUR QUARK STATES IN ADDITIVE POTENTIALS. *Phys. Lett.*, B123:449–451, 1983.
- [9] P. Bicudo and M. Cardoso. Tetraquark bound states and resonances in the unitary and microscopic triple string flip-flop quark model, the light-light-antiheavy-antiheavy $qq\bar{Q}\bar{Q}$ case study. *Phys. Rev. D*, 94(9):094032, 2016.
- [10] P. Bicudo, M. Cardoso, O. Oliveira, and P. J. Silva. Lattice QCD static potentials of the meson-meson and tetraquark systems computed with both quenched and full QCD. *Phys. Rev. D*, 96(7):074508, 2017.

- [11] Pedro Bicudo, Marco Cardoso, Antje Peters, Martin Pflaumer, and Marc Wagner. $ud\bar{b}\bar{b}$ tetraquark resonances with lattice QCD potentials and the Born-Oppenheimer approximation. *Phys. Rev. D*, 96(5):054510, 2017.
- [12] Pedro Bicudo, Krzysztof Cichy, Antje Peters, Björn Wagenbach, and Marc Wagner. Evidence for the existence of $ud\bar{b}\bar{b}$ and the non-existence of $ss\bar{b}\bar{b}$ and $cc\bar{b}\bar{b}$ tetraquarks from lattice QCD. 2015.
- [13] Pedro Bicudo, Krzysztof Cichy, Antje Peters, and Marc Wagner. BB interactions with static bottom quarks from Lattice QCD. *Phys. Rev. D*, 93(3):034501, 2016.
- [14] Pedro Bicudo, Antje Peters, Sebastian Velten, and Marc Wagner. Importance of meson-meson and of diquark-antidiquark creation operators for a $\bar{b}bud$ tetraquark. *Phys. Rev. D*, 103(11):114506, 2021.
- [15] Pedro Bicudo, Jonas Scheunert, and Marc Wagner. Including heavy spin effects in the prediction of a $\bar{b}bud$ tetraquark with lattice QCD potentials. *Phys. Rev. D*, 95(3):034502, 2017.
- [16] Pedro Bicudo and Marc Wagner. Lattice QCD signal for a bottom-bottom tetraquark. *Phys.Rev.*, D87(11):114511, 2013.
- [17] Benoit Blossier, Michele Della Morte, Georg von Hippel, Tereza Mendes, and Rainer Sommer. On the generalized eigenvalue method for energies and matrix elements in lattice field theory. *JHEP*, 04:094, 2009.
- [18] M. Born and R. Oppenheimer. Zur Quantentheorie der Molekeln. *Annalen der Physik*, 389:457, 1927.
- [19] Nora Brambilla, Simon Eidelman, Christoph Hanhart, Alexey Nefediev, Cheng-Ping Shen, Christopher E. Thomas, Antonio Vairo, and Chang-Zheng Yuan. The XYZ states: experimental and theoretical status and perspectives. *Phys. Rept.*, 873:1–154, 2020.

- [20] Raul A. Briceno. Two-particle multichannel systems in a finite volume with arbitrary spin. *Phys. Rev. D*, 89(7):074507, 2014.
- [21] Raul A. Briceno, Jozef J. Dudek, Robert G. Edwards, and David J. Wilson. Isoscalar $\pi\pi$, $K\bar{K}$, $\eta\eta$ scattering and the σ , f_0 , f_2 mesons from QCD. *Phys. Rev. D*, 97(5):054513, 2018.
- [22] Zachary S. Brown, William Detmold, Stefan Meinel, and Kostas Orginos. Charmed bottom baryon spectroscopy from lattice QCD. *Phys. Rev. D*, 90(9):094507, 2014.
- [23] Zachary S. Brown and Kostas Orginos. Tetraquark bound states in the heavy-light heavy-light system. *Phys.Rev.*, D86:114506, 2012.
- [24] M. Cardoso, N. Cardoso, and P. Bicudo. Variational study of the flux tube recombination in the two quarks and two quarks system in Lattice QCD. *Phys.Rev.*, D86:014503, 2012.
- [25] N. Cardoso, M. Cardoso, and P. Bicudo. Colour Fields Computed in SU(3) Lattice QCD for the Static Tetraquark System. *Phys.Rev.*, D84:054508, 2011.
- [26] Nuno Cardoso and Pedro Bicudo. Color fields of the static pentaquark system computed in SU(3) lattice QCD. *Phys.Rev.*, D87(3):034504, 2013.
- [27] Gavin K. C. Cheung, Christopher E. Thomas, Jozef J. Dudek, and Robert G. Edwards. Tetraquark operators in lattice QCD and exotic flavour states in the charm sector. *JHEP*, 11:033, 2017.
- [28] G. F. Chew and F. E. Low. Effective range approach to the low-energy p wave pion - nucleon interaction. *Phys. Rev.*, 101:1570–1579, 1956.
- [29] C. T. H. Davies, K. Hornbostel, A. Langnau, G. P. Lepage, A. Lidsey, J. Shigemitsu, and J. H. Sloan. Precision Upsilon spectroscopy from nonrelativistic lattice QCD. *Phys. Rev. D*, 50:6963–6977, 1994.

- [30] William Detmold, Kostas Orginos, and Martin J. Savage. BB Potentials in Quenched Lattice QCD. *Phys. Rev. D*, 76:114503, 2007.
- [31] Takumi Doi, Toru T. Takahashi, and Hideo Suganuma. Meson-meson and meson-baryon interactions in lattice QCD. *AIP Conf. Proc.*, 842(1):246–248, 2006.
- [32] Anthony Francis, Renwick J. Hudspith, Randy Lewis, and Kim Maltman. Lattice Prediction for Deeply Bound Doubly Heavy Tetraquarks. *Phys. Rev. Lett.*, 118(14):142001, 2017.
- [33] Anthony Francis, Renwick J. Hudspith, Randy Lewis, and Kim Maltman. Evidence for charm-bottom tetraquarks and the mass dependence of heavy-light tetraquark states from lattice QCD. *Phys. Rev. D*, 99(5):054505, 2019.
- [34] Tim Gershon. Exotic hadron naming convention. 6 2022.
- [35] C. Gignoux, B. Silvestre-Brac, and J. M. Richard. Possibility of Stable Multi - Quark Baryons. *Phys. Lett. B*, 193:323, 1987.
- [36] Andrea L. Guerrieri, Mauro Papinutto, Alessandro Pilloni, Antonio D. Polosa, and Nazario Tantalo. Flavored tetraquark spectroscopy. *PoS, LATTICE2014*:106, 2015.
- [37] Maxwell T. Hansen and Stephen R. Sharpe. Multiple-channel generalization of Lellouch-Luscher formula. *Phys. Rev. D*, 86:016007, 2012.
- [38] Maxwell T. Hansen and Stephen R. Sharpe. Relativistic, model-independent, three-particle quantization condition. *Phys. Rev. D*, 90(11):116003, 2014.
- [39] Maxwell T. Hansen and Stephen R. Sharpe. Expressing the three-particle finite-volume spectrum in terms of the three-to-three scattering amplitude. *Phys. Rev. D*, 92(11):114509, 2015.
- [40] R. Hudspith and D. Mohler. Non-perturbative heavy quark action tuning using machine learning. *PoS, LATTICE2022*, 2023.

- [41] R. J. Hudspith, B. Colquhoun, A. Francis, R. Lewis, and K. Maltman. A lattice investigation of exotic tetraquark channels. *Phys. Rev. D*, 102:114506, 2020.
- [42] Ciaran Hughes, Estia Eichten, and Christine T. H. Davies. Searching for beauty-fully bound tetraquarks using lattice nonrelativistic QCD. *Phys. Rev. D*, 97(5):054505, 2018.
- [43] N. Ishii, S. Aoki, and T. Hatsuda. The Nuclear Force from Lattice QCD. *Phys. Rev. Lett.*, 99:022001, 2007.
- [44] Noriyoshi Ishii, Sinya Aoki, Takumi Doi, Tetsuo Hatsuda, Yoichi Ikeda, Takashi Inoue, Keiko Murano, Hidekatsu Nemura, and Kenji Sasaki. Hadron–hadron interactions from imaginary-time Nambu–Bethe–Salpeter wave function on the lattice. *Phys. Lett. B*, 712:437–441, 2012.
- [45] Robert L. Jaffe. Multi-Quark Hadrons. 1. The Phenomenology of (2 Quark 2 anti-Quark) Mesons. *Phys.Rev.*, D15:267, 1977.
- [46] Robert L. Jaffe. Multi-Quark Hadrons. 2. Methods. *Phys. Rev. D*, 15:281, 1977.
- [47] Parikshit Junnarkar, Nilmani Mathur, and M. Padmanath. Study of doubly heavy tetraquarks in Lattice QCD. *Phys. Rev. D*, 99(3):034507, 2019.
- [48] Patrick Koppenburg. New particles discovered at the Lhc.
- [49] C.B. Lang, Daniel Mohler, Sasa Prelovsek, and Matija Vidmar. Coupled channel analysis of the rho meson decay in lattice QCD. *Phys.Rev.*, D84(5):054503, 2011.
- [50] G. Peter Lepage, Lorenzo Magnea, Charles Nakhleh, Ulrika Magnea, and Kent Hornbostel. Improved nonrelativistic QCD for heavy quark physics. *Phys. Rev. D*, 46:4052–4067, 1992.
- [51] Luka Leskovec, Stefan Meinel, Martin Pflaumer, and Marc Wagner. Lattice QCD investigation of a doubly-bottom $\bar{b}bud$ tetraquark with quantum numbers $I(J^P) = 0(1^+)$. *Phys. Rev. D*, 100(1):014503, 2019.

- [52] Randy Lewis and R. M. Woloshyn. $O(1/M^3)$ effects for heavy - light mesons in lattice NRQCD. *Phys. Rev. D*, 58:074506, 1998.
- [53] Randy Lewis and R. M. Woloshyn. Bottom baryons from a dynamical lattice QCD simulation. *Phys. Rev. D*, 79:014502, 2009.
- [54] Martin Luscher. Two particle states on a torus and their relation to the scattering matrix. *Nucl. Phys. B*, 354:531–578, 1991.
- [55] Martin Luscher and Ulli Wolff. How to Calculate the Elastic Scattering Matrix in Two-dimensional Quantum Field Theories by Numerical Simulation. *Nucl. Phys. B*, 339:222–252, 1990.
- [56] M. Lüscher. Volume Dependence of the Energy Spectrum in Massive Quantum Field Theories. 1. Stable Particle States. *Commun.Math.Phys.*, 104:177, 1986.
- [57] M. Lüscher. Volume Dependence of the Energy Spectrum in Massive Quantum Field Theories. 2. Scattering States. *Commun.Math.Phys.*, 105:153–188, 1986.
- [58] Aneesh V. Manohar. The HQET / NRQCD Lagrangian to order α/m^3 . *Phys. Rev. D*, 56:230–237, 1997.
- [59] Christopher Michael and P. Pennanen. Two heavy - light mesons on a lattice. *Phys. Rev. D*, 60:054012, 1999.
- [60] A. Mihaly, H. R. Fiebig, H. Markum, and K. Rabitsch. Interactions between heavy - light mesons in lattice QCD. *Phys. Rev. D*, 55:3077–3081, 1997.
- [61] Colin Morningstar, John Bulava, Bijit Singha, Ruairí Brett, Jacob Fallica, Andrew Hanlon, and Ben Hörz. Estimating the two-particle K -matrix for multiple partial waves and decay channels from finite-volume energies. *Nucl. Phys. B*, 924:477–507, 2017.
- [62] Fumiko Okiharu, Hideo Suganuma, and Toru T. Takahashi. Detailed analysis of the tetraquark potential and flip-flop in $SU(3)$ lattice QCD. *Phys.Rev.*, D72:014505, 2005.

- [63] Fumiko Okiharu, Hideo Suganuma, and Toru T. Takahashi. First study for the pentaquark potential in SU(3) lattice QCD. *Phys. Rev. Lett.*, 94:192001, 2005.
- [64] Bicudo P. Exotic tetraquarks with some heavy quarks in lattice QCD. *TBD*.
- [65] M. Padmanath, Sara Collins, Daniel Mohler, Stefano Piemonte, Sasa Prelovsek, Andreas Schäfer, and Simon Weishaeupl. Identifying spin and parity of charmonia in flight with lattice QCD. *Phys. Rev. D*, 99(1):014513, 2019.
- [66] M. Padmanath, C. B. Lang, and Sasa Prelovsek. X(3872) and Y(4140) using diquark-antidiquark operators with lattice QCD. *Phys. Rev. D*, 92(3):034501, 2015.
- [67] M. Padmanath and S. Prelovsek. Evidence for a doubly charm tetraquark pole in DD^* scattering on the lattice. 2 2022.
- [68] Antje Peters, Pedro Bicudo, Krzysztof Cichy, and Marc Wagner. Investigation of $B\bar{B}$ four-quark systems using lattice QCD. *J. Phys. Conf. Ser.*, 742(1):012006, 2016.
- [69] Antje Peters, Pedro Bicudo, and Marc Wagner. $b\bar{b}u\bar{d}$ four-quark systems in the Born-Oppenheimer approximation: prospects and challenges. *EPJ Web Conf.*, 175:14018, 2018.
- [70] Martin Pflaumer, Luka Leskovec, Stefan Meinel, and Marc Wagner. Existence and Non-Existence of Doubly Heavy Tetraquark Bound States. In *38th International Symposium on Lattice Field Theory*, 8 2021.
- [71] S. Prelovsek, H. Bahtiyar, and J. Petkovic. Z_b tetraquark channel from lattice QCD and Born-Oppenheimer approximation. *Phys. Lett. B*, 805:135467, 2020.
- [72] Sasa Prelovsek, Hüseyin Bahtiyar, and Jan Petkovic. Z_b tetraquark channel and $B\bar{B}^*$ interaction. *PoS, LATTICE2019*:012, 2019.
- [73] Sasa Prelovsek, C. B. Lang, Luka Leskovec, and Daniel Mohler. Study of the Z_c⁺ channel using lattice QCD. *Phys. Rev. D*, 91(1):014504, 2015.

- [74] Sasa Prelovsek and Luka Leskovec. Search for Z_c^+ (3900) in the 1^{+-} Channel on the Lattice. *Phys. Lett. B*, 727:172–176, 2013.
- [75] Mitja Sadl and Sasa Prelovsek. Tetraquark systems $\bar{b}b\bar{d}u$ in the static limit and lattice QCD. *Phys. Rev. D*, 104(11):114503, 2021.
- [76] B. A. Thacker and G. Peter Lepage. Heavy quark bound states in lattice QCD. *Phys. Rev. D*, 43:196–208, 1991.
- [77] M. Wagner, S. Meinel, and M. Pflauman. Search for $\bar{b}\bar{b}us$ and $\bar{b}\bar{c}ud$ tetraquark bound states using lattice QCD.
- [78] Marc Wagner. Forces between static-light mesons. *PoS, LATTICE2010*:162, 2010.
- [79] Marc Wagner. Static-static-light-light tetraquarks in lattice QCD. *Acta Phys. Polon. Supp.*, 4:747–752, 2011.
- [80] S. Zouzou, B. Silvestre-Brac, C. Gignoux, and J.M. Richard. FOUR QUARK BOUND STATES. *Z.Phys.*, C30:457, 1986.