

Electroweak logarithms in OpenLoops

Lorenzo Mai



Parton Shower and Resummation

Milano

06-08/06/2023

Artwork by C.Z.

Introduction

- In the energy range above the EW scale ($\sqrt{s} \gg M_W$), Sudakov logs represent the leading contribution of EW radiative corrections
- Sudakov logarithms from N^n LO EW corrections

$$\alpha^n \log^k \frac{s}{M_W^2}, \quad 1 \leq k \leq 2n$$

- At NLO

Double logs:


$$L(s) = \frac{\alpha}{4\pi} \log^2 \frac{s}{M_W^2},$$

Single logs:


$$l(s) = \frac{\alpha}{4\pi} \log \frac{s}{M_W^2}$$

Introduction


- Significant enhancement of tails of kinematic distributions up to several tens percent



(Bothmann, Napoletano [2006.14635](#); 2020)



(Pagani, Zaro [2110.03714](#); 2021)



(Chiesa et al. [1305.6837](#); 2013)

Framework: notation & conventions

- $n \rightarrow 0$ process

$$\varphi_{i_1}(p_1) \dots \varphi_{i_n}(p_n) \rightarrow 0$$

with not mass suppressed Born matrix-element, i.e. $\mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i_n}} \sim E^d$

- DP algorithm based on logarithmic approximation (LA):

→ Hierarchy scales

$$\mu^2 = s \sim (p_k + p_l)^2 \gg m_t^2, M_H^2 > M_{Z,W}^2 \gg m_f^2 \gg \lambda^2, \quad \forall k, l$$


→ At one-loop keep only double and singular logarithmic corrections

$$\delta^{\text{DL}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} \sim E^d \textcolor{red}{L} \qquad \qquad \delta^{\text{SL}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} \sim E^d \textcolor{red}{l}$$

neglecting constant ($\sim \alpha E^d$) and mass suppressed ($\sim M^n E^{d-n} \textcolor{red}{L}$) contributions


Double Logs (DL)

- DL originate from triangle diagrams where two external legs exchange a *soft and collinear (SC)* gauge boson V



Double Logs (DL)

- DL originate from triangle diagrams where two external legs exchange a *soft and collinear (SC)* gauge boson V
- In the *Eikonal approximation*, the loop integral reduces to the scalar three-point function C_0 , which factorises




$$\delta^{\text{DL}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} \stackrel{\text{LA}}{=} \sum_k \sum_{l < k} \sum_V \sum_{k', l'} \frac{\alpha}{4\pi} I_{kk'}^V I_{ll'}^{\bar{V}} \underbrace{\left[\log^2 \frac{|r_{kl}|}{M_V^2} - 2i\pi \Theta(r_{kl}) \log \frac{|r_{kl}|}{M_V^2} \right]}_{\propto C_0|_{\text{LA}}} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i'_l} \dots \varphi_{i_n}}$$

with $r_{kl} = (p_k + p_l)^2$

Double Logs (DL)

- DL originate from triangle diagrams where two external legs exchange a *soft and collinear (SC) gauge boson V*
- In the *Eikonal approximation*, the loop integral reduces to the scalar three-point function C_0 , which *factorises*



$$\delta^{\text{DL}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} \stackrel{\text{LA}}{=} \sum_k \sum_{l < k} \sum_V \sum_{k', l'} \frac{\alpha}{4\pi} I_{kk'}^V I_{ll'}^{\bar{V}} \underbrace{\left[\log^2 \frac{|r_{kl}|}{M_V^2} - 2i\pi \Theta(r_{kl}) \log \frac{|r_{kl}|}{M_V^2} \right]}_{\propto C_0|_{\text{LA}}} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i'_l} \dots \varphi_{i_n}}$$

with $r_{kl} = (p_k + p_l)^2$

- Consequence of C_0 *factorisation*: DL are *universal*, i.e. process independent

Double Logs: LSC, SSC, S-SSC

- DL can be split into

→ Leading soft-collinear (LSC): angular independent, single sum over external legs

$$\delta^{\text{LSC}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} = \sum_k \sum_{k'} \sum_V \delta_{kk'}^{\text{LSC}, V} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i_n}}, \quad \delta_{kk'}^{\text{LSC}, V} \sim \frac{\alpha}{4\pi} I^V I^{\bar{V}} \log^2 \left(\frac{s}{M_V^2} \right)$$

→ Subleading soft-collinear (SSC) and S-SSC: angular dependent, double sum over external legs

$$\delta^{(\text{S-})\text{SSC}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} = \sum_k \sum_{l < k} \sum_{k', l'} \sum_V \delta_{kk' ll'}^{(\text{S-})\text{SSC}, V} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i'_l} \dots \varphi_{i_n}}$$

$$\delta_{kk' ll'}^{\text{SSC}, V} \sim \frac{\alpha}{4\pi} I^V I^{\bar{V}} \log \left(\frac{s}{M_V^2} \right) \log \left(\frac{|r_{kl}|}{s} \right), \quad \delta_{kk' ll'}^{\text{S-SSC}, V} \sim \frac{\alpha}{4\pi} I^V I^{\bar{V}} \log^2 \left(\frac{|r_{kl}|}{s} \right)$$

Single Logs (SL)

- SL have a triple origin

Single Logs (SL): PR

- SL have a triple origin
→ PR: renormalisation of EW dimensionless parameters

$$\mu_{i,0}^2 = \mu_i^2 + \delta\mu_i^2$$

$$\varphi_{i,0} = \left(1 + \frac{1}{2}\delta Z_{\varphi_i \varphi_j}\right) \varphi_j$$

$$g_{i,0} = g_i + \delta g_i = (1 + \delta Z_{g_i}) g_i$$

$$g_{i,0} = g_i + \delta g_i = (1 + \delta Z_{g_i}) g_i$$

Single Logs (SL): PR

- SL have a triple origin
- PR: renormalisation of EW dimensionless parameters

$$\begin{aligned}\mu_{i,0}^2 &= \mu_i^2 + \delta\mu_i^2 \\ \varphi_{i,0} &= \left(1 + \frac{1}{2}\delta Z_{\varphi_i \varphi_j}\right) \varphi_j \\ g_{i,0} &= g_i + \delta g_i = (1 + \delta Z_{g_i}) g_i\end{aligned}$$

$$g_{i,0} = g_i + \delta g_i = (1 + \delta Z_{g_i}) g_i$$

- WF: wave-function renormalisation of external fields

$$\varphi_{i,0} = \left(1 + \frac{1}{2}\delta Z_{\varphi_i \varphi_j}\right) \varphi_j$$

yields to the factorised correction

$$\delta^{\text{WF}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} = \sum_k \sum_{k'} \delta_{kk'}^{\text{WF}} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i_n}},$$

$$\delta_{kk'}^{\text{WF}} = \frac{1}{2} \delta Z_{kk'}$$

Single Logs (SL): PR

- SL have a triple origin

→ PR: renormalisation of EW dimensionless parameters

$$\mu_{i,0}^2 = \mu_i^2 + \delta\mu_i^2$$

$$\varphi_{i,0} = \left(1 + \frac{1}{2}\delta Z_{\varphi_i \varphi_j}\right) \varphi_j$$

$$g_{i,0} = g_i + \delta g_i = (1 + \delta Z_{g_i}) g_i$$

$$g_{i,0} = g_i + \delta g_i = (1 + \delta Z_{g_i}) g_i$$


→ WF: wave-function renormalisation of external fields

$$\varphi_{i,0} = \left(1 + \frac{1}{2}\delta Z_{\varphi_i \varphi_j}\right) \varphi_j$$

yields to the factorised correction

$$\delta^{\text{WF}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} = \sum_k \sum_{k'} \delta_{kk'}^{\text{WF}} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i_n}},$$


$$\delta_{kk'}^{\text{WF}} = \frac{1}{2} \delta Z_{kk'}$$



Renormalisation of masses and couplings with mass dimensions brings only mass-suppressed corrections

Single Logs (SL): Coll


- SL have a triple origin
→ **Coll**: external leg emission of a collinear gauge boson



Single Logs (SL): Coll

- SL have a triple origin

→ **Coll**: external leg emission of a collinear gauge boson



Its evaluation in *Eikonal approximation* leads to the factorised contribution


$$\delta^{\text{coll}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} = \sum_k \sum_{k'} \delta_{kk'}^{\text{coll}} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i_n}},$$

$$\delta_{kk'}^{\text{coll}} \sim \frac{\alpha}{4\pi} \sum_V I^V I^{\bar{V}} \log \left(\frac{s}{M_V^2} \right)$$

Single Logs (SL): Coll

- SL have a triple origin

→ **Coll**: external leg emission of a collinear gauge boson



Its evaluation in *Eikonal approximation* leads to the factorised contribution

$$\delta^{\text{coll}} \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} = \sum_k \sum_{k'} \delta_{kk'}^{\text{coll}} \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i_n}},$$

$$\delta_{kk'}^{\text{coll}} \sim \frac{\alpha}{4\pi} \sum_V I^V I^{\bar{V}} \log \left(\frac{s}{M_V^2} \right)$$

→ **C**: Full gauge-invariant SL correction associated to external fields:

$$\delta^C \mathcal{M}^{\varphi_{i_1} \dots \varphi_{i_n}} = \sum_k \sum_{k'} \delta_{kk'}^C \mathcal{M}_0^{\varphi_{i_1} \dots \varphi_{i'_k} \dots \varphi_{i_n}},$$

$$\delta_{kk'}^C = (\delta_{kk'}^{\text{coll}} + \delta_{kk'}^{\text{WF}})|_{\mu^2=s}$$

Implementation in OpenLoops: why

- NLO EW corrections have been almost fully automated nowadays

Implementation in OpenLoops: why

- NLO EW corrections have been almost fully automated nowadays
- EW Sudakov logarithms at one-loop already implemented in
 - ▶ ALPGEN: Chiesa *et al*, [1305.6837](#); 2013
 - ▶ Sherpa: Bothmann, Napoletano [2006.14635](#); 2020
 - ▶ MadGraph: Pagani, Zaro [2110.03714](#); 2021

Implementation in OpenLoops: why

- NLO EW corrections have been almost fully automated nowadays
- EW Sudakov logarithms at one-loop already implemented in
 - ▶ ALPGEN: Chiesa *et al*, [1305.6837](#); 2013
 - ▶ Sherpa: Bothmann, Napoletano [2006.14635](#); 2020
 - ▶ MadGraph: Pagani, Zaro [2110.03714](#); 2021

However:

- ▶ Even if automated, one-loop computations can be very complicated (e.g. high multiplicity processes)
- ▶ No NNLO/two-loop level automation available
- ▶ EW Sudakov logs have nice properties: factorisation, being the leading contribution of radiative corrections

Implementation in OpenLoops: why

- NLO EW corrections have been almost fully automated nowadays
- EW Sudakov logarithms at one-loop already implemented in
 - ▶ ALPGEN: Chiesa *et al*, [1305.6837](#); 2013
 - ▶ Sherpa: Bothmann, Napoletano [2006.14635](#); 2020
 - ▶ MadGraph: Pagani, Zaro [2110.03714](#); 2021

However:

- ▶ Even if automated, one-loop computations can be very complicated (e.g. high multiplicity processes)
 - ▶ No NNLO/two-loop level automation available
 - ▶ EW Sudakov logs have nice properties: factorisation, being the leading contribution of radiative corrections
- OpenLoops (OL): automated tool for the calculation of tree and one-loop amplitudes (Buccioni *et al*, [1907.13071](#); 2019)
 - Goal of the implementation: evaluate NLO EW Sudakov corrections via tree amplitudes (w/o loop computations) and make them available to any MC with OL interface

Implementation in OpenLoops: how

- Representation of Denner-Pozzorini algorithm via effective CT vertices

$$\begin{array}{c} V \\ \hline \varphi & \varphi' \end{array} \longrightarrow \begin{array}{c} V \\ \bullet \\ \hline \varphi & \varphi' \end{array} = ieI_{\varphi\varphi'}^V K_{\text{ew}}^V$$

reducing one-loop amplitudes to tree-level ones via double CT insertions


Implementation in OpenLoops: how

- Representation of Denner-Pozzorini algorithm via effective CT vertices

$$\begin{array}{c} V \\ \hbox{\scriptsize wavy line} \\ \hbox{\scriptsize horizontal line} \\ \varphi \qquad \varphi' \end{array} \longrightarrow \begin{array}{c} V \\ \bullet \\ \hbox{\scriptsize horizontal line} \\ \varphi \qquad \varphi' \end{array} = ie I_{\varphi\varphi'}^V K_{\text{ew}}^V$$

reducing one-loop amplitudes to tree-level ones via double CT insertions

Eg.: Drell-Yann




Implementation in OpenLoops: how

- Effective CT vertices are suitable for evaluation of *soft-collinear* and *collinear Sudakov* corrections

Implementation in OpenLoops: how


- Effective CT vertices are suitable for evaluation of *soft-collinear* and *collinear Sudakov* corrections
- Single logs coming from **PR** contributions can be evaluated via generation of standard UV counterterms, e.g.



setting all the **WFRCs** to zero

Implementation in OpenLoops: how



- Effective CT vertices are suitable for evaluation of *soft-collinear* and *collinear Sudakov* corrections
- Single logs coming from **PR** contributions can be evaluated via generation of standard UV counterterms, e.g.




setting all the **WFRCs** to zero

- Alternative way: set $\delta_{kk'}^{\text{WF}}$ to zero and evaluate **WF** + **PR** via standard UV counterterms


Results: $pp \rightarrow W^+W^-Z$



Results: $pp \rightarrow W^+W^-Z$





Results: $pp \rightarrow W^+W^-Z$





$\delta_{kk' ll'}^{\text{S-SSC}, V} \sim \frac{\alpha}{4\pi} I^V I^{\bar{V}} \log^2 \left(\frac{|r_{kl}|}{s} \right)$

 S-SSC relevant for PS regions where
 $s \sim (p_k + p_l)^2 \gg M_{Z,W}^2 \quad \forall k, l$
 is violated.

Results: $pp \rightarrow W^+W^-Z$




Results: $pp \rightarrow W^+W^-Z$





However, no full control on S-SSC term!

Arising also
from box diagrams





$$\sim D_0 \sim \log \left(\frac{|r_{kl}|}{s} \right)$$

Results: $pp \rightarrow \bar{t}tW$



Results: $pp \rightarrow \bar{t}tW$



Implementation in OpenLoops: resonances

Implementation in OpenLoops: resonances


- DP algorithm:
 - ▶ At $\sqrt{s} \gg M_W$, NLO EW radiative corrections are DL and SL
 - ▶ These corrections are *universal*, i.e. are associated to external states only

Implementation in OpenLoops: resonances

- DP algorithm:
 - ▶ At $\sqrt{s} \gg M_W$, NLO EW radiative corrections are DL and SL
 - ▶ These corrections are *universal*, i.e. are associated to external states only
- Consequence: algorithm not suitable for processes dominated by resonant configurations


Implementation in OpenLoops: resonances

- DP algorithm:
 - ▶ At $\sqrt{s} \gg M_W$, NLO EW radiative corrections are DL and SL
 - ▶ These corrections are *universal*, i.e. are associated to external states only
- Consequence: algorithm not suitable for processes dominated by resonant configurations
- E.g.: partonic channel of $pp \rightarrow Zj \rightarrow \bar{l}l j$



Implementation in OpenLoops: resonances

- DP algorithm:
 - ▶ At $\sqrt{s} \gg M_W$, NLO EW radiative corrections are DL and SL
 - ▶ These corrections are *universal*, i.e. are associated to external states only
- Consequence: algorithm not suitable for processes dominated by resonant configurations
- E.g.: partonic channel of $pp \rightarrow Zj \rightarrow \bar{l}l j$



LSC, C:

$$\delta_{kk}^{\text{LSC,C}}, \quad k \in \{q, \bar{q}, l, \bar{l}\}$$

SSC, S-SSC:


$$\delta_{kl}^{(\text{S-})\text{SSC}}, \quad k \neq l \text{ and } k, l \in \{q, \bar{q}, l, \bar{l}\}$$

PR:

CTs for $Z\bar{q}q, Z\bar{l}l$ vertices

Implementation in OpenLoops: resonances

- E.g.: partonic channel of $pp \rightarrow Zj \rightarrow \bar{l}l j$




LSC, C: $\delta_{kk}^{\text{LSC,C}}, \quad k \in \{q, \bar{q}, l, \bar{l}\}$

SSC, S-SSC: $\delta_{kl}^{(\text{S-})\text{SSC}}, \quad k \neq l \text{ and } k, l \in \{q, \bar{q}, l, \bar{l}\}$

PR: CTs for $Z\bar{q}q, Z\bar{l}l$ vertices

- In the kinematic region where the Z boson is nearly on shell



LSC, C: $\delta_{kk}^{\text{LSC,C}}, \quad k \in \{q, \bar{q}, Z\}$

SSC, S-SSC: $\delta_{kl}^{(\text{S-})\text{SSC}}, \quad k \neq l \text{ and } k, l \in \{q, \bar{q}, Z\}$



PR: CT for $Z\bar{q}q$ vertex

Implementation in OpenLoops: resonances

- Solution: evaluation of Sudakov corrections associated to both Z and $\{l, \bar{l}\}$ with different weights $P_i(k_i)$


Implementation in OpenLoops: resonances

- Solution: evaluation of Sudakov corrections associated to both Z and $\{l, \bar{l}\}$ with different weights $P_i(k_i)$




Implementation in OpenLoops: resonances



- Solution: evaluation of Sudakov corrections associated to both Z and $\{l, \bar{l}\}$ with different weights $P_i(k_i)$



$$P_{X_i}(k_i) = \left| \frac{\mu_{X_i}^2 - M_{X_i}^2 \Gamma_{X_i}^2}{(k_i^2 - \mu_{X_i}^2)^2 + \mu_{X_i}^2} \right| = \begin{cases} 1 & \text{if } k_i^2 \rightarrow M_{X_i}^2 \\ 0 & \text{if } k_i^2 \rightarrow \infty \end{cases}$$




Results: $pp \rightarrow e^+e^-j$



- External insertions approach (as well Sudakov corrections to the hard process only) fail in reproducing the full NLO_{VI} prediction for $m_{e^+e^-}$ range "capturing" the resonance
- Issue naturally solved with internal insertions technique via projectors
- Automatic recover of standard algorithm when far from the resonance

Conclusions and outlook

- In the **EW** sector, radiative corrections at high energies are dominated by Sudakov logarithms which significantly enhance tails of kinematic distributions ($> 10\%$)
- Exploiting the universality of Sudakov logs we developed an effective CT vertex approach for the DP algorithm and implemented it in OpenLoops




Reduction of one-loop **EW** corrections to a tree-level problem with percent level of accuracy

- Additional aspects of the implementation:
 - ▶ Model independent
 - ▶ Direct employment in PS Event Generators with OL interface
 - ▶ Can be used together with differential QED radiation at NLO (both mass and dim reg are available)
 - ▶ Support **EW** corrections for resonant processes (novelty)
- Outlook:
 - ▶ Resummation of logarithms to preserve perturbation theory
 - ▶ Suitable for NNLO extension

Backup

Single Logs: PR


- Generic two-point function



$$\frac{i}{(4\pi)^2} B_{\{0,\mu,\mu\nu\}}(p, m_0, m_1) := \mu^{4-D} \int \frac{d^D q}{(2\pi)^D} \frac{\{1, q_\mu, q_\mu q_\nu\}}{(q^2 - m_0^2 + i\varepsilon) [(q + p)^2 - m_1^2 + i\varepsilon]}$$

Single Logs: PR

- Generic two-point function




$$\frac{i}{(4\pi)^2} B_{\{0,\mu,\mu\nu\}}(p, m_0, m_1) := \mu^{4-D} \int \frac{d^D q}{(2\pi)^D} \frac{\{1, q_\mu, q_\mu q_\nu\}}{(q^2 - m_0^2 + i\varepsilon) [(q + p)^2 - m_1^2 + i\varepsilon]}$$

- In LA $\mu^2 = s \gg p^2, m_0^2, m_1^2 \Rightarrow$ four possible hierarchy of masses

- (a) $m_i^2 \ll p^2$ and $p^2 - m_{1-i}^2 \ll p^2$ for $i = 0$ or $i = 1$,
- (b) not (a) and $m_i^2 \not\asymp p^2$ for $i = 0, 1$,
- (c) $m_0^2 = m_1^2 \gg p^2$
- (d) $m_i^2 \gg p^2 \not\asymp m_{1-i}^2$ for $i = 0$ or $i = 1$

Single Logs: PR



- Generic two-point function

$$\frac{i}{(4\pi)^2} B_{\{0,\mu,\mu\nu\}}(p, m_0, m_1) := \mu^{4-D} \int \frac{d^D q}{(2\pi)^D} \frac{\{1, q_\mu, q_\mu q_\nu\}}{(q^2 - m_0^2 + i\varepsilon) [(q+p)^2 - m_1^2 + i\varepsilon]}$$

- Results for two point functions and their derivatives

$$B_0(p, m_0, m_1) \stackrel{\text{LA}}{=} \log \frac{\mu^2}{M^2},$$

$$B_1(p^2, m_0, m_1) \stackrel{\text{LA}}{=} -\frac{1}{2} \log \frac{\mu^2}{M^2},$$

$$\frac{1}{p^2} B_{00}(p^2, m_0, m_1) \stackrel{\text{LA}}{=} \frac{3m_0^2 + 3m_1^2 - p^2}{12p^2} \log \frac{\mu^2}{M^2},$$



$$\frac{1}{p^2} g^{\mu\nu} B_{\mu\nu}(p^2, m_0, m_1) \stackrel{\text{LA}}{=} \frac{m_0^2 + m_1^2}{p^2} \log \frac{\mu^2}{M^2}$$

$$p^2 B'_0(p, m_0, m_1) \stackrel{\text{LA}}{=} \frac{1}{2} \log \frac{m_{1-i}^2}{m_i^2} = \frac{1}{2} \log \frac{p^2}{\lambda^2},$$

$$p^2 B'_1(p^2, m_0, m_1) + \frac{1}{2} p^2 B'_0(p, m_0, m_1) \stackrel{\text{LA}}{=} -\frac{1}{4} \log \frac{m_0^2}{m_1^2}$$

Implementation in OpenLoops: resonances

- Solution: evaluation of Sudakov corrections associated to both Z and $\{l, \bar{l}\}$ with different weights $P_i(k_i)$



Implementation in OpenLoops: projectors

- Explicit expression of the projectors for unstable particles X

$$P_{X_i}(k_i) = \left| \frac{\mu_{X_i}^2 - M_{X_i}^2 \Gamma_{X_i}^2}{(k_i^2 - \mu_{X_i}^2)^2 + \mu_{X_i}^2} \right| = \begin{cases} 1 & \text{if } k_i^2 \rightarrow M_{X_i}^2 \\ 0 & \text{if } k_i^2 \rightarrow \infty \end{cases}$$

Implementation in OpenLoops: projectors

- Explicit expression of the projectors for unstable particles X

$$P_{X_i}(k_i) = \left| \frac{\mu_{X_i}^2 - M_{X_i}^2 \Gamma_{X_i}^2}{(k_i^2 - \mu_{X_i}^2)^2 + \mu_{X_i}^2} \right| = \begin{cases} 1 & \text{if } k_i^2 \rightarrow M_{X_i}^2 \\ 0 & \text{if } k_i^2 \rightarrow \infty \end{cases}$$



- Unitarity is violated but it can be restored:

- ▶ Evaluation of $P_{X_i}(k_i)$ for a given psp



- ▶ Generation of random number $0 \leq a \leq 1$

- ▶ Choice $P_{X_i} = \begin{cases} 1 & \text{if } P_{X_i} \geq a \\ 0 & \text{if } P_{X_i} \leq a \end{cases}$



Results: $pp \rightarrow Z + j$





Results: $pp \rightarrow Z + j$





Results: $pp \rightarrow ZZj$





Results: $pp \rightarrow ZZj$



Results: $pp \rightarrow ZZ$





Results: $pp \rightarrow ZZ$





Results: $pp \rightarrow e^+e^-j$

$75 \text{ GeV} \leq m_{e^+e^-} \leq 107 \text{ GeV}$





Results: $pp \rightarrow e^+e^-j$

$75 \text{ GeV} \leq m_{e^+e^-} \leq 107 \text{ GeV}$



Results: $pp \rightarrow e^+e^-j$

$200 \text{ GeV} \leq m_{e^+e^-} \leq 500 \text{ GeV}$



Results: $pp \rightarrow e^+e^-j$

$75 \text{ GeV} \leq m_{e^+e^-} \leq 107 \text{ GeV}$

