


# Probing light Yukawa couplings at the (HL-) LHC

Ramona Gröber


based on work with L. Alasfar and  
R. Corral Lopez, JHEP 11 (2019)088

L. Alasfar, C. Grojean, A. Paul,  
Z. Qian JHEP 11 (2022)045


work with E. Balzaní, M. Vitti, to  
appear



12/04/2023



# Higgs couplings



3rd generation fermion and gauge boson couplings to Higgs boson fairly good measured

2nd generation fermion couplings first results available

Higgs self-couplings?

First generation Yukawa couplings?

# Light quark Yukawa couplings

HL-LHC prospects for measurement of 1st and 2nd generation quark Yukawa couplings

$$\kappa = y_q/y_q^{SM}$$

Ide Blas, Cepeda, d'Hondt et al '19]

$$|\kappa_u| \leq 570, \quad |\kappa_d| \leq 270, \quad |\kappa_s| \leq 13, \quad |\kappa_c| \leq 1.2$$

global fit, not completely model-independent

# Light quark Yukawa couplings

HL-LHC prospects for measurement of 1st and 2nd generation quark Yukawa couplings

$$\kappa = y_q / y_q^{SM}$$

[de Blas, Cepeda, d'Hondt et al '19]

$$|\kappa_u| \leq 570, \quad |\kappa_d| \leq 270, \quad |\kappa_s| \leq 13, \quad |\kappa_c| \leq 1.2$$

global fit, not completely model-independent

Alternative ways:

- Higgs kinematics: Higgs + jet transverse momentum distribution

[Bishara Haisch, Monni, Re '16;  
Soreq, Zhu, Zupan '16]

- Higgs decays to photon and vector mesons

[Bodwin, Pietrello, Stoynev, Velasco '13; Kagan,  
Perez, Pietrello, Soreq, Stoynev, Zupan '14;  
Alte, König, Neubert '16  
ATLAS 1712.02758, CMS 2007.05122]

- Charm tagging (strange tagging at lepton colliders)

[Perez, Soreq, Stamou, Tobioka '15;  
Brivio, Goertz, Isidori '15;  
ATLAS 1802.04329, CMS 1912.01662;  
Duarte-Camderros, Perez, Schlaffer, Soffer '18]

- various other proposals

[Yu '17, Aguilar-Saavedra, Cano, No '18, Falkowski et al. '20, Vignaroli '22]



# Light quark Yukawa couplings

HL-LHC prospects for measurement of 1st and 2nd generation quark Yukawa couplings

$$\kappa = y_q/y_q^{SM}$$

[de Blas, Cepeda, d'Hondt et al '19]

$$|\kappa_u| \leq 570, \quad |\kappa_d| \leq 270, \quad |\kappa_s| \leq 13, \quad |\kappa_c| \leq 1.2$$

global fit, not completely model-independent

Alternative ways:

- Higgs kinematics: Higgs+jet transverse momentum distribution

[Bishara Haisch, Monni, Re '16;  
Soreq, Zhu, Zupan '16]

- Higgs decays to photon and vector mesons

[Bodwin, Pietrello, Stoynev, Velasco '13; Kagan,  
Perez, Pietrello, Soreq, Stoynev, Zupan '14;  
Alte, König, Neubert '16  
ATLAS 1712.02758, CMS 2007.05122]

- Charm tagging (strange tagging at lepton colliders)

[Perez, Soreq, Stamou, Tobioka '15;  
Brivio, Goertz, Isidori '15;  
ATLAS 1802.04329, CMS 1912.01662;  
Duarte-Campanerros, Perez, Schlaffer, Soffer '18]

In this talk: explore the potential of Higgs pair production and off-shell Higgs production for constraining first generation quarks

# SMEFT

$$\mathcal{L}_{SM} \supset -y_{ij}^u \bar{Q}_L^i \tilde{\phi} u_R^j - y_{ij}^d \bar{Q}_L^i \phi d_R^j + h.c.$$

At dim-6 level the Higgs couplings to fermions are modified by the operator

$$\mathcal{L}_{dim6} \supset \frac{c_{ij}^u}{\Lambda^2} (\phi^\dagger \phi) \bar{Q}_L^i \tilde{\phi} u_R^j + \frac{c_{ij}^d}{\Lambda^2} (\phi^\dagger \phi) \bar{Q}_L^i \phi d_R^j + h.c.$$

mass eigenbasis:

$$\tilde{c}_{ij}^q = (V_q^L)^*_{ki} c_{kl}^q V_{lj}^R$$

Couplings:

$$g_{h\bar{q}_i q_j} = \frac{m_{q_i}}{v} \delta_{ij} - \frac{v^2}{\Lambda^2} \frac{\tilde{c}_{ij}^q}{\sqrt{2}}$$

$$g_{hh\bar{q}_i q_j} = -\frac{3}{2\sqrt{2}} \frac{v^2}{\Lambda^2} \tilde{c}_{ij}^q$$

direct coupling to Higgs pair

$$g_{G_0 G_0 \bar{q}_i q_j} = -\frac{1}{2\sqrt{2}} \frac{v^2}{\Lambda^2} \tilde{c}_{ij}^q$$


In the following consider only flavour diagonal case.

Notation:


$$g_{h\bar{q}q} = \kappa_q g_{h\bar{q}q}^{SM}$$

$$g_{hh\bar{q}q} = -\frac{3}{2} \frac{1 - \kappa_q}{v} g_{h\bar{q}q}^{SM}$$


# Large light quark Yukawas



heavy scalar



vectorlike quarks



vectorlike quark + heavy scalar

concrete models:

2HDM with spontaneous flavour violation

[Egana-Usgrinovic,  
Homiller, Meade '18, '19]



vector-like quarks + flavour symmetries

[Bar-Shalom, Soni '18]




Higgs pair production

# Higgs pair production

Higgs pair production in SM, gluon fusion dominated by heavy quark loops



enhanced light Yukawa couplings




contribution most important for 1st generation (given the coupling limits)

# Higgs pair production


$pp \rightarrow hh (q\bar{q}A) \quad \sqrt{s} = 14 \text{ TeV}$

[Alasfar, Corral Lopez, RG '19]



increase of cross section,  
(also modified distributions)

$hh \rightarrow b\bar{b}\gamma\gamma$



[Alasfar, Corral Lopez, RG '19]

decrease of BR for typical di-  
Higgs final state

cut and count analysis:  $\kappa_u < 1251, \kappa_d < 610$



# Machine Learning

Can extract both light quark Yukawas and trilinear Higgs self-coupling from di-Higgs?

We use Boosted Decision Trees  
and a comprehensive set of  
kinematic variables instead of  
four-momenta

we are interested in  
interpretability of our result

$$p_T^{b_1}, p_T^{b_2}, p_T^{\gamma_1}, p_T^{\gamma\gamma}$$

$$\eta_{b_{j1}}, \eta_{b_{j2}}, \eta_{\gamma_1}, \eta_{\gamma\gamma}$$

$$n_{bjet}, n_{jet}, \Delta R_{min}^{b\gamma}, \Delta\phi_{min}^{bb}$$

$$m_{\gamma\gamma}, m_{bb}, m_{b_1, h}, m_{b\bar{b}h}, H_T$$

To learn the shapes of the various contributions  
we divide into several categories:

$$Q\bar{Q}h = b\bar{b}h(h \rightarrow \gamma\gamma), t\bar{t}h(h \rightarrow \gamma\gamma)$$

$$b\bar{b}\gamma\gamma$$

$$d\bar{d}hh, u\bar{u}hh$$

$$hh_{tri}^{ggF}, hh_{int}^{ggF}$$

$$hh_{box}^{ggF}$$

background

background

signal for enhanced light quark couplings

signal for trilinear Higgs self-coupling


background

# Interpretable ML: Shapley values


measure of importance of a variable from game theory

[L.S. Shapley '51]


The value of each player and each combination of players





The value of the player in each game



marginalised  
values

$$(7+7+10+3+9+10) / 6 =$$




$$(4+0+4+4+4+3) / 6 =$$


$$(8+12+5+12+6+6) / 6 =$$




most  
important  
player

# Interpretable ML: Shapley values




relative importance of the various kinematic variables for signal background discrimination

# Results

[Alasfar, RG, Grojean,  
Paul, Qian '22]

$$\kappa_d = [-392.84, 394.73]$$




HL-LHC  
Best Fit Point:  
 $\kappa_d = 1.0$   
 $\kappa_\lambda = 1.0$

We performed several one-/two-  
and three-parameter fits


$\kappa_\lambda = [0.53, 1.73]$   
1 parameter fit

$$\kappa_\lambda = [0.64, 2.20]$$



here we can see that the  
sensitivity on the trilinear Higgs  
self-coupling is diluted in two-  
parameter fit

# Results



huge improvement over cut-and count analysis

Off-shell Higgs production



# Off-shell Higgs production

Considered as probe of Higgs width

[Kauer, Passarino '12, Carla, Melniko '13,  
Campbell, Ellis, Williams '13]

$$\frac{\mu_{on}}{\mu_{off}} \propto \frac{\kappa_{ggh}^2(m_h) \kappa_{hZZ}^2(m_h)}{\Gamma_h / \Gamma_h^{SM}} \frac{1}{\kappa_{ggh}^2(m_{4\ell}) \kappa_{hZZ}^2(m_{4\ell})}$$

works for

[Englert, (Soreq), Spannowsky '14]

$$\kappa_{ggh}(m_h) = \kappa_{ggh}(m_{4\ell})$$



$$\kappa_{hZZ}(m_h) = \kappa_{hZZ}(m_{4\ell})$$

$$\text{CMS: } 3.2_{-1.7}^{+2.4} \text{ GeV}$$

[CMS in Nature 18 (2022) 1392]

$$\text{ATLAS: } 4.6_{-2.6}^{+2.6} \text{ GeV}$$

[ATLAS-CONF-2022-068]



For enhanced light quark Yukawa couplings it does not work:

new production channel to be added, spoils the “model-independence” of width measurement

# Off-shell Higgs production

Considered as probe of Higgs width

[Kauer, Passarino '12, Carla, Melniko '13,  
Campbell, Ellis, Williams '13]

$$\frac{\mu_{on}}{\mu_{off}} \propto \frac{\kappa_{ggh}^2(m_h) \kappa_{hZZ}^2(m_h)}{\Gamma_h / \Gamma_h^{SM}} \frac{1}{\kappa_{ggh}^2(m_{4\ell}) \kappa_{hZZ}^2(m_{4\ell})}$$

works for

[Englert, (Soreq), Spannowsky '14]

$$\kappa_{ggh}(m_h) = \kappa_{ggh}(m_{4\ell})$$



$$\kappa_{hZZ}(m_h) = \kappa_{hZZ}(m_{4\ell})$$

$$\text{CMS: } 3.2_{-1.7}^{+2.4} \text{ MeV}$$

[CMS in Nature 18 (2022) 1392]

$$\text{ATLAS: } 4.6_{-2.6}^{+2.6} \text{ MeV}$$

[ATLAS-CONF-2022-068]



For enhanced light quark Yukawa couplings it does not work:

use instead kinematic properties of off-shell production

[works nicely also for other BSM scenarios see  
Haisch, Koole '21 '22]


# Kinematic discriminants

$$D_s^d = \log_{10} \left( \frac{P_{sig}^{d\bar{d}}}{P_{back}^{q\bar{q}} + P_{back}^{gg}} \right)$$

Poisson ratio of likelihoods

$$Z_i = \sqrt{2 \left[ (s_i + b_i) \ln \frac{(s_i + b_i)(b_i + \sigma_{b_i}^2)}{b_i^2 + (s_i + b_i)\sigma_{b_i}^2} - \frac{b_i^2}{\sigma_{b_i}^2} \ln \left( 1 + \frac{s_i \sigma_{b_i}^2}{b_i(b_i + \sigma_{b_i}^2)} \right) \right]}$$

$$\sigma_{b_i} = \Delta_{b_i} b_i$$



# Kinematic discriminants


$$D_s^d = \log_{10} \left( \frac{P_{sig}^{d\bar{d}}}{P_{back}^{q\bar{q}} + P_{back}^{gg}} \right)$$

Poisson ratio of likelihoods


$$Z_i = \sqrt{2 \left[ (s_i + b_i) \ln \frac{(s_i + b_i)(b_i + \sigma_{b_i}^2)}{b_i^2 + (s_i + b_i)\sigma_{b_i}^2} - \frac{b_i^2}{\sigma_{b_i}^2} \ln \left( 1 + \frac{s_i \sigma_{b_i}^2}{b_i(b_i + \sigma_{b_i}^2)} \right) \right]}$$

$$\sigma_{b_i} = \Delta_{b_i} b_i$$

Best probe of first  
generation quark  
Yukawa couplings!



# Conclusion



Higgs pair production: ML helps to improve on the sensitivity, possible to measure both trilinear and light quark Yukawa couplings at the HL-LHC


Off-shell Higgs: Kinematic discriminants extremely helpful to distinguish signal from background

Thanks for your attention!


Backup



# Invariant mass distribution




# SMEFT trilinear bounds




[Alasfar, RG, Grojean,  
Paul, Qian '22]

# Invariant mass distribution $ZZ$



# up Yukawa coupling



# EFT validity

