

## C<sub>5</sub>N<sup>-</sup> in collision with He: rotational transitions in the ISM

L. González-Sánchez,<sup>a</sup> A. Veselinova,<sup>a</sup> A. Martín Santa Daría,<sup>a</sup> F. A. Gianturco<sup>b</sup>


<sup>a</sup>Departamento de Química Física, University of Salamanca, Plaza de los Caídos sn, 37008 Salamanca, Spain

<sup>b</sup>Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck, A-6020 Innsbruck, Austria  
e-mail: lgonsan@usal.es

**Keywords:** inelastic collisions, computational chemistry, rotational excitations

In the last years, several C-bearing and (C,N)-bearing chains of molecular anions have been detected in the interstellar medium (ISM). Since experimental treatments are still challenging, computational methods have to be used to understand their chemistry. The C<sub>5</sub>N<sup>-</sup> anion is one of the largest (C,N)-bearing chains and, although there are several theoretical studies [1-3] that describe the dynamics of the smaller (CN<sup>-</sup>, C<sub>3</sub>N<sup>-</sup>) + He/H<sub>2</sub> systems, for the moment there has been no study reported for the collision between this molecular anion with He or H<sub>2</sub>. We have recently published our results [4].

We have performed quantum scattering calculations using a new ab initio potential energy surface (PES) where the interaction potential between C<sub>5</sub>N<sup>-</sup> and He was obtained using CCSD(T) approach and the complete basis set (CBS) limit (see Figure 1). Given the ISM conditions of this system, we calculate the state-to-state (de-)excitation cross sections and the respective rate coefficients as a function of temperature. These results have been also compared by those obtained for the C<sub>3</sub>N<sup>-</sup>/C<sub>5</sub>N<sup>-</sup> + H<sub>2</sub> system.



**Figure 1.** Potential energy contours for the C<sub>5</sub>N<sup>-</sup>/He system.

### References

- [1] L. González-Sánchez et al., *Rotationally Inelastic Collisions of CN<sup>-</sup> with He: Computing Cross Sections and Rates in the Interstellar Medium*. APJ **897** (2020) 75.
- [2] J. Klos and F. Lique, *First rate coefficients for an interstellar anion: application to the CN<sup>-</sup>-H<sub>2</sub> collisional system*. MNRAS **418** (2011) 271.
- [3] M. Lara-Moreno, T. Stoecklin, P. Halvick, *Rotational (de-)excitation of C<sub>3</sub>N<sup>-</sup> by collision with He atoms*. MNRAS **467** (2017) 4174.
- [4] R. Biswas, K. Giri, L. González-Sánchez, et al., *Rotational state-changes in C<sub>5</sub>N<sup>-</sup> by collisions with He and H<sub>2</sub>*. MNRAS **522** (2023) 5775.