

Heavy flavour production and decays at colliders



Mario Campanelli
University College London
African School of Physics 2022




African School of Fundamental
Physics and Applications













Outline

- What are heavy flavors and why they are interesting
- Discrete symmetries and CP violation
- CKM matrix and unitarity triangle
- Meson-antimeson oscillations
- b-production in ATLAS and CMS

What are heavy flavors?

- Matter comes in three generations of quarks and three of leptons, that we order in mass. A quark (or lepton) type (for quarks: u,d,c,s,t,b) is called flavor, to distinguish it from color.
- The heaviest quark is the top, so heavy that it decays before forming bound states. b and c are the heaviest to form mesons and baryons



	Quarks		Leptons	
Generation 3	 t Top	 b Bottom	 τ Tau	 ν_τ Tau-neutrino
Generation 2	 c Charm	 s Strange	 μ Muon	 ν_μ Muon-neutrino
Generation 1	 u Up	 d Down	 e Electron	 ν_e Electron-neutrino

Why interesting: Heavy Quark Effective Theory

- Quantum ChromoDynamics has an intrinsic scale, $\Lambda_{\text{QCD}} \sim 200 \text{ MeV}$, above which perturbative expansion can be applied, and below which (soft QCD) only empirical models can be used.
- For quark masses $m_Q \gg \Lambda_{\text{QCD}}$ Perturbative expansions can be used, and calculations easier
- For states with two heavy quarks (J/ψ , Y), Non-Relativistic QCD is used.
- No time to describe HQET here; refer to e.g. A.V. Manohar and M.B. Wise, Heavy Quark Physics, Cambridge University Press (2000)

Symmetries in Physics

- An operator can be applied to a Lagrangian representing a physical system; if the Lagrangian is invariant under this transformation, the operator corresponds to a conserved quantity (Noether's theorem).
- Ex. invariance of Lagrangian under translation
 $x \rightarrow x+a$ leads to momentum conservation
- If the Lagrangian is not conserved under an operator, the symmetry is broken, and the physics will be different. In some cases, symmetry breaking is subtle and can be treated as a perturbation

Discrete symmetries

Three discrete symmetries can be applied to a Lagrangian:

- Parity
- Charge conjugation
- Time reversal

In classical physics, all these symmetries are conserved at microscopic level; macroscopically, the concept of entropy breaks T-symmetry.

Things are more complicated in quantum mechanics

Parity: \mathcal{P}

- Reflection through a mirror, followed by a rotation of π around an axis defined by the mirror plane.
 - Space is isotropic, so we care if physics is invariant under a mirror reflection.



$$\mathbf{r} \rightarrow -\mathbf{r}$$

$$\mathbf{p} \rightarrow -\mathbf{p}$$

$$\mathbf{L} \rightarrow \mathbf{L}$$

- \mathcal{P} is violated in weak interactions:

$$[\mathcal{P}, \mathcal{H}_w] \neq 0$$

- Vectors change sign under a \mathcal{P} transformation, pseudo-vectors or axial-vectors do not.
- \mathcal{P} is a unitary operator: $\mathcal{P}^2=1$.


T. D. Lee & G. C. Wick Phys. Rev. **148** p1385 (1966) showed that there is no operator \mathcal{P} that adequately represents the parity operator in QM.

Charge Conjugation: C

- ◆ Change a quantum field ϕ into ϕ^\dagger , where ϕ^\dagger has opposite U(1) charges:
 - ◆ *baryon number, electric charge, lepton number, flavour quantum numbers like strangeness & beauty etc.*
- ◆ Change particle into antiparticle.
 - ◆ *the choice of particle and antiparticle is just a convention.*
- ◆ C is violated in weak interactions, so matter and antimatter behave differently, and:

$$[C, \mathcal{H}_w] \neq 0$$

- ◆ C is a unitary operator: $C^2=1$.



Combining Charge and Parity: CP

The fundamental point is that CP symmetry is broken in any theory that has complex coupling constants in the Lagrangian which cannot be removed by any choice of phase redefinition of the fields in the theory.


- Weak interactions are left-right asymmetric.
 - *It is not sufficient to consider \mathcal{C} and \mathcal{P} violation separately in order to distinguish between matter and antimatter.*
 - *i.e. if helicity is negative (left) or positive (right).*
- \mathcal{CP} is a unitary operator: $\mathcal{CP}^2=1$

Time Reversal: \mathcal{T}

Not to be confused with the classical consideration of the entropy of a macroscopic system.

- 'Flips the arrow of time'
 - *Reverse all time dependent quantities of a particle (momentum/spin).*
 - *Complex scalars (couplings) transform to their complex conjugate.*
 - *It is believed that weak decays violate \mathcal{T} , but EM interactions do not.*

- \mathcal{T} is an anti-unitary operator: $\mathcal{T}^2 = -1$.



Combining all symmetries: CPT

- All locally invariant Quantum Field Theories conserve CPT .¹
- CPT is anti-unitary: $CPT^2 = -1$.
- CPT can be violated by non-local theories like quantum gravity. These are hard to construct.
 - ⊙ *see work by Mavromatos, Ellis, Kostelecky etc. for more detail.*
- If CPT is conserved, a particle and its antiparticle will have
 - ⊙ *The same mass and lifetime .*
 - ⊙ *Symmetric electric charges.*
 - ⊙ *Opposite magnetic dipole moments (or gyromagnetic ratio for point-like leptons).*

Applying CP to physical states

$$CP | u \rangle = | \bar{u} \rangle$$

The u quark has $J^P = 1/2^+$, so the \mathcal{P} operator acting on u has an eigenvalue of $+1$. The \mathcal{C} operator changes particle to antiparticle.

$$CP | \pi^0 \rangle = - | \pi^0 \rangle$$

The π^0 has $J^{PC} = 0^{-+}$, so the minus sign comes from the parity operator acting on the π^0 meson. The \mathcal{C} operator changes particle to antiparticle. A π^0 is its own antiparticle.

$$CP | \pi^\pm \rangle = - | \pi^\mp \rangle$$

The π^\pm has $J^P = 0^-$, so the minus sign comes from the parity operator acting on the π meson. The \mathcal{C} operator changes the particle to antiparticle.

Flavour interactions in the SM: the CKM matrix

- In the SM Lagrangian, charged-current interactions, mediated by the W boson, allow interactions between U-like and D-like quarks

$$\mathcal{L}_{CC} = -\frac{g}{\sqrt{2}} \left(\bar{\tilde{U}}_L \gamma^\mu W_\mu^+ V \tilde{D}_L + \bar{\tilde{D}}_L \gamma^\mu W_\mu^- V^\dagger \tilde{U}_L \right).$$

- Where V is a non-diagonal mixing matrix

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$

How many parameters in the matrix?

in general, an $n \times n$ unitary matrix has n^2 real and independent parameters:

- ▶ a $n \times n$ matrix would have $2n^2$ parameters
- ▶ the unitary condition imposes n normalization constraints
- ▶ $n(n - 1)$ conditions from the orthogonality between each pair of columns:

thus $2n^2 - n - n(n - 1) = n^2$.

In the CKM matrix, not all of these parameters have a physical meaning:

- ▶ given n quark generations, $2n - 1$ phases can be absorbed by the freedom to select the phases of the quark fields
 - ▷ Each u , c or t phase allows for multiplying a row of the CKM matrix by a phase, while each d , s or b phase allows for multiplying a column by a phase.

thus: $n^2 - (2n - 1) = (n - 1)^2$.

Among the n^2 real independent parameters of a generic unitary matrix:

- ▶ $\frac{1}{2} n(n - 1)$ of these parameters can be associated to real rotation angles, so the number of independent phases in the CKM matrix case is:

$n^2 - \frac{1}{2} n(n - 1) - (2n - 1) = \frac{1}{2} (n - 1)(n - 2)$

$n(\text{families})$	Total indep. params. $(n - 1)^2$	Real rot. angles $\frac{1}{2}n(n - 1)$	Complex phase factors $\frac{1}{2}(n - 1)(n - 2)$
2	1	1	0
3	4	3	1
4	9	6	3

The matrix in terms of angles

- There are many ways of writing the matrix as a function of 4 parameters, but the most common (from PDG) uses 4 angles:

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}$$

- Where $c_{ij} = \cos q_{ij}$, $s_{ij} = \sin q_{ij}$, with all angles real
- The only complex part of the matrix comes from the $e^{i\delta}$ terms, and that is responsible for CP violation!
- Indeed, the CP swapping on the Lagrangian gives a different result if the matrix has one or more complex terms

Approximate parametrisation


- Experimentally, mixing is larger for nearby generations, $1 \gg \theta_{12} \gg \theta_{23} \gg \theta_{13}$. Wolfenstein expanded in $\lambda = \sin \theta_{12}$

$$V = \begin{pmatrix} 1 - \lambda^2/2 & +\lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \lambda^2/2 & +A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4)$$

- Diagonal terms ~ 1 , $V_{12}, V_{21} \sim \lambda$, $V_{23}, V_{32} \sim \lambda^2$, $V_{13}, V_{31} \sim \lambda^3$
- At second order, use $\rho = \rho (1 - \lambda^2/2)$, $\eta = \eta (1 - \lambda^2/2)$
- Now the complex component is only in V_{13} and V_{31} (third family!)

The CKM triangle

- The Wolfenstein parametrisation is graphically represented as a triangle with base at (0,0) and (1,0) and apex at (r,η).




Measuring triangle angles


$b \rightarrow c$ interfering with $b \rightarrow u$
 $B \rightarrow D^{(*)}K^{(*)}$
 $B^0 \rightarrow D^-K^0\pi^+$
 $B^0 \rightarrow D^{(*)}\pi$
 $B^0 \rightarrow D^{(*)}\rho$
 + charmless

$b \rightarrow u\bar{u}d$ $B \rightarrow a_1\pi$
 $B \rightarrow \pi\pi$ $B \rightarrow a_1\rho$
 $B \rightarrow \rho\pi$ $B \rightarrow b_1\pi$
 $B \rightarrow \rho\rho$ $B \rightarrow b_1\rho$
 $B \rightarrow a_1a_1$

$b \rightarrow c\bar{c}s$ $B \rightarrow J/\psi\pi^0$
 $B^0 \rightarrow J/\psi K_L^0$ $B \rightarrow D^{(*)+}D^{(*)-}$
 $B^0 \rightarrow J/\psi K_S^0$ $B \rightarrow \eta'K^0$
 $B^0 \rightarrow \psi(2S)K_S^0$ $B \rightarrow \rho K^0$
 $B^0 \rightarrow \chi_{1c}K_S^0$ $B \rightarrow \omega K^0$
 $B^0 \rightarrow \eta_c K_S^0$ $B \rightarrow \pi^0 K^0$
 $B^0 \rightarrow J/\psi K^{*0}$ $B \rightarrow \phi K^{(*)0}$
 $B \rightarrow KKK^0$
 $B \rightarrow f^0(980)K^0$



Measuring triangle sides



Current constraints on triangle (from www.pdg.org)






Figure 12.2: Constraints on the $\bar{\rho}$, $\bar{\eta}$ plane. The shaded areas have 99% CL.

- Several ways to independently measure sides and angles
- All point to a coherent picture: CP violation well understood in the SM

Producing B mesons

- The “clean” way: e^+e^- collisions at the $\Upsilon(4S)$ peak, followed by decay into B^+B^- or $B^0\bar{B}^0$ (Babar, BELLE)
- The “dirty” way: proton collisions followed by b-tagging (CDF, D0, LHCb, ATLAS, CMS)
 - Advantages: large rates, can produce B_s , B_c etc.
 - Disadvantages: large BG




Oscillations of neutral mesons in QM

- We have flavour eigenstates M^0 and \bar{M}^0 :
 - ⊙ M^0 can be K^0 (sd), D^0 (cu), B_d^0 (bd) or B_s^0 (bs)

flavour states \neq H_{eff} eigenstates:
 (defined flavour) (defined $m_{1,2}$ and $\Gamma_{1,2}$)

- if we consider only strong or electromagnetic interactions only, these flavour eigenstates would correspond to the physical ones
- However due to the weak interaction, the physical eigenstates are different from the flavour ones. This means that they can mix into each other:
 - ⊙ via short-distance or long-distance processes
- and then the flavour superposition decays

$$M = p M^0 \pm q \bar{M}^0$$



Schroedinger equation for oscillation

- We have flavour eigenstates M^0 and \bar{M}^0 :
 - ◎ M^0 can be K^0 (sd), D^0 (cu), B_d^0 (bd) or B_s^0 (bs)

flavour states \neq H_{eff} eigenstates:
(defined flavour) (defined $m_{1,2}$ and $\Gamma_{1,2}$)

- Time-dependent Schrödinger eqn. describes the evolution of the system:

$$i \frac{\partial}{\partial t} \begin{pmatrix} M^0 \\ \bar{M}^0 \end{pmatrix} = H \begin{pmatrix} M^0 \\ \bar{M}^0 \end{pmatrix} = \left(M - \frac{i}{2} \Gamma \right) \begin{pmatrix} M^0 \\ \bar{M}^0 \end{pmatrix}$$

- ◎ H is the hamiltonian; M and Γ are 2x2 hermitian matrices ($a_{ij} = \bar{a}_{ji}$)

$$M = \frac{1}{2} (H + H^\dagger) \text{ and } \Gamma = i(H - H^\dagger)$$

- CPT theorem: $M_{11} = M_{22}$ and $\Gamma_{11} = \Gamma_{22}$
 - ◎ particle and antiparticle have equal masses and lifetimes

Solutions for physical states

⊙ Physical states: eigenstates of effective Hamiltonian:

$$M_{S,L} \text{ (or } M_{L,H}) = p M^0 \pm q \bar{M}^0$$

label can be either S,L (short-, long-lived) or L,H (light, heavy) depending on values of Δm & $\Delta\Gamma$ (labels 1,2 usually reserved for CP eigenstates)

p & q complex coefficients that satisfy $|p|^2 + |q|^2 = 1$

● CP conserved if physical states = CP eigenstates ($|q/p| = 1$)

⊙ Eigenvalues (μ) and mass (Δm) and lifetime ($\Delta\Gamma$) differences can be derived with this formalism:

$$\mu_{L,H} = m_{L,H} - i/2 \Gamma_{L,H} = (M_{11} - i/2 \Gamma_{11}) \pm (q/p) (M_{12} - i/2 \Gamma_{12})$$

$$\Delta m = m_H - m_L \text{ and } \Delta\Gamma = \Gamma_H - \Gamma_L$$

$$(\Delta m)^2 - 1/4 (\Delta\Gamma)^2 = 4 (|M_{12}|^2 + 1/4 |\Gamma_{12}|^2)$$

$$\Delta m \Delta\Gamma = 4 \Re (M_{12} \Gamma_{12}^*)$$

$$(q/p)^2 = (M_{12}^* - i/2 \Gamma_{12}^*) / (M_{12} - i/2 \Gamma_{12})$$


other useful definitions:

$$x \equiv \Delta m / \Gamma$$




$$y \equiv \Delta\Gamma / 2\Gamma$$

Oscillation probability


- Bd and Bs oscillations formally identical, frequency very different



probab. to observe an initially produced X^0 as X^0 after time t
 probab. to observe an initially produced X^0 as \bar{X}^0 after time t




Oscillations in e^+e^- collisions




- $\beta\gamma = 0.56$ (BaBar)
= 0.425 (Belle)
- $t = t_1$ corresponds to the time that B_{TAG} decays.
- $t_2 - t_1 = \Delta t$

Bd oscillations

HFAG, <http://www.slac.stanford.edu/xorg/hfag/>



$$\frac{d\Gamma(B^0 \rightarrow f)/d\Delta t - d\Gamma(\bar{B}^0 \rightarrow f)/d\Delta t}{d\Gamma(B^0 \rightarrow f)/d\Delta t + d\Gamma(\bar{B}^0 \rightarrow f)/d\Delta t} = (1 - 2w) \cos(x\Delta t) \otimes R(\Delta t)$$



$$\Delta m_d = (0.507 \pm 0.005) \text{ ps}^{-1}$$

$$x = \Delta m_d \cdot \tau_{B_d} = 0.774 \pm 0.008$$

Bs oscillations

At the Tevatron on the B_s:

- amplitude method, instead of extracting directly Δm_s (*à la* LEP)


$$\frac{1}{|A_f|^2} \frac{d\Gamma(P^0(\bar{P}^0) \rightarrow f)}{d\Delta t} = [1 \pm A(1 - 2w) \cos(x\Delta t)] e^{-\Delta t}$$

- fit A at different values of Δm_s ;
if A=1
⇒ oscillations at this Δm_s value

Very precise determination
from the Tevatron:




$$\Delta m_s = (17.77 \pm 0.10 \pm 0.07) \text{ ps}^{-1}$$

$$x = \Delta m_s \cdot \tau_{B_s} = 25.5 \pm 0.6$$







ATLAS and CMS: production of heavy quark-antiquark systems

- For both ATLAS and CMS experiments, **dimuon decays** provide a particularly clean signature to trigger on in order to reconstruct **quarkonium states**




Prompt-non prompt quarkonia

- Also, to measure **prompt** and **non-prompt** yields simultaneously and disentangle the two contributions both **CMS & ATLAS** exploit a **2D mass and pseudo-proper time fit**.





Observations of bottomium systems




- The $\chi_b(3P)$ was observed by **ATLAS** in 2011 as a new structure in the $Y(1S)\gamma$ and $Y(2S)\gamma$ decay modes.

- LHCb** observed the $\chi_b(3P) \rightarrow Y(3S) \gamma$ decay channel.



- DØ** saw the $\chi_b(3P)$ in the $\chi_b(3P) \rightarrow Y(1S) \gamma$ decay channel.

- CMS** saw the $\chi_b(3P)$ in the $Y(1S)$, $Y(2S)$, and $Y(3S)$ radiative decays, in the 7 TeV data





Mesons with beauty and charm: B_c

- B^+ (B^-) is the b-quark meson with the largest production rate composed of $u\bar{b}$ ($\bar{u}b$). B_c^+ (B_c^-) meson is a ground state of $\bar{b}c$ ($b\bar{c}$) system and contains **two** heavy quarks of **different flavours** and its production is **then much rarer** [$\bar{b}b + \bar{c}c$]. CMS has reported the *inclusive* and *differential* (γ & p_T) $\sigma \cdot \mathcal{B}$

$$B_c^\pm \rightarrow J/\psi (\rightarrow \mu\mu) \pi^\pm \quad B^\pm \rightarrow J/\psi (\rightarrow \mu\mu) K^\pm$$

Theoretical prediction uncertainties up to 40%: renormalization, factorization scales and the m_b dependencies.

- Results from 4.77 fb^{-1} Run I pp collisions @ **7 TeV** : event selection based on displaced dimuon triggers.




Kinematic region


$p_T > 10 \text{ GeV}/c$ and $|y| < 1.5$ to maximize B_c^+ significance $[S/\sqrt{(S+B)}]$

S from Gaussian fit to MC
 [BCVEGPY $gg \rightarrow B_c + b + c$]
 B from $J/\psi \pi^\pm$ sidebands in data



B jets at the LHC



4 main production mechanisms




Request b-tagging




Use vertex mass and distance to distinguish bb, b, c and light quarks in jet


Invariant mass of two jets each identified as b




$g \rightarrow bb$ splitting with both b in a single jet



Study composition of jets by fitting separately each bin and compare to theory





Boosted and/or displaced topologies




Boosted $Z \rightarrow bb$ in a single jet

- To reduce BG, require production of a photon and a very energetic Z decaying into bb
- Being very energetic, the $Z \rightarrow bb$ is reconstructed as a single jet
- The jet must have:
 - a two-prong structure
 - two displaced vertices
- Z peak visible in the jet mass despite huge background



Boosted Higgs->bb




The Higgs coupling is proportional to the mass squared, so the largest decay probability is in 2 b-quarks final states

This decay mode suffers from large BG from b quarks produced in normal strong interactions (see above)


To reduce BG, search for b-jets produced in association with a W or Z boson (associate production)

Since signal/BG ratio improves with Higgs p_T , we only look at cases where the Higgs is **boosted**: its decay products are reconstructed in a single jet with a 2-prong sub-structure




Results for $H \rightarrow b\bar{b}$

- Events divided in 6 categories:
 - 0, 1 or two leptons
 - Medium- or high jet p_T



Background estimated from data and subtracted.
 Then the various categories are combined in an optimised way.
 First observation of the main decay mode of the Higgs boson!



Conclusions

- Heavy quarks a fundamental “laboratory” due to large mass (perturbative calculations) and long lifetime (reconstruction of secondary vertex)
- Also, CP-violating effects particularly relevant in third family
- Dedicated experiments (and accelerators!) built to extensively study the CKM matrix, all coherent with SM picture
- b and c production in jets studied in general-purpose LHC experiments
- The use of boosted topologies allowed observation of Z and H decays into b pairs despite a very large QCD background