


MSSM Higgs and Beyond




Marcela Carena

Theoretical Physics Department, Fermilab
Enrico Fermi Institute, University of Chicago

PHENO 2011 SYMPOSIUM
University of Wisconsin-Madison, May 9-11, 2011

Outline

- Introduction:
 - Higgs in the minimal SUSY model (MSSM)
 - Tevatron and LHC reach
- MSSM Higgs Extensions: A model-independent approach
 - The EFT at NLO
 - Masses and couplings
 - Collider phenomenology



The Higgs Sector in the MSSM

2 Higgs SU(2) doublets ϕ_1 and ϕ_2

$$\tan \beta = v_2 / v_1$$

→ 2 CP-even h, H with mixing angle α
1 CP-odd A and a charged pair H^\pm

$$\Rightarrow v = \sqrt{v_1^2 + v_2^2} = 246 \text{ GeV}$$

At tree level,

one Higgs doublet couples only to down quarks, the other couples only to up quarks

$$-L = \bar{\psi}_L^i \left(\hat{h}_d^{ij+} \phi_1 d_R^j + \hat{h}_u^{ij+} \phi_2 u_R^j \right) + h.c.$$

**Since the up and down sectors are diagonalized independently,
the Higgs interactions remain flavor diagonal at tree level.**

Couplings to gauge bosons & fermions (SM normalized)	$hZZ, hWW, ZHA, WH^\pm H$	→ $\sin(\beta - \alpha)$
	$HZZ, HWW, ZhA, WH^\pm h$	→ $\cos(\beta - \alpha)$
	$(h, H, A) u\bar{u} \rightarrow \cos \alpha / \sin \beta, \sin \alpha / \sin \beta, 1 / \tan \beta$ $(h, H, A) d\bar{d}/l^+ l^- \rightarrow -\sin \alpha / \cos \beta, \cos \alpha / \cos \beta, \tan \beta$	

Decoupling limit $m_A \gg m_Z$

Lightest (SM-like) Higgs $m_h \leq m_Z$, others heavy and roughly degenerate

Radiative Corrections to Higgs Boson Masses

Important quantum corrections due to incomplete cancellation of particles and superparticles in the loops

Main effects: stops; and sbottoms at large tan beta

$$m_h^2 = M_Z^2 \cos^2 2\beta + \frac{2g_2^2 m_t^4}{8\pi^2 M_W^2} \left[\ln(M_S^2/m_t^2) + \frac{X_t^2}{M_S^2} \left(1 - \frac{X_t^2}{12 M_S^2} \right) \right] + \text{h.o.}$$


$$M_S^2 = \frac{1}{2}(m_{t_1}^2 + m_{t_2}^2) \text{ and } X_t = A_t - \mu/\tan\beta \rightarrow \text{stop mixing}$$

- m_t^4 enhancement
- log sensitivity to stop masses M_S
- depend. on stop mass mixing X_t

2-loop corrections: $m_h \leq 135 \text{ GeV}$

$$M_S = 1 \rightarrow 2 \text{ TeV} \Rightarrow \Delta m_h \simeq 2 - 5 \text{ GeV}$$

Brignole, M.C., Degrandi, Diaz, Ellis, Haber, Hempfling, Heinemeyer, Hollik, Espinosa, Martin, Quiros, Ridolfi, Slavich, Wagner, Weiglein, Zhang, Zwirner, ...



Radiative Corrections to the Higgs Couplings

- 1) Important effects through radiative corrections to the CP-even mass matrix δM_{ij}^2 , which defines the mixing angle α

$$\sin \alpha \cos \alpha = M_{12}^2 / \sqrt{(\text{Tr } M^2)^2 - 4 \det M^2}$$

The off diagonal elements are prop. to

$$M_{12}^2 \propto -(m_A^2 + m_Z^2) \cos \beta \sin \beta + \frac{m_t^4}{16\pi^2 v^2} \frac{\mu X_t}{M_S^2} \left(\frac{X_t^2}{M_S^2} - 6 \right)$$

M.C. Mrenna, Wagner

Important effects of rad. correc. on $\sin \alpha$ or $\cos \alpha$ depending on the sign of μX_t and the magnitude of X_t / M_S and μ / M_S

==> govern couplings of Higgs to fermions and vector bosons

When off-diagonal elements vanish, either $\sin \alpha$ or $\cos \alpha$ vanish

==> strong suppression of the SM-like Higgs boson coupling to b-quarks and taus

Enhancement of BR ($h/H \rightarrow WW/\gamma\gamma$) for $m_{h/H} < 135 \text{ GeV}$

Radiative Corrections to the Higgs Couplings

2) Important vertex corrections to Higgs-fermion couplings from SUSY loops, relevant for large $\tan\beta$ (induce FCNC and CC effects)

$$g_{hbb} \approx \frac{-m_b \sin\alpha}{(1 + \Delta_b) v \cos\beta} (1 - \Delta_b / \tan\alpha \tan\beta)$$

destroy basic relation

$$g_{h,H,Abb} / g_{h,H,A\tau\tau} \propto m_b / m_\tau$$

$$g_{Hbb} \approx \frac{m_b \cos\alpha}{(1 + \Delta_b) v \cos\beta} (1 - \Delta_b \tan\alpha / \tan\beta)$$

$$\Delta_b = (\varepsilon_0^3 + \varepsilon_Y h_t^2) \tan\beta$$

$$g_{Abb} \approx \frac{m_b \tan\beta}{(1 + \Delta_b) v}$$

$$\varepsilon_0^i \approx \frac{2\alpha_s}{3\pi} \frac{\mu^* M_{\tilde{g}}^*}{\max[m_{\tilde{d}_1^i}^2, m_{\tilde{d}_2^i}^2, M_{\tilde{g}}^2]}$$

$$\varepsilon_Y \approx \frac{\mu^* A_t^*}{16\pi^2 \max[m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2, \mu^2]}$$

Strong suppression of $h(H)$ -bottom coupling

M.C. Mrenna, Wagner
 Haber, Herrero, Logan, Penaranda, Rigolin, Temes
 Noth, Spira: Muhlleitner, Rzehak, Spira

$$\tan\alpha \simeq \Delta_b / \tan\beta \rightarrow g_{hbb} \simeq 0; g_{h\tau\tau} \simeq \Delta_b m_\tau / v \quad (\text{Similar for } H)$$

SM-like Higgs decays into b - and tau-pairs can be drastically changed

Enhancement of BR ($h/H \rightarrow WW/\gamma\gamma$) for $m_{h/H} < 135 \text{ GeV}$

SM-like MSSM Higgs: SUSY benchmark scenarios

M.C., Heinemeyer, Wagner, Weiglein

- The m_h^{\max} scenario: [Maximizes m_h]

$$M_S = 1 \text{ TeV} ; \quad X_t = 2.4 M_S ; \quad m_{\tilde{g}} = 0.8 M_S ; \quad M_2 = -\mu = 200 \text{ GeV} ; \quad A_t = A_b$$

$$g_{hbb}, g_{h\tau\tau} \sim \sin \alpha_{\text{eff}} / \cos \beta \quad \textit{enhanced for low } m_A \text{ and intermediate to large } \tan \beta \text{ (analogous for H)}$$

hence, strong suppression of $\text{BR}(h \rightarrow \gamma\gamma)$ and $\text{BR}(h \rightarrow WW)$ with respect to SM

- The m_h^{\min} scenario: [zero mixing in the stop sector]

Similar coupling's behaviour as m_h^{\max} , but minimizes m_h .


- The small $\sin \alpha_{\text{eff}}$ scenario: (specially interesting for early phase LHC)

$$M_S = 800 \text{ GeV} ; \quad X_t = -1.2 \text{ TeV} ; \quad \mu = 2.5 M_S ; \quad m_{\tilde{g}} = M_2 = 500 \text{ GeV} ; \quad A_t = A_b$$


$$g_{hbb}, g_{h\tau\tau} \quad \textit{importantly suppressed for large } \tan \beta \text{ and small } m_A, \\ \text{and in different ways due to } \Delta_b \text{ corrections}$$

hence, $\text{BR}(h \rightarrow \gamma\gamma)$ and $\text{BR}(h \rightarrow WW)$ enhanced with respect to SM

SM-like Higgs production at the Tevatron



with $H \rightarrow b\bar{b}, WW^*$




with $H \rightarrow WW^{(*)}$.

*The Tevatron
projections for
a SM-like Higgs*

2011 and $\mathcal{L} = 10 \text{ fb}^{-1}$

~~(2014 and $\mathcal{L} = 16 \text{ fb}^{-1}$)~~




Draper


Tevatron reach for the MSSM SM-like Higgs

All channels included in CDF/DO combination.

The m_h^{\max} scenario: $m_h \sim 125$ GeV






The m_h^{\min} scenario: $m_h < 120$ GeV




The small $\sin \alpha_{eff}$: interesting coverage from $h \rightarrow WW$ for low mass range

Draper, Liu, Wagner + MC

SM-like Higgs at LHC




For a large region of parameter space
suppression of the $\gamma\gamma$ mode at the LHC




Suppression still sizable for m_A as large as 500 GeV

LHC reach for the MSSM SM-like Higgs

2×ATLAS 95%CL MSSM Higgs Reach
7 TeV, 5fb^{-1} , $\gamma\gamma+WW+\tau\tau+ZZ+bb$,
Max. Mixing




2×ATLAS 95%CL MSSM Higgs Reach
7 TeV, 5fb^{-1} , $\gamma\gamma+WW+\tau\tau+ZZ+bb$,
Min. Mixing



Important to improve on early LHC reach in tau tau mode

*Tevatron - early LHC combined reach :
MSSM SM-like Higgs*





3 sigma evidence of the SUSY Higgs responsible for EWSB

Non-Standard Higgs Production at the Tevatron and LHC

- Enhanced couplings to b quarks and tau-leptons
- Considering value of running bottom mass and 3 quark colors

$$BR(A \rightarrow b\bar{b}) \cong \frac{9}{9 + (1 + \Delta_b)^2}$$

$$BR(A \rightarrow \tau^+\tau^-) \cong \frac{(1 + \Delta_b)^2}{9 + (1 + \Delta_b)^2}$$



$$\sigma(b\bar{b}A) \times BR(A \rightarrow b\bar{b}) \cong \sigma(b\bar{b}A)_{SM} \times \frac{\tan\beta^2}{(1 + \Delta_b)^2} \times \frac{9}{(1 + \Delta_b)^2 + 9}$$

Strong dependence on the SUSY parameters in the bb channel.


$$\sigma(b\bar{b}, gg \rightarrow A) \times BR(A \rightarrow \tau\tau) \cong \sigma(b\bar{b}, gg \rightarrow A)_{SM} \times \frac{\tan\beta^2}{(1 + \Delta_b)^2 + 9}$$

Robust predictions in the tau-tau channel

Excellent coverage at both colliders in the di-tau inclusive channel

MSSM Higgs at the Tevatron and LHC

CDF + D0 combination: $A/H \rightarrow \text{di-taus}$





Limits are robust under variation
of SUSY parameter space

All channels combined:

H/A to tau pairs plus SM-like Higgs searches;

Significant coverage by Tevatron

LHC: almost all space explored at 3σ with $\sim 5 \text{ fb}^{-1}$.



LEP bounds on SM-like Higgs are in some tension with upper bound on m_h in the MSSM

Extensions of the MSSM Higgs Sector

- MSSM with Explicit CP violation
- Additional SM singlets
- Additional gauged U(1)'s
- Models with enhanced weak gauge symmetries
- Effective field theory with higher dimensional operators:
A more model-independent approach

More general MSSM Higgs extensions: EFT approach

- The non-minimal part of Higgs sector is parametrically heavier than the weak scale (understood as $v = 174$ GeV)

- SUSY breaking is of order v , hence heavy masses nearly supersymmetric

M : overall “heavy” scale **SUSY breaking mass splittings** $\Delta m \sim v \ll M$

In practice: formalism applies for e.g. $M \sim 1$ TeV

Low energy superpotential: at leading order in $1/M$

$$W = \mu H_u H_d + \frac{\omega_1}{2M} (H_u H_d)^2$$

- can include SUSY breaking via a spurion $X = m_s \theta^2$ $W_X \supset \alpha_1 \frac{\omega_1}{2M} X (H_u H_d)^2$

Only two new parameters: ω_1 and X

M.C, Kong, Ponton, Zurita

see also Dine, Seiberg, Thomas;
Antoniadis, Dudas, Ghilencea, Tziveloglou

With CP violation: Altmannshofer, M.C, Gori, de la Puente

- At NLO, Kähler potential only:

$$K = H_d^\dagger e^{2V} H_d + H_u^\dagger e^{2V} H_u + \Delta K^{\text{CV}} + \Delta K^{\text{Cust}}$$

Custodially violating (tree level) :

$$\Delta K^{\text{CV}} = \frac{c_1}{2|M|^2} (H_d^\dagger e^{2V} H_d)^2 + \frac{c_2}{2|M|^2} (H_u^\dagger e^{2V} H_u)^2 + \frac{c_3}{|M|^2} (H_u^\dagger e^{2V} H_u) (H_d^\dagger e^{2V} H_d)$$

Custodially preserving (tree level) :

$$\Delta K^{\text{Cust}} = \frac{c_4}{|M|^2} |H_u H_d|^2 + \left[\frac{c_6}{|M|^2} H_d^\dagger e^{2V} H_d + \frac{c_7}{|M|^2} H_u^\dagger e^{2V} H_u \right] (H_u H_d) + \text{h.c.}$$

Plus SUSY breaking terms obtained by multiplication by spurion, with new coefficients

$$X \rightarrow \gamma_i, \quad X^\dagger X \rightarrow \beta_i$$

- EFT coefficients can be essentially arbitrary, if UV theory complicated enough

Why to go beyond LO in the EFT approach

Quartic interactions of 2HDM can be written as

$$V \supset \frac{1}{2}\lambda_1(H_d^\dagger H_d)^2 + \frac{1}{2}\lambda_2(H_u^\dagger H_u)^2 + \lambda_3(H_u^\dagger H_u)(H_d^\dagger H_d) + \lambda_4(H_u H_d)(H_u^\dagger H_d^\dagger) \\ + \left\{ \frac{1}{2}\lambda_5(H_u H_d)^2 + \left[\lambda_6(H_d^\dagger H_d) + \lambda_7(H_u^\dagger H_u) \right] (H_u H_d) + \text{h.c.} \right\}$$

At $O(1/M)$, only $\lambda_5, \lambda_6, \lambda_7$ modified

At $O(1/M^2)$ all λ_i 's receive contributions

But at tree-level in MSSM: $\lambda_1, \lambda_2, \lambda_3, \lambda_4 \propto g^2$ (small)

**NLO effects can be relevant without indicating breakdown of EFT
(however, higher order effects should be small...)**


Higgs Spectra in EFT extensions of the MSSM

The lightest tree level Higgs mass can be well above the LEP bound!!.

Expansion parameters: μ/M and m_s/M (m_s is the spurion F term)

Second order terms can have a relevant impact.

Large deviations from the MSSM mass values, specially for low $\tan\beta$



Higgs Spectra in EFT extensions of the MSSM

M.C., Kong, Ponton, Zurita

The lightest tree level Higgs mass is well above M_Z .


Expansion parameters: μ/M and m_s/M

Second order terms can have a relevant impact.

Large deviations from the MSSM mass values,
specially for low $\tan\beta$

Scanning over model parameters

Scan: $|\omega_1|, |c_i| \in [0, 1]$ and $|\alpha_1|, |\beta_i|, |\gamma_i|, |\delta_i| \in [1/3, 1]$ for $i = 1, 2, 3, 4, 6, 7$



Lightest Higgs Mass after LEP and the Tevatron



GREEN → LEP excluded

MAGENTA → Tevatron excluded

[Higgsbounds: Bechtle, Brein, Heinemeyer, Weiglein, Williams]

RED → Tevatron with 10 fb⁻¹ and eff. = 1.5

BLUE → LHC




Most magenta and red regions at Tevatron reach in the $h \rightarrow WW$ channel


Heavy CP Even and Charged Higgs Masses

H and H^\pm follow MSSM trend (with m_A), but

- large spreading at smaller m_A (heavier H) **Multi-Higgs chain decays**
- non-negligible deviations throughout



Heavy CP-even Higgs



Charged Higgs

BMSSM Higgs at the Tevatron and LHC

- **At Tevatron:**

SM-like searches: 1) $h \rightarrow b\bar{b}$ 2) $h \rightarrow W\bar{W}$ 3) $H \rightarrow W\bar{W}$

Non-SM-like searches: A, H and h to tau pairs,

- **At the LHC:**

SM-like reach in di-photons, tau pairs and di-bosons

Non-SM-like Higgs boson in di-tau pairs or top-bottom and tau-neutrinos

Multi-Higgs chain decays

Benchmark Scenarios



Many benchmarks similar to MSSM ones, or with larger mass splitting in the A/H/H⁺ system (MSSM with ~~CP~~) → need to detect light new spectra.

Here, only examples of non-MSSM like scenarios, including ~~CP~~ BMSSM.

CP-even Higgs Bosons: low $\tan\beta$


Tevatron searches in the $h/H \rightarrow WW$ channel, ($h/H \rightarrow bb$ remains borderline)

Tevatron allowed (blue): good LHC coverage in $h/H \rightarrow WW/ZZ$ channels




Lightest Higgs Boson: low $\tan\beta$

- Important variations in the BR of h into bottom pairs



- In small regions of parameter space, **enhancement of order 2-5 in BR ($h \rightarrow \text{di-photons}$)** \rightarrow at Tevatron/LHC reach in the near future

Suppression of the hbb couplings: both for large and (unlike the MSSM) for intermediate/low tanb





Cancellation between tree level and h.o. operators contributions
yields enhancement in gluon fusion: lack of b-loops + light SUSY

For large tanb: enhanced hbb coupling as in the MSSM, when h is non-SM like



Enhancement of $h \rightarrow WW/ZZ$ and $h \rightarrow$ di-photon channels (also due to hbb coupling suppression)

$\tan \beta = 20$, $M = 1 \text{ TeV}$, $\mu = m_s = 200 \text{ GeV}$, $M_{\text{SUSY}} = 300 \text{ GeV}$, $A_t = A_b = 0$



Interesting reach in $h \rightarrow WW$ and in di-photon signals at the Tevatron and, of course, at the LHC

A-h inversion of hierarchy at low tan β



The MSSM channels $A/H \rightarrow hh$ and $H^\pm \rightarrow hW^\pm$
replaced by $h/H \rightarrow AA$ and $H^\pm \rightarrow AW^\pm$ in BMSSM
with parameter sets of BR's of order one

Benchmark point 1 (LHC signal)

Heavy h and H, non-SM like h in WW/ZZ channels at LHC

m_A (GeV)	m_h (GeV)	m_H (GeV)	m_{H^\pm} (GeV)
134	181	205	165
g_{hWW}^2	g_{HWW}^2	g_{hgg}^2	g_{Hgg}^2
0.03	0.95	0.79	0.99
channel	BMSSM (SM)	channel	BMSSM (SM)
$h \rightarrow b\bar{b}$	0.23 (0.005)	$h \rightarrow \tau\bar{\tau}$	0.03 (0.0005)
$h \rightarrow WW$	0.68 (0.92)	$h \rightarrow ZZ$	0.04 (0.07)
$H \rightarrow WW$	0.72 (0.73)	$H \rightarrow ZZ$	0.27 (0.27)
$A \rightarrow b\bar{b}$	0.89	$A \rightarrow \tau\bar{\tau}$	0.10
$H^\pm \rightarrow t\bar{b}$	0.57	$H^\pm \rightarrow \tau\nu_\tau$	0.40

$\tan\beta = 2$

- All Higgs CP-even Higgs masses well above the MSSM limit and $m_h > m_A$
- H very SM-like, first to be seen at LHC
- hWW and hZZ very suppressed, still $h \rightarrow ZZ/WW$ possible at LHC

Not such a heavy SM-like Higgs in the MSSM, specially with light SUSY

Benchmark point 2 (LHC signal)

No Tevatron reach. Two $ZZ \rightarrow 4$ lepton peaks at the LHC

m_A (GeV)	m_h (GeV)	m_H (GeV)	m_{H^\pm} (GeV)
184	204	234	203
g_{hWW}^2	g_{HWW}^2	g_{hgg}^2	g_{Hgg}^2
0.3	0.7	1.39	0.36
channel	BMSSM (SM)	channel	BMSSM (SM)
$h \rightarrow WW$	0.73 (0.72)	$h \rightarrow ZZ$	0.25 (0.27)
$H \rightarrow WW$	0.70 (0.71)	$H \rightarrow ZZ$	0.29 (0.29)
$A \rightarrow b\bar{b}$	0.87	$H^\pm \rightarrow t\bar{b}$	0.99

$\tan\beta = 2$

- All Masses in similar mass range and beyond LEP/Tevatron reach
- Lightest Higgs ~ 200 GeV
- $\text{BR}(h/H \rightarrow WW/ZZ) \sim \text{SM value}$ but hWW suppressed
- Any decay $H/A/H^\pm \rightarrow h X$ is closed due to heavy h

Two Higgs signals in the ZZ channel at LHC, both in the 200 GeV range

Benchmark point 3 (LHC signal)

Multi Higgs signal: chain decays

m_A (GeV)	m_h (GeV)	m_H (GeV)	m_{H^\pm} (GeV)
64	135	155	125
g_{hWW}^2	g_{HWW}^2	g_{hgg}^2	g_{Hgg}^2
0.002	0.991	0.65	1.17
channel	BMSSM	channel	BMSSM
$h \rightarrow b\bar{b}$	0.15	$h \rightarrow AA$	0.84
$H \rightarrow WW$	0.12	$H \rightarrow AA$	0.84
$H \rightarrow b\bar{b}$	0.02	$A \rightarrow b\bar{b}$	0.92
$H^\pm \rightarrow \tau\nu_\tau$	0.56	$H^\pm \rightarrow W^\pm + A$	0.40

$\tan\beta = 2$

- $h \rightarrow AA$ and $H \rightarrow AA$, with subsequent decays into di-taus + b pairs
- Also $gg \rightarrow H \rightarrow WW$ but needs large luminosity (about 100 fb^{-1})
- **$H^\pm \rightarrow A W^\pm$ possible**

Benchmark point 4 (LHC signal)

SM-like light higgs with enhanced di-photon signal

m_A (GeV)	m_h (GeV)	m_H (GeV)	m_{H^\pm} (GeV)
210	111.3	215	225
g_{hWW}^2	g_{HWW}^2	g_{hgg}^2	g_{Hgg}^2
0.98	0.02	1.39	0.84
channel	BMSSM (SM)	channel	BMSSM (SM)
$h \rightarrow b\bar{b}$	0.03 (0.79)	$h \rightarrow \gamma\gamma/10^{-3}$	12.1 (2.1)
$h \rightarrow \text{jets}$	0.56 (0.07)	$h \rightarrow WW$	0.36 (0.05)
$H \rightarrow b\bar{b}$	0.86	$H \rightarrow \tau\bar{\tau}$	0.14
$A \rightarrow b\bar{b}$	0.86	$A \rightarrow \tau\bar{\tau}$	0.14
$H^\pm \rightarrow \tau\nu_\tau$	0.35	$H^\pm \rightarrow t\bar{b}$	0.64

$\tan\beta = 20$

- strong suppression of $h \rightarrow b\bar{b}$ channel (escaped LEP bound)
- *Similar scenario with heavier A/H will allow $A/H \rightarrow hh$ decays*

Benchmark point 5 (LHC signal)

CP violation: All three Higgs decay dominantly to WW

Scenario I	H_1	H_2	H_3
M_{H_i} [GeV]	157	177	202
$\xi_{ZZH_i}^2$	0.94	0.04	0.02
$\xi_{ggH_i}^2$	0.72	0.62	0.47
$\text{BR}(H_i \rightarrow bb)$	15%	34%	24%
$\text{BR}(H_i \rightarrow WW)$	76%	58%	53%
$\text{BR}(H_i \rightarrow ZZ)$	6%	2%	19%

	Sc. I
$ \alpha $	1
$ \omega $	2
$\text{Arg}(\alpha)$	$\pi/2$
$\text{Arg}(\omega)$	$-\pi/10$
$\tan \beta$	2
M_{H^\pm} [GeV]	195
M [TeV]	2.5
μ [GeV]	160
m_S [GeV]	160

New CP phases allowed by EDM's

Large region of phases for similar but smaller mass values excluded by Tevatron

Heaviest Higgs can also be seen in ZZ channel

Outlook

Some type of SM-like Higgs is probably around the corner

*The Higgs sector can shed light to many SM puzzles
the origin of mass, flavor, dark matter ...*

*Many types of experiments are exploring the Higgs sector
impressive results from the Tevatron and LHC*

*The SM and many new physics models,
in particular SUSY Models, are being constrained*

Some corners of SUSY parameter space may be elusive

*But the Tevatron and ultimately the LHC will have the final
word on multi Higgs searches*

Higgs Spectra in EFT extensions of the MSSM


M.C., Kong, Ponton, Zurita

The lightest tree level Higgs mass is well above M_Z .

Expansion parameters: μ/M and m_s/M

Second order terms can have a relevant impact.

Smaller effects for large $\tan\beta$
main contributions
proportional to $1/M^2$.



Scanning over model parameters

Scan: $|\omega_1|, |c_i| \in [0, 1]$ and $|\alpha_1|, |\beta_i|, |\gamma_i|, |\delta_i| \in [1/3, 1]$ for $i = 1, 2, 3, 4, 6, 7$

Precision Electroweak Constraints

1. Tree-level effects due to new physics:

$$\alpha T^{\text{Tree}} = -\frac{v^2}{2M^2} \sin^4 \beta \left[c_2 - 2(\tan \beta)^{-2} c_3 + (\tan \beta)^{-4} c_1 \right]$$

2. Effects from MSSM Higgs sector:

- Heavier SM-like Higgs
 - Mass splittings among non-standard Higgses
- } Loop-level contr. to S and T

3. Custodially violating mass splittings in SUSY sector

Medina, Shah, Wagner

Here: require that $-0.4 < T^{\text{Tree}} + T^{\text{Higgs}} < 0.3$ (S is small)

Consistent with $-0.2 < T^{\text{Total}} < 0.3$ (95% *C.L.*) for $0 < T^{\text{SUSY}} < 0.2$

Examples

Example 1: singlets

$$W = \mu H_u H_d + \frac{1}{2} M_S S^2 + \lambda_S S H_u H_d - \underbrace{X \left(a_1 \mu H_u H_d + \frac{1}{2} a_2 M_S S^2 + a_3 \lambda_S S H_u H_d \right)}_{B_\mu\text{-term}}$$

$$K = H_u^\dagger e^V H_u + H_d^\dagger e^V H_d + S^\dagger S - \underbrace{X^\dagger X \left(b_1 H_d^\dagger H_d + b_2 H_u^\dagger H_u + b_3 S^\dagger S \right)}_{\text{Soft masses: } m_{H_d}^2, m_{H_u}^2, m_S^2}$$

Integrating out the singlet:

$$\begin{aligned} M &= M_S, & \omega_1 &= -\lambda_S^2, & \alpha_1 &= a_2 - 2a_3, \\ c_4 &= |\lambda_S|^2, & \gamma_4 &= a_2 - a_3, & \beta_4 &= |a_2 - a_3|^2 - b_3 \end{aligned}$$

Note $c_4 > 0$, other arbitrary

Example 2: triplets with $Y = \pm 1$

$$W \supset M_T T \bar{T} + \frac{1}{2} \lambda_T H_u T H_u + \frac{1}{2} \lambda_{\bar{T}} H_d \bar{T} H_d \\ + X \left(a_2 M_T T \bar{T} + \frac{1}{2} a_3 \lambda_T H_u T H_u + \frac{1}{2} a_4 \lambda_{\bar{T}} H_d \bar{T} H_d \right)$$

$$K \supset T^\dagger e^{2V} T + \bar{T}^\dagger e^{2V} \bar{T} + X X^\dagger (b_3 T^\dagger T + b_4 \bar{T}^\dagger \bar{T})$$

Integrating out the triplets:







$$\left. \begin{array}{lll} M = M_T, & \omega_1 = \frac{1}{4} \lambda_{\bar{T}}, & \alpha_1 = a_2 - a_3 - a_4, \\ c_1 = \frac{1}{4} |\lambda_{\bar{T}}|^2, & \gamma_1 = a_2 - a_4, & \beta_1 = |a_2 - a_4|^2 - b_3, \\ c_2 = \frac{1}{4} |\lambda_T|^2, & \gamma_2 = a_2 - a_3, & \beta_2 = |a_2 - a_3|^2 - b_4, \end{array} \right\} \begin{array}{l} \text{Induce custodially violating ops.} \\ \text{Note } c_1, c_2 > 0, \text{ other arbitrary} \\ (\Delta T < 0) \end{array}$$

For triplets with $Y = 0 \rightarrow \lambda_T H_u T H_d$

$$\left. \begin{array}{lll} M = M_T, & \omega_1 = -\frac{1}{4} \lambda_T^2, & \alpha_1 = a_2 - 2a_3, \\ c_3 = \frac{1}{2} |\lambda_T|^2, & \gamma_3 = a_2 - a_3, & \beta_3 = |a_2 - a_3|^2 - b_3, \\ c_4 = -\frac{1}{4} |\lambda_T|^2, & \gamma_4 = a_2 - a_3, & \beta_4 = |a_2 - a_3|^2 - b_3, \end{array} \right\} \begin{array}{l} \text{Induce custodially violating ops.} \\ \text{Note } c_3 > 0 \text{ } (\Delta T > 0), \\ \text{and } c_4 < 0! \end{array}$$





CP-even Higgs Couplings to gauge bosons and fermions

Variations of couplings
with respect to SM and
MSSM can lead to
important variations in
the production
processes and BR's
relevant for
Higgs searches



Gluon Fusion Production

A generic enhancement of the production for the Higgs that is SM-like (the one with largest coupling to WW/ZZ)




- The couplings of the CP-odd and charged Higgs bosons differ from the MSSM due to corrections to their kinetic terms only at order $1/M^2$
 → much less significant
- The main effects involving A and H^\pm are those related to new decay modes due to variations in the mass spectrum
- New decay channels such as $H \rightarrow AA/AZ$, $h \rightarrow AA$ and $H^\pm \rightarrow W^\pm A$ open with BR's of order one (low $\tan\beta$, A/h inversion)
- Regular MSSM channels with decays into h are closed at low $\tan\beta$ and open at large $\tan\beta$: $A \rightarrow hZ$; $H^\pm \rightarrow W^\pm h$; $H \rightarrow hh$
- At sufficiently large m_A (> 300 GeV) behavior similar to MSSM

MSSM Higgs at the Tevatron


All channels combined: H/A to tau pairs plus SM-like Higgs searches,
CDF + D0 projections for 10 fb⁻¹

Draper, Liu, Wagner

$$a_t = \sqrt{6} M_S, \mu = 200 \text{ GeV}, M_S = 1 \text{ TeV}$$



$$a_t = 0 \text{ GeV}, \mu = 200 \text{ GeV}, M_S = 2 \text{ TeV}$$



The Tevatron can explore a significant region of the MSSM Higgs parameter space

LHC rapidly becoming superior in tau-tau non SM-like Higgs channel.

LHC with all channels combined → almost all space explored at 3 σ with 5 fb⁻¹.

Benchmark point 1a (Tevatron signal)

Heavy Higgs SM-like, but Tevatron reach in $h \rightarrow WW$

m_A (GeV)	m_h (GeV)	m_H (GeV)	m_{H^\pm} (GeV)
135	174	186	164
g_{hWW}^2	g_{HWW}^2	g_{hgg}^2	g_{Hgg}^2
0.11	0.89	1.05	0.65
channel	BMSSM (SM)	channel	BMSSM (SM)
$h \rightarrow b\bar{b}$	0.12 (0.01)	$h \rightarrow WW$	0.84 (0.96)
$H \rightarrow WW$	0.81 (0.82)	$H \rightarrow ZZ$	0.17 (0.17)
$A \rightarrow b\bar{b}$	0.90	$A \rightarrow \tau\bar{\tau}$	0.10
$H^\pm \rightarrow \tau\nu_\tau$	0.59	$H^\pm \rightarrow t\bar{b}$	0.38

$\tan\beta = 2$

- All Higgs CP-even Higgs masses well above the MSSM limit and $m_h > m_A$
- hWW coupling very suppressed but still sizable $\text{BR}(h \rightarrow WW)$
- $H \rightarrow WW$ too heavy for the Tevatron, but good at LHC in $H \rightarrow ZZ \rightarrow 4\text{-leptons}$

Not such a heavy SM-like Higgs in the MSSM, specially with light SUSY