



Quarkonium measurements at Belle-II

Quarkonia as Tools 2023 (QaT2023+)

Pavel Oskin
(IJCLab)





KEKB and Belle



Collected data from 1999 to 2010:

- 121 fb^{-1} at $Y(5S) \sim 7.11 \times 10^6 B_s B_s$
- 711 fb^{-1} at $Y(4S) \sim 771 \times 10^6 BB$
- 3 fb^{-1} at $Y(3S)$
- 24 fb^{-1} at $Y(2S)$
- 6 fb^{-1} at $Y(1S)$
- 26 fb^{-1} scan above $Y(4S)$




SuperKEKB and Belle II

Beam current increased by **x2**

$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*}\right) \left(\frac{I_{\pm} \xi_{y\pm}}{\beta_y^*}\right) \left(\frac{R_L}{R_{\xi_y}}\right)$$

Vertical beta function at IP reduced by **1/20**
"Nano-beam" scheme

x40 instant luminosity increase





Collected data:

- 424 fb^{-1} at Y(4S)
- 19 fb^{-1} scan above Y(4S)

Goal:

- 50 ab^{-1} at Y(4S)



PTEP 2020 (2020) 2, 029201




Theoretical models

Hadronic molecule

Compound state of two hadrons. The most promising model.

The charmonium-like states can be described as a mixture of pure charmonium and a molecular component:


$$X(3872) = C_1 \cdot |c\bar{c}\rangle + C_2 \cdot |D^0 \bar{D}^{0*}\rangle$$



Quarkonium measurements at Belle-II / Pavel Oskin / QaT 2023


Hybrid

Conventional quark-antiquark mesons with excited gluon degrees of freedom.



Compact tetraquark

States containing four constituent quarks irrespective of their clustering.



Hadroquarkonium





Compact quarkonium core surrounded by an excited light-quark cloud.



Charmonium production at Belle II

At Belle / Belle II charmonium events comes “for free”:




- We don't need to develop special triggers.
- We don't need to tune accelerator.
- Uniquely clean environment compare to pp production.



Unique for Belle II !

B decays

- $B \rightarrow K X_{cc^-}$ is CKM favoured process with large branching fractions $\sim 10^{-3} - 10^{-4}$.
- Due to the known number of B pairs, it is possible to measure absolute BF[$B \rightarrow X_{cc^-} K$].
- Access to X(3872) lineshape with $D^0 \bar{D}^0 \pi^0$ channel.
- Access to unknown / hardly reconstructable modes with the B-tagging technique.





Phys. Rev. D 84, 052004 (2011)

Charmonium + B-tagging

PRD 97, 012005 (2018)

Precise measurement of the e^+e^- energy allows to observe mass peaks with partial reconstruction:


$$M_{X_{cc}} = \sqrt{p_{e^+e^-} - p_{B_{tag}} - p_K}$$




Mode	Yield	Significance (σ)	$\epsilon (10^{-3})$	$\mathcal{B} (10^{-4})$	World average for $\mathcal{B} (10^{-4})$ [10]
η_c	2590 ± 180	14.2	2.73 ± 0.02	$12.0 \pm 0.8 \pm 0.7$	9.6 ± 1.1
J/ψ	1860 ± 140	13.7	2.65 ± 0.02	$8.9 \pm 0.6 \pm 0.5$	10.26 ± 0.031
χ_{c0}	430 ± 190	2.2	2.67 ± 0.02	$2.0 \pm 0.9 \pm 0.1 (<3.3)$	$1.50^{+0.15}_{-0.14}$
χ_{c1}	1230 ± 180	6.8	2.68 ± 0.02	$5.8 \pm 0.9 \pm 0.5$	4.79 ± 0.23
$\eta_c(2S)$	1050 ± 240	4.1	2.77 ± 0.02	$4.8 \pm 1.1 \pm 0.3$	3.4 ± 1.8
$\psi(2S)$	1410 ± 210	6.6	2.79 ± 0.02	$6.4 \pm 1.0 \pm 0.4$	6.26 ± 0.24
$\psi(3770)$	-40 ± 310	-	2.76 ± 0.02	$-0.2 \pm 1.4 \pm 0.0 (<2.3)$	4.9 ± 1.3
$X(3872)$	260 ± 230	1.1	2.79 ± 0.01	$1.2 \pm 1.1 \pm 0.1 (<2.6)$	(<3.2)
$X(3915)$	80 ± 350	0.3	2.79 ± 0.01	$0.4 \pm 1.6 \pm 0.0 (<2.8)$	-

Initial state radiation

- 50 ab^{-1} data corresponds to 2000-2800 pb^{-1} /10 MeV at 4-5 GeV, which is compatible with BES III (500 pb^{-1} /10 MeV, but higher efficiency).
- Access to 1^{--} states with masses > 4.6 GeV.
- Effective luminosity and detection efficiency are relatively low.
- c-baryons ($\Lambda_c^+ \Sigma_c^-, \Sigma_c^+ \Sigma_c^-$) and cs-mesons pairs production ($D_s^+ D_{s2}^- (2573), D_s^- D_{s0}^* (2317)$).




Golden Channels	$E_{c.m.}$ (GeV)	Statistical error (%)	Related XYZ states
$\pi^+ \pi^- J/\psi$	4.23	7.5 (3.0)	$Y(4008), Y(4260), Z_c(3900)$
$\pi^+ \pi^- \psi(2S)$	4.36	12 (5.0)	$Y(4260), Y(4360), Y(4660), Z_c(4050)$
$K^+ K^- J/\psi$	4.53	15 (6.5)	Z_{cs}
$\pi^+ \pi^- h_c$	4.23	15 (6.5)	$Y(4220), Y(4390), Z_c(4020), Z_c(4025)$
$\omega \chi_{c0}$	4.23	35 (15)	$Y(4220)$
		10 ab^{-1} 50 ab^{-1}	




PTEP 2020 (2020) 2, 029201

Two-photon production

- Access to a wide range of quantum numbers. $J/\psi\omega$ and $J/\psi\phi$ in particular.
- Provides a better S/B ratio compare to B-decays.
- Allows to measure the Q^2 dependence of $c\bar{c}$ production.




B-decay



PRL 94, 182002 (2005)




Two photon



PRD 86, 072002 (2012)




Double charmonium production

- All observed processes are of the type $e^+e^- \rightarrow c\bar{c} (J=1) c\bar{c} (J=0)$. Belle II will allow to study $e^+e^- \rightarrow \eta_c X_{(c\bar{c})}$ and $e^+e^- \rightarrow \chi_{c0} X_{(c\bar{c})}$ processes.
- $e^+e^- \rightarrow \psi(2S) X_{(c\bar{c})}$ can be studied at Belle II with larger statistics.
- Large cross section revealed importance of the next order corrections in NRQCD.




Double charmonium production

- All observed processes are of the type $e^+e^- \rightarrow c\bar{c} (J=1) c\bar{c} (J=0)$. Belle II will allow to study $e^+e^- \rightarrow \eta_c X_{(c\bar{c})}$ and $e^+e^- \rightarrow \chi_{c0} X_{(c\bar{c})}$ processes.
- $e^+e^- \rightarrow \psi(2S) X_{(c\bar{c})}$ can be studied at Belle II with larger statistics.
- Large cross section revealed importance of the next order corrections in NRQCD.



Bottomonium



We have two type of states:

- Below $B\bar{B}$ threshold
- Above $B\bar{B}$ threshold


The states below $B\bar{B}$ mesons threshold are well described by the potential models.

Conventional bottomonium (pure bb states)

Bottomonium-like states (mixture of bb and $B\bar{B}$)

Purely exotic states

Bottomonium



Process	Γ , MeV
$\Upsilon(5S) \rightarrow \Upsilon(1S)\eta$	0.039 ± 0.011
$\Upsilon(5S) \rightarrow \Upsilon(2S)\eta$	0.204 ± 0.44
$\Upsilon(4S) \rightarrow \Upsilon(1S)\eta$	0.004 ± 0.0008
$\Upsilon(4S) \rightarrow h_b(1P)\eta$	0.045 ± 0.007
$\Upsilon(3S) \rightarrow \Upsilon(1S)\eta$	$< 0.002 \cdot 10^{-3}$
$\Upsilon(2S) \rightarrow \Upsilon(1S)\eta$	$(0.0093 \pm 0.0015) \cdot 10^{-3}$


Process	Γ , MeV
$\Upsilon(5S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	$0.59 \pm 0.04 \pm 0.09$
$\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^+\pi^-$	$0.85 \pm 0.07 \pm 0.16$
$\Upsilon(5S) \rightarrow \Upsilon(3S)\pi^+\pi^-$	$0.52^{+0.20}_{-0.17} \pm 0.10$
$\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.0060 ± 0.0005
$\Upsilon(3S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.0009 ± 0.00008
$\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.0019 ± 0.0002

Conventional bottomonium (pure bb states)
 Bottomonium-like states (mixture of bb and BB)
 Purely exotic states

PRL 100, 112001 (2008)

Bottomonium


Mod. Phys. Lett. A 32, No04, 1750025



PRL 108 (2012) 122001

Process	Γ , MeV
$\Upsilon(5S) \rightarrow \Upsilon(1S)\eta$	0.039 ± 0.011
$\Upsilon(5S) \rightarrow \Upsilon(2S)\eta$	0.204 ± 0.44
$\Upsilon(4S) \rightarrow \Upsilon(1S)\eta$	0.004 ± 0.0008
$\Upsilon(4S) \rightarrow h_b(1P)\eta$	0.045 ± 0.007
$\Upsilon(3S) \rightarrow \Upsilon(1S)\eta$	$< 0.002 \cdot 10^{-3}$
$\Upsilon(2S) \rightarrow \Upsilon(1S)\eta$	$(0.0093 \pm 0.0015) \cdot 10^{-3}$

Process	Γ , MeV
$\Upsilon(5S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	$0.59 \pm 0.04 \pm 0.09$
$\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^+\pi^-$	$0.85 \pm 0.07 \pm 0.16$
$\Upsilon(5S) \rightarrow \Upsilon(3S)\pi^+\pi^-$	$0.52^{+0.20}_{-0.17} \pm 0.10$
$\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.0060 ± 0.0005
$\Upsilon(3S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.0009 ± 0.00008
$\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	0.0019 ± 0.0002




PRL 100, 112001 (2008)

Proceed via intermediate exotic state
 $Y(5S) \rightarrow [Z_b^+ \rightarrow Y(nS)\pi^+]\pi^-$

Hadronic molecule

Compound state of two hadrons. The most promising model. The bottomonium-like states can be described as a mixture of pure bottomonium and a molecular component:




$$Y(10860) = C_1 \cdot |bb\rangle + C_2 \cdot |B_{(s)}^{((*)*)} B_{(s)}^{((*)*)}\rangle$$

Z_b decay mode	Branching fraction
$Z_b^+(10610) \rightarrow \Upsilon(nS)/h_b(mP)\pi^+$	$14.4_{-1.9}^{+2.5}\%$
$Z_b^+(10610) \rightarrow B^+\bar{B}^{*0}/B^+\bar{B}^0$	$85.6_{-2.9}^{+2.1}\%$
$Z_b^+(10650) \rightarrow \Upsilon(nS)/h_b(mP)\pi^+$	$26.6_{-4.7}^{+5.0}\%$
$Z_b^+(10650) \rightarrow B^{*+}\bar{B}^{*0}$	$74_{-6}^{+4}\%$

PRL, 108, 122001 (2012)

Hybrid

Conventional quark-antiquark mesons with excited gluon degrees of freedom.



Compact tetraquark

States containing four constituent quarks irrespective of their clustering.




Hadroquarkonium

Compact quarkonium core surrounded by an excited light-quark cloud.



Bottomonium production

Prompt production




Direct production in e^+e^- collisions






- Provides a unique clean environment (we have only bottomonium in the event);
- Precise measurement of the beam energy gives access to “unreconstructable” particles and decay modes;
- Allows tuning CM energy
- **Only $Y(nS)$ states can be produced with quantum numbers of the photon 1^-**
- Other quantum numbers can be obtained via hadronic or radiative transitions from 1^- states;

gives access to exotic states above $Y(4S)$



Bottomonium production



- Provides a unique clean environment (we have only bottomonium in the event);
- Precise measurement of the beam energy gives access to “unreconstructable” particles and decay modes;
- Allows tuning CM energy
- **Only $Y(nS)$ states can be produced with quantum numbers of the photon 1^-**
- Other quantum numbers can be obtained via hadronic or radiative transitions from 1^- states;



Missing mass technique for bottomonium

Precise measurement of the e^+e^- energy allows to observe mass peaks with partial reconstruction:

$\pi^+ \pi^-$ missing mass has better resolution than $M(\mu^+ \mu^-)$

$$M_{\text{miss}}(\pi\pi) = \sqrt{(E_{\text{c.m.}} - E_{\pi\pi}^*)^2 - p_{\pi\pi}^{*2}}$$

$\Upsilon(5S) \rightarrow X_{b\bar{b}} \pi^+ \pi^-$ transitions:



PRL 108 (2012) 032001




Very important for unreconstructable bottomonium states $\eta_b(nS)$, $h_b(nP)$

Discovery of the $Y(10753)$

$Y(10753)$ state was observed in the $e^+e^- \rightarrow Y(nS) \pi^+ \pi^-$ ($n = 1, 2, 3$) cross section energy dependence.



JHEP 10 (2019) 220

measurement at $\sqrt{s} = 10.864$ GeV with $L = 47.647$ fb $^{-1}$:






Belle II energy scan above $Y(4S)$

- Unique data provide an opportunity to study $Y(10753)$ in different final states and understand its nature.
- Scan above $Y(4S)$ has a good potential for early physics impact by Belle II even with small statistics.




Observation of $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$ at $\sqrt{s} = 10.745$ GeV

- Similar to Y(10753) structure named Y(4220) was observed in $e^+e^- \rightarrow J/\psi \pi^+ \pi^-$ cross section dependence by BES III (PRL 118, 092001 (2017)).
- BES III also observed the Y(4220) peak in $\gamma X(3872)$ and $\omega\chi_{c0}$ final states (PRL, 122, 232002 (2019), PRD 99, 091103(R) (2019)).
- $\omega\chi_{b1,2}$ production was found to be enhanced near Y(5S) (PRL 113, 142001 (2014)).
- We expect Y(10753) to decay into $\gamma[X_b \rightarrow \omega Y(1S)]$ and $\omega\chi_{bJ}$ final states.




Observation of $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$ at $\sqrt{s} = 10.745$ GeV





Strongly enhanced at 10.745 GeV.

$$\sigma(e^+e^- \rightarrow \omega\chi_{b1}) = 3.6 \pm 0.7 \pm 0.5 \text{ pb}$$

$$\sigma(e^+e^- \rightarrow \omega\chi_{b2}) = 2.8_{-1.0}^{+1.2} \pm 0.4 \text{ pb}$$



No evidence of X_b ($b\bar{b}$ partner of $X(3872)$) state. These are reflections of $\omega\chi_{bJ}(1P)$.




arXiv:2208.13189




$B\bar{B}$ decomposition with B-tagging

Study the energy dependence of the $B\bar{B}$ pairs production.

- B-tagging can be used to measure the $B_{(s)}^{(*)} \bar{B}_{(s)}^{(*)}$ cross section energy dependence.
- Yet another unique way to study bottomonium at Belle / Belle II.
- A good probe for bottomonium models (especially the molecular model).



$$M_{bc} = \sqrt{(E_{cm}/2)^2 - p_B^2}$$



JHEP 06 (2021) 137

Belle II potential charmonium

The main source of improvements and new results is the increase in statistics.

- Measure double charmonium production cross sections with $e^+e^- \rightarrow \psi(2S)X_{(c\bar{c})}$ and search for $e^+e^- \rightarrow \eta_c X_{(c\bar{c})}$ and $e^+e^- \rightarrow \chi_{c0} X_{(c\bar{c})}$.
- Measure $X(3872)$ lineshape with $D^0\bar{D}^0\pi^0$ channel.
- Study high energy region ($E > 4.6$ GeV) unapproachable for BES III with ISR production.
- Search for new charmonium-like states or new decay modes. Confirm the states and transitions obtained with low statistics. Measure quantum numbers of observed charmonium-like states.
- Search for new charmonium state with improved B-tagging method.
- Larger statistics will allow to measure $p_{c\bar{c}}^*$, Q^2 production dependence.

Belle II potential bottomonium

Scan above $Y(4S)$ gives an opportunity for a lot of unique studies:

- $Y(10753)$ decays to different exclusive and inclusive final states. Study of its properties;
- Energy dependence of the various final state cross sections;
- $B\bar{B}$ decomposition and its cross section dependence on CM energy;

Wide range of long-term non- $Y(4S)$ possibilities:

- Increase the above- $Y(4S)$ scan statistics;
- $Y(6S)$ region study with high statistics after accelerator upgrade;

Golden Modes
$e^+e^- \rightarrow \pi^+\pi^-\Upsilon(pS)(\rightarrow \ell^+\ell^-)$
$B\bar{B}$ decomposition
$\pi^+\pi^-$ Dalitz
$Y_b \rightarrow \omega\eta_b(1S)$
$Y_b \rightarrow \omega\chi_{bJ}(1P)$
Silver Modes
$Y_b \rightarrow \pi^+\pi^-X$ (inclusive)
$Y_b \rightarrow \eta X$ (inclusive)
$Y_b \rightarrow \eta\Upsilon(1S, 2S)(\rightarrow \ell^+\ell^-)$
$Y_b \rightarrow \eta'\Upsilon(1S)(\rightarrow \ell^+\ell^-)$
$Y_b \rightarrow \Upsilon(1S)$ (inclusive)
Bronze Modes
$Y_b \rightarrow \gamma X_b$
$Y_b \rightarrow \pi^0\pi^0\Upsilon(pS)(\rightarrow \ell^+\ell^-)$
$Y_b \rightarrow KK(\phi)\Upsilon(pS)(\rightarrow \ell^+\ell^-)$
$Y_b \rightarrow \pi^0\pi^0X$ (inclusive)
$Y_b \rightarrow \pi^0X$ (incl. or excl.)
...

Backup

X(3872) lineshape with $D^0\bar{D}^0\pi^0$ channel


Γ_{tot}
< 1.2 MeV (90% C.L.)

<



Mass resolution
 $1.86 \pm 0.01 \text{ MeV}/c^2$

Phys. Rev. D 84, 052004 (2011)

- Improvement of mass resolution is more essential than that of statistics.
- We can use channel with good mass resolution:
 - Mass resolution: $684 \pm 8 \text{ keV}$
 - Signal yield with 1 ab^{-1} : 64.5 ± 23.9



Comes from large error of
 $\text{Br}(B^\pm \rightarrow K^\pm X(3872))$
 $\times \text{Br}(X(3872) \rightarrow D^0\bar{D}^0\pi^0)$



(Assuming a Breit-Wigner shape)

B decays


S - seen (observed)

NS - not seen or excluded

N - not performed (in a given mass range)

MF - missing fit (fit have not been extended to this mass range)

– no search has been performed in this mode



State	J^{PC}	$\psi\pi\pi$	$\psi\omega$	$\psi\gamma$	$\psi\phi$	$\psi\eta$	$\psi'\pi\pi$	$\psi'\omega$	$\psi'\gamma$	$\chi_{c\gamma}$	$p\bar{p}$	$\Lambda\bar{\Lambda}$	$\Lambda_c\bar{\Lambda}_c$	$D\bar{D}$	$D\bar{D}^*$	$2D^*$	$2D_s^{(*)}$	$\gamma\gamma$
X(3872)	1^{++}	S	S	S	—	NS	—	—	S	NS	MF	MF	—	—	S	—	—	NS
Y(3940)	J^{P+}	MF	S	NS	—	—	—	—	MF	—	MF	MF	—	MF	NS	—	N	N
Z(3930)	2^{++}	MF	MF	NS	—	—	—	—	MF	—	MF	MF	—	MF	MF	—	N	N
Y(4140)	J^{P+}	MF	MF	N	S	—	N	—	N	—	MF	MF	—	MF	N	N	N	N
X(4160)	0^{P+}	MF	MF	N	MF	—	N	—	N	—	MF	MF	—	MF	N	N	N	N
Y(4260)	1^{--}	NS	—	—	—	MF	N	—	—	N	MF	MF	—	N	N	N	N	—
X(4350)	J^{P+}	MF	MF	N	MF	—	N	N	N	—	MF	MF	—	N	N	N	N	N
Y(4350)	1^{--}	MF	—	—	—	MF	N	—	—	N	MF	MF	—	N	N	N	N	—
Y(4660)	1^{--}	N	—	—	—	MF	N	—	—	N	MF	MF	MF	N	N	N	N	—

Initial state radiation


S - seen (observed)

NS - not seen or excluded

N - not performed (in a given mass range)

MF - missing fit (fit have not been extended to this mass range)

– no search has been performed in this mode



State	J^{PC}	$\psi\pi\pi$	$\psi'\pi\pi$	$\psi\eta$	$\chi_{c\gamma}$	$p\bar{p}$	$\Lambda\bar{\Lambda}$	$\Lambda_c\bar{\Lambda}_c$	$D\bar{D}$	$D\bar{D}^*$	$2D^*$	$2D_s^{(*)}$
Y(4260)	1^{--}	S	NS	NS	NS	NS	MF	—	NS	NS	NS	NS
Y(4350)	1^{--}	NS	S	MF	MF	MF	MF	—	MF	MF	MF	MF
Y(4660)	1^{--}	NS	S	MF	MF	MF	MF	S	MF	MF	MF	MF

Two-photon production


S - seen (observed)

NS - not seen or excluded

N - not performed (in a given mass range)

MF - missing fit (fit have not been extended to this mass range)

– no search has been performed in this mode



State	J^{PC}	$\psi\pi\pi$	$\psi\omega$	$\psi\gamma$	$\psi\phi$	$\psi'\pi\pi$	$\psi'\omega$	$\psi'\gamma$	$p\bar{p}$	$\Lambda\bar{\Lambda}$	$\Lambda_c\bar{\Lambda}_c$	$D\bar{D}$	$D\bar{D}^*$	$2D^*$	$2D_s^{(*)}$
X(3872)	1^{++}	N	hard	hard	—	—	—	hard	MF	MF	—	MF	N	—	—
Y(3915)	0^{++}	N	S	hard	—	—	—	hard	MF	MF	—	MF	N	—	N
Z(3930)	2^{++}	N	MF	hard	—	—	—	hard	MF	MF	—	S	N	—	N
Y(4140)	J^{P+}	N	MF	hard	NS	N	—	hard	N	N	—	MF	N	N	N
X(4160)	0^{P+}	N	MF	hard	NS	N	—	hard	N	N	—	MF	N	N	N
X(4350)	J^{P+}	N	N	hard	S	N	N	hard	N	N	N	N	N	N	N

Double charmonium production


S - seen (observed)

NS - not seen or excluded

N - not performed (in a given mass range)

MF - missing fit (fit have not been extended to this mass range)

– no search has been performed in this mode



State	J^{PC}	$\psi\pi\pi$	$\psi\omega$	$\psi\gamma$	$\psi\phi$	$\psi'\pi\pi$	$\psi'\omega$	$\psi'\gamma$	$\chi_{c\gamma}$	$p\bar{p}$	$\Lambda\bar{\Lambda}$	$\Lambda_c\bar{\Lambda}_c$	$D\bar{D}$	$D\bar{D}^*$	$2D^*$
X(3872)	1^{++}	hard	N	hard	—	hard	—	hard	hard	hard	hard	—	MF	MF	—
X(3940)	0^{-+}	hard	N	hard	—	hard	—	hard	hard	hard	hard	—	NS	S	—
Z(3930)	2^{++}	hard	N	hard	—	hard	—	hard	hard	hard	hard	—	MF	MF	—
Y(4140)	J^{P+}	hard	N	hard	N	hard	—	hard	hard	hard	hard	—	MF	MF	MF
X(4160)	0^{P+}	hard	N	hard	N	hard	—	hard	hard	hard	hard	—	MF	S	MF
X(4350)	J^{P+}	hard	N	hard	N	hard	N	hard	hard	hard	hard	hard	MF	MF	MF