



# Goldstino condensation

Fotis Farakos

University of Padova

@ SUSY '23, Southampton

## Plan:

→ Motivation

→ Goldstino condensation at large  $N$

→ Outlook

## *Motivation*

Dark Energy in String Theory/Supergravity means

$$\rho_{DE} \sim V_{4D} = f^2 - 3m_{3/2}^2 > 0 \implies \text{Supersymmetry Breaking.}$$

We will focus on non-linear SUSY because:

1. NL-SUSY underlines many **EFTs with broken SUSY**.

*See e.g. Wess, Bagger '92, Dudas, Dall'Agata, FF '16,  
Dall'Agata, FF, Cribiori '17*

2. In specific **“anti-brane uplifts”** the supersymmetry breaking is described by sectors with non-linear supersymmetry.

*See e.g. Kallosh, Wrase '14, Bergshoeff, Dasgupta, Kallosh,  
Van Proeyen, Wrase '15, Dasgupta, Emelin and McDonough '16*

## Goldstino: An indispensable part of uplifts

Salient features of NL-SUSY in Volkov–Akulov model ('73)

- ▶ The (N=1) goldstino Lagrangian

$$\mathcal{L}_{VA} = -f^2 - i\bar{G}\not{\partial}G + \frac{1}{4f^2}\bar{G}^2\partial^2G^2 - \frac{1}{16f^6}G^2\bar{G}^2\partial^2G^2\partial^2\bar{G}^2,$$

and generates the **uplift** when coupled to SG.

- ▶ The non-linear SUSY is

$$\delta G_\alpha = -f\xi_\alpha - (i/2f)\not{\partial}_{\alpha\dot{\alpha}}G^2\bar{\xi}^{\dot{\alpha}} + \dots$$


We want to understand if the **non-linear terms** are spectators or they have any physical significance/impact.

## Goldstino condensation: A threat to uplifts

- ▶ If there is a condensate of the form

$$\langle i\bar{G}\not{\partial}G \rangle \neq 0,$$

then this may potentially **ruin uplifts**.



(uplift Kachru, Kallosh, Linde, Trivedi '03)

- ▶ Due to the non-linear terms, the goldstini may produce such condensates.
- ▶ We will discuss **the simplest model** where goldstino condensation can be studied convincingly.

*Goldstino condensation at large  $N$*

FF, Matteo Moritsu '22

- ▶ We can work with N copies of 4D Dirac goldstini

$$\mathcal{L} = -Nf^2 \det[\mathcal{A}_m^a] = -Nf^2 + i\bar{G}^A \not{\partial} G^A + \dots$$

with  $A = 1, \dots, N$ , with cut-off  $\Lambda \ll \sqrt{f}$ , where

$$\mathcal{A}_m^a = \delta_m^a + \frac{i}{2Nf^2} \left( \bar{G}^A \gamma^a \partial_m G^A - \partial_m \bar{G}^A \gamma^a G^A \right),$$

and  $\delta G^A = -f\xi^A + \dots$

- ▶ The theory can be written as

$$\mathcal{L} = -Nf^2 \det[e_m^a] + Nf^2 C_a^m (e_m^a - \mathcal{A}_m^a),$$

making the path integral **Gaussian in the fermions**.



- ▶ We can perform the Gaussian integral over the fermions, to find formally

$$Z_F \sim (\det[iC_a^m \gamma^a \partial_m])^N,$$

and then the action for the bosons reads

$$S = N \times \left\{ -f^2 \int d^4x \left[ \det[e_n^b] - C_a^m (e_m^a - \delta_m^a) \right] - i \operatorname{tr} \log [iC_a^m \gamma^a \partial_m] \right\}.$$

- ▶ Restore momentarily  $\hbar$  and notice an “effective”  $\hbar$  as

$$\hbar_{\text{eff}} = \hbar/N,$$

infinitesimal **at large N** and **theory behaves “classically”**.

- ▶ We want to find **new stationary points** therefore we can focus on

$$C_a^m = (1 + h) \delta_a^m, \quad e_m^a = (1 - \phi) \delta_m^a.$$

- ▶ The potential reads

$$V(h, \phi) = N \left\{ f^2 \left[ (1 - \phi)^4 + 4(1 + h)\phi \right] - \frac{\Lambda^4}{16\pi^2} \log \left[ (1 + h)^2 \right] \right\}.$$

- ▶ We can readily eliminate  $h$  to get an effective  $V(\phi)$ ; an “effective potential” for the condensate to search for critical points.


- ▶ The potential has a new stationary point at

$$\phi_{GC} \simeq 1,$$

with

$$V(\phi_{GC}) \ll N f^2,$$

thus ruining uplifts.



- ▶ We interpret this as a goldstino condensate because classically

$$\phi = -\frac{i}{8Nf^2} \left( \bar{G}^A \gamma^m \partial_m G^A - \partial_m \bar{G}^A \gamma^m G^A \right).$$

## *Outlook*

- Large N: GC quite robust under higher derivative goldstino self-couplings / typical matter coupling of NL-SUSY.
- Using “exact RG flow”, we are studying the 4D goldstino condensate also at  $N=1$ , which is physically more relevant.  
*Dall’Agata, Emelin, FF, Moritsu ’22, work in progress*
- Our results resonate with gravitino condensation. *E.g.*  
*Jasinski, Smith ’83, Alexandre, Houston, Mavromatos ’13-’15*
- String theory interpretation? Tachyon mass/Decay time?  
Other dimensions? *Emelin, FF, Moritsu, work in progress*

Thank you