



SUSY2023

Quantum information & CP measurement in Higgs to tau tau at future lepton colliders

[*Phys.Rev.D* 107 (2023) 9, 093002]

Mohammad Mahdi Altakach

In collaboration with: P. Lamba, F. Maltoni, K. Mawatari and K. Sakurai


18 Jul. 2023

Part I


Introduction

Spin

- In **classical mechanics**, the components of **angular momentum** (I_x, I_y, I_z) take continuous real numbers.
- A striking fact, found in the **Stern-Gerlach** experiment, is that the measurement outcome of **spin** component is either **+1** or **-1** (in the $\hbar/2$ unit).



Alice & Bob




- **Alice** and **Bob** receive particles α and β , respectively, and measure the **spin z-component** of their particles. Repeat the process many times.
- **Alice** and **Bob** will find their results are completely random (**+1** and **-1**, **50-50%**).
- Nevertheless, their result is **100%** anti-correlated due to the **angular momentum** conservation. If **Alice**'s result is **+1**, **Bob**'s result is always **-1** and vice versa.

Alice	+	+	-	+	-	-	+	+	+	-	+
Bob	-	-	+	-	+	+	-	-	-	+	-
$S_z^\alpha \cdot S_z^\beta$	-	-	-	-	-	-	-	-	-	-	-


$$\langle S_z^\alpha \cdot S_z^\beta \rangle = -1$$

Hidden variable theory



Alice	+	+	-	+	-	-	+	+	+	-	+
Bob	-	-	+	-	+	+	-	-	-	+	-
$S_z^\alpha \cdot S_z^\beta$	-	-	-	-	-	-	-	-	-	-	-


Hidden variable theory



$$\overbrace{\underbrace{\hspace{10em}}_{\lambda}}^{\substack{\{\lambda_{+-}\} \quad \{\lambda_{-+}\}}} \\ P(\lambda \in \{\lambda_{+-}\}) = P(\lambda \in \{\lambda_{-+}\}) = \frac{1}{2}$$

Alice	+	+	-	+	-	-	+	+	+	-	+
Bob	-	-	+	-	+	+	-	-	-	+	-
$S_z^\alpha \cdot S_z^\beta$	-	-	-	-	-	-	-	-	-	-	-

Hidden variable theory



$$\overbrace{\underbrace{\hspace{10em}}_{\lambda}}^{\lambda}$$

$$\underbrace{\hspace{10em}}_{\{\lambda_{+-}\}} \quad \underbrace{\hspace{10em}}_{\{\lambda_{-+}\}}$$

$$P(\lambda \in \{\lambda_{+-}\}) = P(\lambda \in \{\lambda_{-+}\}) = \frac{1}{2}$$

- Particles have definite properties regardless of the measurement.
- **Alice**'s measurement has no influence on **Bob**'s particle.

(**realism**)

(**locality**)

Quantum mechanics (QM)

- Although their outcomes are different in each decay, **QM** says the **state** of the particles are exactly the same for all decays:


$$|\Psi^{(0,0)}\rangle \doteq \frac{|\overset{\alpha}{+} \overset{\beta}{-}\rangle_z - | - + \rangle_z}{\sqrt{2}}$$

↑
up to a phase $e^{i\theta}$

- Before the measurements, particles have no definite **spin**. Outcomes are undetermined.

(no realism)

- At the moment when Alice makes her measurement, the state collapses into:



Bob's outcome is completely determined (before his measurement) and **100%** anti-correlated with **Alice's**.

(non-local)

Entanglement

- The origin of this bizarre feature is **entanglement**.

general

$$|\Psi\rangle \doteq c_{11}|++\rangle_z + c_{12}|+-\rangle_z + c_{21}| - + \rangle_z + c_{22}|--\rangle_z$$

separable

$$|\Psi_{\text{sep}}\rangle \doteq [c_1^\alpha|+\rangle_z + c_2^\alpha|-\rangle_z] \otimes [c_1^\beta|+\rangle_z + c_2^\beta|-\rangle_z]$$

entangled

$$|\Psi_{\text{ent}}\rangle \not\doteq [c_1^\alpha|+\rangle_z + c_2^\alpha|-\rangle_z] \otimes [c_1^\beta|+\rangle_z + c_2^\beta|-\rangle_z]$$

entangled


$$|\Psi^{(0,0)}\rangle \doteq \frac{|+-\rangle_z - |-+\rangle_z}{\sqrt{2}}$$

Bob's measurement collapses the **state** of β to $|+\rangle_z$ or $|-\rangle_z$ but does not influence the **state** of α .


Bell inequalities

- It seems difficult to experimentally discriminate **QM** and **general hidden variable theories**.
- **John Bell (1964)** derived simple inequalities that can discriminate **QM** from any **local-real hidden variable theories**: **Bell inequalities**.

John Bell and his famous theorem in 1982 (Image: CERN)



Bell inequalities



- Finally we construct:

$$R_{\text{CHSH}} \equiv \frac{1}{2} \left| \langle s_a s_b \rangle - \langle s_a s_{b'} \rangle + \langle s_{a'} s_b \rangle + \langle s_{a'} s_{b'} \rangle \right|$$

One can show in **hidden variable theories** that: $R_{\text{CHSH}} \leq 1$
[Clauser, Horne, Shimony, Holt, 1969].

Bell inequalities

- In **QM**, for:

$$|\Psi^{(0,0)}\rangle \doteq \frac{|+-\rangle_z - |-+\rangle_z}{\sqrt{2}}$$

- One can show:


$$\langle s_a s_b \rangle = \langle \Psi^{(0,0)} | s_a s_b | \Psi^{(0,0)} \rangle = (\hat{\mathbf{a}} \cdot \hat{\mathbf{b}})$$

- Therefore:

$$\begin{aligned} R_{\text{CHSH}} &= \frac{1}{2} \left| \langle s_a s_b \rangle - \langle s_a s_{b'} \rangle + \langle s_{a'} s_b \rangle + \langle s_{a'} s_{b'} \rangle \right| \\ &= \frac{1}{2} \left| \underbrace{(\hat{\mathbf{a}} \cdot \hat{\mathbf{b}})}_{\frac{1}{\sqrt{2}}} - \underbrace{(\hat{\mathbf{a}} \cdot \hat{\mathbf{b}}')}_{-\frac{1}{\sqrt{2}}} + \underbrace{(\hat{\mathbf{a}}' \cdot \hat{\mathbf{b}})}_{\frac{1}{\sqrt{2}}} + \underbrace{(\hat{\mathbf{a}}' \cdot \hat{\mathbf{b}}')}_{\frac{1}{\sqrt{2}}} \right| \end{aligned}$$

violates the
upper
bound of
**hidden
variable
theories!**

$$= \sqrt{2}$$



Part II

Spin $1/2$ biparticle system

Density operator

Probability of having $|\psi_1\rangle$

- For a statistical ensemble $\{ \{p_1 : |\psi_1\rangle\}, \{p_2 : |\psi_2\rangle\}, \{p_3 : |\psi_3\rangle\}, \dots \}$, we define the **density operator/matrix**:

$$\hat{\rho} \equiv \sum_k p_k |\Psi_k\rangle \langle \Psi_k| \quad \rho_{ab} \equiv \langle e_a | \hat{\rho} | e_b \rangle$$

$$0 \leq p_k \leq 1$$

$$\sum_k p_k = 1$$

$$\langle e_a | e_b \rangle = \delta_{ab}$$

- **Density matrices** satisfy the conditions:

$$\hat{\rho}^\dagger = \hat{\rho}$$

$$\text{Tr } \hat{\rho} = 1$$

$$\hat{\rho} \text{ is positive definite, that is } \forall |\varphi\rangle; \langle \varphi | \hat{\rho} | \varphi \rangle \geq 0.$$

- The expectation of an observable \hat{O} is calculated by:

$$\langle \hat{O} \rangle = \text{Tr} [\hat{O} \hat{\rho}]$$

Spin 1/2 biparticle system

- The spin system of α and β particles has 4 independent bases:

$$(|e_1\rangle, |e_2\rangle, |e_3\rangle, |e_4\rangle) = (|++\rangle, |+-\rangle, |-+\rangle, |--\rangle)$$

3x3
matrix

- $\Rightarrow \rho_{ab}$ is a 4x4 matrix (hermitian, $\text{Tr}=1$). It can be expanded as

$$\rho = \frac{1}{4} (1 \otimes 1 + B_i \cdot \sigma_i \otimes 1 + \bar{B}_i \cdot 1 \otimes \sigma_i + C_{ij} \cdot \sigma_i \otimes \sigma_j), \quad B_i, \bar{B}_i, C_{ij} \in \mathbb{R}$$

- For the spin operators \hat{s}^α and \hat{s}^β :

$$\langle \hat{s}_i^\alpha \rangle = \text{Tr}[\hat{s}_i^\alpha \hat{\rho}] = B_i \quad \langle \hat{s}_i^\beta \rangle = \text{Tr}[\hat{s}_i^\beta \hat{\rho}] = \bar{B}_i \quad \langle \hat{s}_i^\alpha \hat{s}_j^\beta \rangle = \text{Tr}[\hat{s}_i^\alpha \hat{s}_j^\beta \hat{\rho}] = C_{ij}$$

spin-spin
correlation

Entanglement

- If the **state** is **separable** (not **entangled**):

$$\rho = \sum_k p_k \rho_k^\alpha \otimes \rho_k^\beta, \quad 0 \leq p_k \leq 1 \text{ and } \sum_k p_k = 1$$

- Then, a modified **matrix** by the partial transpose:

$$\rho^{T_\beta} = \sum_k p_k \rho_k^\alpha \otimes [\rho_k^\beta]^T$$

is also a physical **density matrix**, i.e. **Tr=1** and non-negative.

- For biparticle systems, **entanglement** $\iff \rho^{T_\beta}$ to be non-positive. [Peres-Horodecki (1996,1997)].
- A simple sufficient condition for **entanglement** is:

$$E \equiv C_{11} + C_{22} - C_{33} > 1 \quad [\text{Eur. Phys. J. C 82, 666 (2022)}]$$

$$H \rightarrow \tau^+ \tau^-$$

- Generic $H\tau\tau$ interaction:

$$\mathcal{L}_{int} = -\frac{m_\tau}{v_{SM}} \kappa H \bar{\Psi}_\tau (\cos \delta + i\gamma_5 \sin \delta) \Psi_\tau$$

$$\rho_{mn, \bar{m}\bar{n}} = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & e^{-i2\delta} & 0 \\ 0 & e^{-i2\delta} & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\text{SM: } (\kappa, \delta) = (1, 0)$$

$$B_i = \bar{B}_i = 0, \quad C_{ij} = \begin{pmatrix} \cos 2\delta & \sin 2\delta & 0 \\ -\sin 2\delta & \cos 2\delta & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad E = 2\cos 2\delta + 1$$

Estimation of C_{ij}

- Let's suppose a **spin 1/2** particle α is **at rest** and spinning in the \mathbf{s} direction.
- α decays into a measurable particle \mathbf{l}_α and the rest \mathbf{X} : $\alpha \rightarrow \mathbf{l}_\alpha + (\mathbf{X})$
- The decay distribution is generally given by : $\frac{d\Gamma}{d\Omega} \propto 1 + x_\alpha (\hat{I}_\alpha \cdot \mathbf{s})$

$x \in [-1, 1]$ is called spin-analysing power and depends on the decay
 $x = 1$ for $\tau^\pm \rightarrow \pi^\pm \nu$

Unit direction vector of \mathbf{l}_α
 measured at the rest frame of α

- One can show for $\alpha + \beta \rightarrow [\mathbf{l}_\alpha + (\mathbf{X})] + [\mathbf{l}_\beta + (\mathbf{X})]$:

$$\langle \hat{S}_i^\alpha \hat{S}_j^\beta \rangle = -9 \cdot \langle \hat{I}_i^\alpha \hat{I}_j^\beta \rangle$$

measurable at colliders, but

needs to reconstruct the α
 (β) rest frames


Part III

Higgs to tau tau @ lepton colliders


Why lepton colliders?

- Background $Z/\gamma \rightarrow \tau^+\tau^-$ is much smaller at **lepton** colliders
- We need to reconstruct each τ rest frame to measure \hat{I} . This is challenging at **hadron** colliders since partonic CoM energy is unknown for each event

LHC



ILC



Simulation

	ILC	FCC-ee
energy (GeV)	250	240
luminosity (ab^{-1})	3	5
beam resolution e^+ (%)	0.18	0.83×10^{-4}
beam resolution e^- (%)	0.27	0.83×10^{-4}
$\sigma(e^+e^- \rightarrow HZ)$ (fb)	240.1	240.3
# of signal ($\sigma \cdot \text{BR} \cdot L \cdot \epsilon$)	385	663
# of background ($\sigma \cdot \text{BR} \cdot L \cdot \epsilon$)	20	36

- Events were generated with **Madgraph5**
- We incorporate the detector effect by smearing energies of visible particles

$$E^{true} \rightarrow E^{obs} = (1 + \sigma_E \cdot \omega) \cdot E^{true} \quad \sigma_E = 0.03$$

Random number from a normal distribution

- We perform **100 pseudo-experiment** to estimate the statistical uncertainties

Solving kinematical constraints

- To determine the tau momenta, we have to reconstruct the unobserved neutrino momenta: $(p_x^\nu, p_y^\nu, p_z^\nu), (p_x^{\bar{\nu}}, p_y^{\bar{\nu}}, p_z^{\bar{\nu}})$.
- **6** unknowns can be constrained by **2** mass-shell conditions and **4** energy-momentum conservation:

$$m_\tau^2 = (p_{\tau^+})^2 = (p_{\pi^+} + p_{\bar{\nu}})$$



$$m_\tau^2 = (p_{\tau^-})^2 = (p_{\pi^-} + p_\nu)$$

$$(p_{ee} - p_Z)^\mu = p_H^\mu = [(p_{\pi^-} + p_\nu) + (p_{\pi^+} + p_{\bar{\nu}})]^\mu$$

- With the reconstructed momenta, we define $(\hat{r}, \hat{n}, \hat{k})$ basis at the Higgs rest frame.


$$C_{ij} = \langle \hat{s}_i^{(\tau^-)} \hat{s}_j^{(\tau^+)} \rangle = -9 \cdot \langle \hat{I}_i^- \hat{I}_j^+ \rangle$$

$(i, j = r, n, k)$



Impact parameter (IP)

- We use the information of the **impact parameter** \vec{b}_{\pm} measurement of π^{\pm} to “correct” the observed energies of τ^{\pm} and Z decay products.
- We check whether the reconstructed τ momenta are consistent with the measured impact parameters.
- We construct the likelihood function and search for the most likely τ momenta.



$$E_{\alpha}(\delta_{\alpha}) = (1 + \sigma_{\alpha}^E \cdot \delta_{\alpha}) \cdot E_{\alpha}^{\text{obs}}$$

$$\vec{b}_{+} = |\vec{b}_{+}| (\sin^{-1} \Theta_{+} \cdot \vec{e}_{\tau^{+}} - \tan^{-1} \Theta_{+} \cdot \vec{e}_{\pi^{+}})$$

$$\vec{\Delta}_{b_{+}}^i(\{\delta\}) \equiv \vec{b}_{+} - |\vec{b}_{+}| (\sin^{-1} \Theta_{+}^i(\{\delta\}) \cdot \vec{e}_{\tau^{+}}^i(\{\delta\}) - \tan^{-1} \Theta_{+}^i(\{\delta\}) \cdot \vec{e}_{\pi^{+}})$$

$$L_{\pm}^i(\{\delta\}) = \frac{[\Delta_{b_{\pm}}^i(\{\delta\})]_x^2 + [\Delta_{b_{\pm}}^i(\{\delta\})]_y^2}{\sigma_{b_T}^2} + \frac{[\Delta_{b_{\pm}}^i(\{\delta\})]_z^2}{\sigma_{b_z}^2}$$

$$L^i(\{\delta\}) = L_{+}^i(\{\delta\}) + L_{-}^i(\{\delta\})$$

Results

	ILC	FCC-ee
C_{ij}	$\begin{pmatrix} 0.830 \pm 0.176 & 0.020 \pm 0.146 & -0.019 \pm 0.159 \\ -0.034 \pm 0.160 & 0.981 \pm 0.1527 & -0.029 \pm 0.156 \\ -0.001 \pm 0.158 & -0.021 \pm 0.155 & -0.729 \pm 0.140 \end{pmatrix}$	$\begin{pmatrix} 0.925 \pm 0.109 & -0.011 \pm 0.110 & 0.038 \pm 0.095 \\ -0.009 \pm 0.110 & 0.929 \pm 0.113 & 0.001 \pm 0.115 \\ -0.026 \pm 0.122 & -0.019 \pm 0.110 & -0.879 \pm 0.098 \end{pmatrix}$
E_k	2.567 ± 0.279 > 5σ	2.696 ± 0.215 > 5σ
R_{CHSH}	1.103 ± 0.163	1.276 ± 0.094 $\sim 3\sigma$

SM values: $C_{ij} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix}$

$E = 3$ Entanglement $\implies E > 1$

$R_{\text{CHSH}} = \sqrt{2} \simeq 1.414$ Bell-nonlocal $\implies R_{\text{CHSH}} > 1$

The superiority of **FCC-ee** over **ILC** is due to a better **beam resolution**

	ILC	FCC-ee
energy (GeV)	250	240
luminosity (ab^{-1})	3	5
beam resolution e^+ (%)	0.18	$0.83 \cdot 10^{-4}$
beam resolution e^- (%)	0.27	$0.83 \cdot 10^{-4}$

CP measurement

- Under **CP**, the spin correlation matrix transforms: $C \xrightarrow{CP} C^T$
- This can be used for a **model-independent** test of **CP** violation. We define:

$$A \equiv (C_{rn} - C_{nr})^2 + (C_{nk} - C_{kn})^2 + (C_{kr} - C_{rk})^2 \geq 0$$

Observation of $A \neq 0$ immediately confirms **CP** violation

- From our simulation, we observe:

$$A = \begin{cases} 0.168 \pm 0.131 & (\text{ILC}) \\ 0.081 \pm 0.061 & (\text{FCC-ee}) \end{cases}$$

Consistent with the absence of **CPV**

CP measurement

- This model independent bounds can be translated to the constraint on the **CP**-phase δ :

$$\mathcal{L}_{\text{int}} \propto H \bar{\psi}_\tau (\cos \delta + i \gamma_5 \sin \delta) \psi_\tau \quad \rightarrow \quad C_{ij} = \begin{pmatrix} \cos 2\delta & \sin 2\delta & 0 \\ -\sin 2\delta & \cos 2\delta & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \rightarrow \quad A(\delta) = 4 \sin^2 2\delta$$

- Focusing on the region near $|\delta| = 0$, we find the 1- σ bounds:

$$\delta < \begin{cases} 7.9^\circ & (\text{ILC}) \\ 5.4^\circ & (\text{FCC-ee}) \end{cases}$$

Part IV

Summary

Summary

- High energy tests of **entanglement** and **Bell inequality** has recently attracted an attention.
- We investigated feasibility of **quantum** property tests @ **ILC** and **FCC-ee**.
- **Quantum tests** require a precise reconstruction of the τ rest frames and **IP** information is crucial to achieve this.
- Spin correlation is sensitive to **CP**-phase and we can measure the **CP**-phase as a byproduct of the quantum property measurement.

	Entanglement	Bell-inquality	CP-phase
FCC-ee	$> 5\sigma$	$\sim 3\sigma$	7.9°
ILC	$> 5\sigma$		5.4°