

From the exact solution for the Schrödinger equation to NLO cross sections

Yair Mulian
University of Santiago de Compostela



"Singularity is almost invariably a clue" (Sherlock Holmes)




How it all have started?

- **The initial project**: computing the virtual NLO cross section for dijet (inclusive / diffractive) production.
- **Previous knowledge**:

In the work (with E. Iancu) “***Dihadron production in DIS at NLO: the real corrections***” it has been observed that an elegant pattern appears for amplitudes:

$$\begin{aligned}
 |\gamma_T^i(Q, q^+, \mathbf{w})\rangle_{q\bar{q}g}^{\text{reg}} = & -\frac{ee_f g q^+}{2(2\pi)^4} \int_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \int_0^1 d\vartheta \int_0^{1-\vartheta} d\xi \frac{\vartheta}{\sqrt{\xi}} \frac{\mathbf{R}^j \mathbf{Y}^n}{\mathbf{Y}^2} \delta^{(2)}(\mathbf{w} - \mathbf{c}) \\
 & \times \left\{ \Phi_{\lambda_1 \lambda_2}^{ijmn}(\vartheta, \xi) \frac{Q K_1(QD)}{D} [U^{ab}(z)V(x)t^b V^\dagger(\mathbf{y}) - t^a]_{\alpha\beta} - (\mathbf{y}, \mathbf{z} \rightarrow \mathbf{y}') \right\} \\
 & \times \left| \bar{q}_{\lambda_2}^\beta((1-\vartheta-\xi)q^+, \mathbf{y}) g_m^a(\xi q^+, \mathbf{z}) q_{\lambda_1}^\alpha(\vartheta q^+, \mathbf{x}) \right\rangle,
 \end{aligned}$$




Opening Pandora's box

In practice, the calculation based on U leads to one contribution for which these properties **were missing** (!):


- 1) *The elegant pattern did not appear.*
- 2) *JIMWLK could not be observed at the amplitude level.*
- 3) *There were multiple poles for degenerate configurations.*

All that happened due to one (sub-)contribution:


Two WW fields (simple pole)



Branch-cut (multiple poles)



$(x, z \rightarrow w)$ Not working!



The Schrödinger Equation

The fundamental equation:

$$i \frac{d}{dt} |\Psi(t)\rangle = \hat{H}(t) |\Psi(t)\rangle, \quad \text{or} \quad \frac{d\hat{O}(t, t_0)}{dt} = -i\hat{H}(t)\hat{O}(t, t_0).$$

Predicts a dynamics which is **differentiable** (and therefore **continuous**) at each moment of the evolution.

- As long as the Hamiltonian involves no time dependence:

$$|\Psi(t)\rangle = \hat{U}(t, t_0) |\Psi(t_0)\rangle, \quad \hat{U}(t, t_0) = e^{-i\hat{H}(t-t_0)}.$$

- When time dependence is involved it is customary to use the solution

$$\hat{O}(t, t_0) = \hat{U}(t, t_0) \equiv \text{T exp} \left[-i \int_{t_0}^t dt' \hat{H}(t') \right]$$

Which can also be written in the expanded form as (Dyson series)

$$\hat{U}(t, t_0) = \hat{1} - i \int_{t_0}^t dt' \hat{H}(t') - \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') + \dots$$

How do we conclude that U is unitary?

- ***Exact unitarity is a crucial property for any valid quantum description.***

Using the Dyson expansion:

$$\hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) = \hat{\mathbf{1}} + \left(\int_{t_0}^t dt' \hat{H}(t') \right)^2 - \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') - \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t'') \hat{H}(t') + \dots$$

Changing variables:

$$\int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t'') \hat{H}(t') = \int_{t_0}^t dt' \int_{t'}^t dt'' \hat{H}(t') \hat{H}(t'')$$

After adding integrals:

$$\int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') + \int_{t_0}^t dt' \int_{t'}^t dt'' \hat{H}(t') \hat{H}(t'') = \left(\int_{t_0}^t dt' \hat{H}(t') \right)^2$$

So, at least allegedly, U is a unitary operator,

$$\hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) = \hat{\mathbf{1}}$$




Figure 4.1. Geometric interpretation of Eq. (4.21).

From Peskin and Schroeder, P. 85.

The underlying assumptions

1) Linearity:

$$\int_{t_0}^t dt' f(t') + \int_{t_0}^t dt' g(t') = \int_{t_0}^t dt' (f(t') + g(t'))$$

2) Changing order of integrations (Fubini theorem):

$$\int_{t_0}^t dt' \int_{t_0}^{t'} dt'' f(t', t'') = \int_{t_0}^t dt' \int_{t'}^t dt'' f(t', t'')$$

3) Adding integration intervals (additivity):

$$\int_{t_0}^{t'} dt'' f(t'') + \int_{t'}^t dt'' f(t'') = \int_{t_0}^t dt'' f(t'')$$

4) Exact Hermiticity of the Hamiltonian:

$$H^\dagger(t) = H(t)$$

- Properties 1-4 are **valid** for Hamiltonians expressed by **functions**.
- Properties 2-4 are **not valid** for **distributions**.

Linearity

Function: an object which assign exactly one **defined** (finite) element from a set Y to each element of set X .

- **Example:** $f(x) = \frac{1}{x-a}$ is a function defined on the union of two intervals $(-\infty, a) \cup (a, \infty)$, but is **NOT** a function on an interval that contains $x = a$.
- **What happens to linearity for intervals which cross the singularity point?**

```
FullSimplify[Integrate[x/(x-a), {x, -1, 1}] + Integrate[-a/(x-a), {x, -1, 1}]]
```

```
2 + a (-Log[-1-a] + Log[1-a] - Log[-1+a] + Log[1+a]) if Re[a] > 1 || Re[a] < -1 || a ∉ ℝ = 2 + a log(-1)
```

```
FullSimplify[Integrate[1, {x, -1, 1}]]
```

2

By using complex deformation (rotating the pole aside):

$$f(x) = \frac{1}{x-a} \quad \longrightarrow \quad O(x) = \frac{1}{x - ae^{i\epsilon}}$$

For ϵ finite, $O(x)$ is just a general complex function. For ϵ taken to 0, $O(x)$ is a sum of a function and distribution (*Sokhotski-Plemelj theorem*):

$$\lim_{\epsilon \rightarrow 0} O(x) = \lim_{\epsilon \rightarrow 0} \frac{1}{x - a - i\epsilon} = P.v. \left(\frac{1}{x - a} \right) + i\pi\delta(x - a)$$

Preliminaries in Analysis

Absolute convergence:

Both $\sum_n a_n$ and $\sum_n |a_n|$ converges, alternatively, both $\int f(x)$ and $\int |f(x)|$.

Conditional convergence:

$\sum_n a_n$ converges but not $\sum_n |a_n|$, alternatively $\int f(x)$ converges but not $\int |f(x)|$.

- **Riemann series theorem:** “if an infinite series of real numbers is conditionally convergent, then its terms can be arranged in a permutation so that the new series converges to an arbitrary real number, or diverges.”
- **Conclusion:** conditionally convergent contributions are very fragile.



Absolute conv.



Conditionally conv.

The Fubini Theorem

Exchanging the ordering of integrations is allowed only for absolute convergent integrals.

$$\int dx \left(\int dy f(x, y) \right) = \int dy \left(\int dx f(x, y) \right) \quad \text{if} \quad \int dx \int dy |f(x, y)| < \infty$$

- Example:

$$\int_0^2 \left(\int_0^1 \frac{xy(x^2 - y^2)}{(x^2 + y^2)^3} dy \right) dx = -\frac{1}{20} \quad \int_0^1 \left(\int_0^2 \frac{xy(x^2 - y^2)}{(x^2 + y^2)^3} dx \right) dy = \frac{1}{5}$$



Dr Peyam ✓
@drpeyam
150K subscribers

Check: "**Fubini Counterexample (full version)**" @ *Dr Payam* (youtube).

- Distributions typically **do not allow** to exchange the ordering of integrations.

For Y.M. theories, the terms of the perturbative expansion are usually divergent before regularization and **conditionally convergent** afterwards (but not absolutely convergent).

- **The simplifications for establishing unitarity are not always valid!**

Additivity

Additivity **does not hold** for distributions:

$$\int_{t_0}^{t'} dt'' f(t'')\delta(t'' - t_1) + \int_{t'}^t dt'' f(t'')\delta(t'' - t_1) \neq \int_{t_0}^t dt'' f(t'')\delta(t'' - t_1)$$

Since the if t_1 coincide with t' the integral may regarded as '*badly defined*', or the distribution considered to *contribute twice*.

```
FullSimplify[Integrate[f (t') * DiracDelta[t' - a], {t', 0, t1}] + Integrate[f (t') * DiracDelta[t' - a], {t', t1, t}], Assumptions -> t > 0]
```

```
a f ((-1 + 2 HeavisideTheta[t1]) HeavisideTheta[a - t1 HeavisideTheta[-t1]] HeavisideTheta[-a + t1 HeavisideTheta[t1]] + (-1 + 2 HeavisideTheta[t - t1]) HeavisideTheta[a - t1 HeavisideTheta[t - t1] - t HeavisideTheta[-t + t1]] HeavisideTheta[-a + t HeavisideTheta[t - t1] + t1 HeavisideTheta[-t + t1]]) if a ∈ ℝ && t1 ∈ ℝ
```

```
FullSimplify[Integrate[f (t') * DiracDelta[t' - a], {t', 0, t}], Assumptions -> t > 0]
```

```
a f HeavisideTheta[a, -a + t] if a ∈ ℝ
```

- Check: "*When functions have no value(s)*" by Steven G. Johnson.

Handling distributions with Dyson series

A hint on the incapability of the Dyson series to cope with distributions is evident since it involves a measure-0 direct product of two Hamiltonians (contact term).

$$\hat{U}(t, t_0) = \hat{\mathbf{1}} - i \int_{t_0}^t dt' \hat{H}(t') - \int_{t_0}^t dt' \hat{H}(t') \int_{t_0}^{t'} dt'' \hat{H}(t'') + \dots$$

The square of a distribution cannot be defined consistently:


"L. Schwartz, Sur l'impossibilité de la multiplication des distributions".

If our Hamiltonian is expressed via a distribution,

$$H(t) = h \delta(t - t_a)$$

As $\int_0^x dx' \delta(x' - a) = \Theta(x - a)$, the Dyson series becomes non-sense,

$$\int_0^x dx' \delta(x' - a) \Theta(x' - a) \longrightarrow \text{undefined.}$$



On the “Pitaron”

A unitarized version of U can be found by multiplying U by an additional operator:

$$\hat{\mathcal{P}}(t, t_0) \equiv \hat{\mathcal{N}}(t, t_0) \hat{U}(t, t_0)$$

With the Hermitian normalization operator:

$$\hat{\mathcal{N}}(t, t_0) \equiv \sqrt{\hat{U}^{\dagger-1}(t, t_0) \hat{U}^{-1}(t, t_0)}$$

Note that unitarity is **manifest** (unbreakable):

$$\hat{\mathcal{P}}^{\dagger}(t, t_0) \hat{\mathcal{P}}(t, t_0) = \hat{U}^{\dagger}(t, t_0) \hat{U}^{\dagger-1}(t, t_0) \hat{U}^{-1}(t, t_0) \hat{U}(t, t_0) = \hat{\mathbf{1}}$$

Regardless of the choice of Hamiltonian.

Wider class of exotic quantum field theories, in which unitarity is proven to be broken, can now be studied normally, such as QFT with **open systems**, **fractional dimensions** or **non-commutative spaces**.



How to take a square root?

The square root of a matrix extends the notion of square root from numbers to matrices. A matrix B is said to be a square root of A if the matrix product BB is equal to A .

- Hermitian operators can always be diagonalized with real entries on the diagonal, and therefore always have a square root.

$$\mathcal{N}_q^2 = a^2 |q\rangle \langle q| + b^2 |qg\rangle \langle qg| \longrightarrow \mathcal{N}_q = \pm a |q\rangle \langle q| \pm b |qg\rangle \langle qg|.$$

- A unique result is obtained by using the initial conditions.

The Schrodinger-Liouville's unification

Let us see that \hat{P} solves the Schrodinger equation:

$$\frac{d\hat{\mathcal{P}}(t, t_0)}{dt} = -i\hat{H}(t)\hat{\mathcal{P}}(t, t_0)$$


According to Leibniz product rule:

$$\frac{d\hat{\mathcal{P}}(t, t_0)}{dt} = \hat{\mathcal{N}}(t, t_0)\frac{d\hat{U}(t, t_0)}{dt} + \frac{d\hat{\mathcal{N}}(t, t_0)}{dt}\hat{U}(t, t_0) \quad \left[\hat{H}(t), \hat{\mathcal{N}}(t, t_0) \right] + \hat{\mathcal{N}}(t, t_0)\hat{H}(t) = \hat{H}(t)\hat{\mathcal{N}}(t, t_0)$$


We arrive at:

$$\hat{\mathcal{N}}(t, t_0) \left\{ \frac{d\hat{U}(t, t_0)}{dt} + i\hat{H}(t)\hat{U}(t, t_0) \right\} = \left\{ \frac{d\hat{\mathcal{N}}(t, t_0)}{dt} + i \left[\hat{H}(t), \hat{\mathcal{N}}(t, t_0) \right] \right\} \hat{U}(t, t_0)$$

Implying “two equations at once”:



$$\frac{d\hat{U}(t, t_0)}{dt} = -i\hat{H}(t)\hat{U}(t, t_0)$$



$$\frac{d\hat{\mathcal{N}}(t, t_0)}{dt} = -i \left[\hat{H}(t), \hat{\mathcal{N}}(t, t_0) \right]$$



Generalizing the Magnus expansion

Since P is an exact unitary operator it can be written **without** the time-ordering “T operator”:

$$\hat{P}(t, t_0) = e^{-i\hat{\Omega}(t, t_0)} \quad \Omega(t, t_0) = \int_{t_0}^t dt' \hat{H}(t') - \frac{i}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' [\hat{H}(t'), \hat{H}(t'')]]$$

Computing the inverse:

$$\hat{U}^{-1}(t, t_0) = \hat{\mathbf{1}} + i \int_{t_0}^t dt' \hat{H}(t') - \left(\int_{t_0}^t dt' \hat{H}(t') \right)^2 + \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') + \dots$$

Then:

$$\hat{N}(t, t_0) = \hat{\mathbf{1}} - \frac{1}{2} \left(\int_{t_0}^t dt' \hat{H}(t') \right)^2 + \frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \{ \hat{H}(t'), \hat{H}(t'') \} + \dots$$

One arrives at:

$$\hat{P}(t, t_0) = \hat{\mathbf{1}} - i \int_{t_0}^t dt' \hat{H}(t') - \frac{1}{2} \left(\int_{t_0}^t dt' \hat{H}(t') \right)^2 - \frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' [\hat{H}(t'), \hat{H}(t'')] + \dots$$



“*On the Exponential Solution of Differential Equations for a Linear Operator*” by Wilhelm Magnus (1954). **With this expansion the anticipated JIMWLK structure is restored!**

N via Truncation

There are several different ways in which the operator N can get a non-trivial value.

A trivial demonstration is via truncation of the perturbative expansion at first order, assuming **nilpotent Hamiltonian** or **negligible correction at order g^2** :

$$\left\| \int_{t_0}^t dt' \hat{H}(t') \right\| \gg \left\| \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') \right\|$$

In these cases only one term should be kept:

$$\hat{U}(t, t_0) \approx \hat{\mathbf{1}} - i \int_{t_0}^t dt' \hat{H}(t')$$

The operator N can be computed via

$$\hat{N}(t, t_0) = \left(\sqrt{\hat{U}(t, t_0) \hat{U}^\dagger(t, t_0)} \right)^{-1}$$

Assuming Hermiticity, leads to:

$$\hat{N}(t, t_0) \approx \hat{\mathbf{1}} - \frac{1}{2} \left\| \int_{t_0}^t dt' \hat{H}(t') \right\|^2$$

The world of Hamiltonians

- **Bounded**

Hermitian Hamiltonian with ‘*functional entries*’ and therefore contain no singularities. No poles or distributions are involved.

$$\hat{U}^\dagger(t, t_0) \frac{d\hat{U}(t, t_0)}{dt} = -i\hat{U}^\dagger(t, t_0) \hat{H}(t) \hat{U}(t, t_0), \quad \frac{d\hat{U}^\dagger(t, t_0)}{dt} \hat{U}(t, t_0) = i\hat{U}^\dagger(t, t_0) \hat{H}(t) \hat{U}(t, t_0)$$

$$\frac{d}{dt} \left(\hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) \right) = 0 \quad \longrightarrow \quad \hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) = \text{const}$$

- **Unbounded**

Hamiltonian that contain at least one point of singularity as distributions or $\hat{H}(t) = \frac{t - t_1}{(t - t_1)^2 - \Delta^2} \hat{\mathbf{1}}$
Singular Hamiltonians break the unitarity of U (but not of P) for a certain intervals.

$$\hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) \Big|_{-\infty < t_0 \leq t < t_1 + |\Delta| - \epsilon} = \hat{\mathbf{1}}, \quad \hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) \Big|_{t_1 + |\Delta| - \epsilon < t_0 \leq t < t_1 + |\Delta| + \epsilon} \neq \hat{\mathbf{1}},$$
$$\hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) \Big|_{t_1 + |\Delta| + \epsilon < t_0 \leq t < \infty} = \hat{\mathbf{1}},$$

- **Non-Hermitian Hamiltonians**


Generalization which allows us to study the singular case by taking a limit (**carefully**).

Breaking unitarity in QM

The Dirac comb potential is defined by:

$$V(t) = \sum_{i=1}^n V_i \delta(t - t_i)$$

with $\hat{H}(t, \hat{x}, \hat{p}) = \frac{\hat{p}^2}{2m} + \hat{V}(t, \hat{x}, \hat{p})$.




`H[t_] = 0.4*DiracDelta[t - 1] + 0.6*DiracDelta[t - 2] + 0.8*DiracDelta[t - 3] + 0.5*DiracDelta[t - 4]`


`0.5 DiracDelta[-4 + t] + 0.8 DiracDelta[-3 + t] + 0.6 DiracDelta[-2 + t] + 0.4 DiracDelta[-1 + t]`

`f[t_] := 1 + Integrate[H[t1]*H[t2], {t1, 0, t}, {t2, 0, t}] - 2 Integrate[H[t1]*H[t2], {t1, 0, t}, {t2, 0, t1}]`

`Plot[f[t], {t, 0, 5}]`



$$\begin{aligned} \hat{U}^\dagger(t, t_0) \hat{U}(t, t_0) &= \hat{1} + \left(\int_{t_0}^t dt' \hat{H}(t') \right)^2 \\ &\quad - \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') - \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t'') \hat{H}(t') + \dots \end{aligned}$$



Time independent potential

The familiar argument why time-independent Hamiltonians preserve unitarity:

$$\left(\exp\left[-i\hat{G}\right]\right)^\dagger = \left(\hat{\mathbf{1}} + \sum_{n=1}^{\infty} \frac{(-i\hat{G})^n}{n!}\right)^\dagger = \hat{\mathbf{1}} + \sum_{n=1}^{\infty} \frac{(i\hat{G})^n}{n!} = \left(\exp\left[i\hat{G}\right]\right)^{-1}$$

With $\hat{G} \equiv (t - t_0)\hat{H}$.

One of the canonical examples in QM:

$$\hat{V}(\hat{x}) = \alpha\delta(\hat{x})$$

However, the Taylor expansion can no longer be made and:

$$\left(\exp\left[-i\int_{t_0}^t dt' \delta(\hat{x})\right]\right)^\dagger \neq \left(\exp\left[-i\int_{t_0}^t dt' \delta(\hat{x})\right]\right)^{-1}$$

How to perform complex deformations?


The pole becomes a winding/portal point via the complexification

$$\hat{H}(t) \in \mathbb{R} \longrightarrow \hat{H}(z) \in \mathbb{C}$$

If the complex limit exist, one can interpret it as the original Hamiltonian:

$$\lim_{\text{Im } z \rightarrow 0} \hat{H}(z) = \hat{H}(t).$$

- *In order for a complex limit to exist, each way in which z can approach z_0 must yield the same limiting value.*



$$\hat{N}(t, t_0) = \hat{\mathbf{1}} - \frac{1}{2} \left(\int_{t_0}^t dt' \hat{H}(t') \right)^\dagger \int_{t_0}^t dt' \hat{H}(t') + \frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') + \frac{1}{2} \left(\int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') \right)^\dagger + \dots$$

$$\begin{aligned} \hat{P}(t, t_0) &= \hat{\mathbf{1}} - i \int_{t_0}^t dt' \hat{H}(t') - \frac{1}{2} \left(\int_{t_0}^t dt' \hat{H}(t') \right)^\dagger \left(\int_{t_0}^t dt' \hat{H}(t') \right) \\ &\quad - \frac{1}{2} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') + \frac{1}{2} \left(\int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \hat{H}(t') \hat{H}(t'') \right)^\dagger + \dots \end{aligned}$$

An example in QM

Let us look on the case of a free particle subject for two kinds of perturbations:

$$\begin{aligned} H_0(t) &= \frac{p^2}{2m}, & H(t) &= \frac{g}{\Delta} (\sin(\omega(t-t_1))\hat{\sigma}_2 + \hat{\sigma}_3), \\ H_0(t) &= \frac{p^2}{2m}, & H(t) &= g \frac{t-t_1}{(t-t_1)^2 - \Delta^2} \hat{\mathbf{1}}. \end{aligned}$$

One note that the first perturbation keep unitarity exact:

$$\begin{aligned} \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \sin(\omega(t'-t_1)) \sin(\omega(t''-t_1)) &= \int_{t_0}^t dt' \int_{t'}^t dt'' \sin(\omega(t'-t_1)) \sin(\omega(t''-t_1)) \\ &= \frac{1}{2\omega^2} (\cos(\omega(t_1-t_0)) - \cos(\omega(t-t_1)))^2. \end{aligned}$$

This is no longer valid for the singular Hamiltonian. In order to study a singular Hamiltonian we study a larger class of non-hermitian Hamiltonians:

$$\hat{H}(t) = g \frac{t-t_1}{(t-t_1)^2 - \Delta^2} \hat{\mathbf{1}} \quad \longrightarrow \quad \hat{H}(t, i\epsilon) = g \frac{t-t_1}{(t-t_1)^2 - \Delta^2 - i\epsilon} \hat{\mathbf{1}}$$

The original dynamics is obtained as a private case of a vanishing deformation:

$$\hat{H}(t) = \lim_{\epsilon \rightarrow 0} \hat{H}(t, i\epsilon)$$

$$\hat{\mathcal{N}}(t, t_0) = \hat{\mathbf{1}} - \frac{1}{2}g^2\pi^2 \left(\int_{t_0}^t dt' (t' - t_1) \delta((t' - t_1)^2 - \Delta^2) \right)^2$$

$$- g^2\pi^2 \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' (t' - t_1)(t'' - t_1) \delta((t' - t_1)^2 - \Delta^2) \delta((t'' - t_1)^2 - \Delta^2).$$

$$\hat{\mathcal{N}}(t, t_0) = \hat{\mathbf{1}} - \frac{1}{8}g^2\pi^2 \text{Boole} [t_0 < -\Delta + t_1 < t \ || \ t < -\Delta + t_1 < t_0]$$




Figure 4. Plot from mathematica describing the time evolution of $\hat{\mathcal{N}}$ from eq. (4.18) for $t \in [0, 4]$. The following parametrization was used $g = 0.4$, $\Delta = 0.5$, $t_0 = 0$, $t_1 = 1$.

$$H(t) = \frac{t - t_1}{(t - t_1)^2 - \Delta^2 - i\epsilon} + \frac{t - t_2}{(t - t_2)^2 - \Delta^2 - i\epsilon}$$

$$\begin{aligned} \hat{\mathcal{N}}(t, t_0) = & \hat{\mathbf{1}} - \frac{1}{2}g^2\pi^2 \left(\int_{t_0}^t dt' \left((t' - t_1)\delta((t' - t_1)^2 - \Delta^2) + (t' - t_2)\delta((t' - t_2)^2 - \Delta^2) \right) \right)^2 \\ & - g^2\pi^2 \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' \left((t' - t_1)\delta((t' - t_1)^2 - \Delta^2) + (t' - t_2)\delta((t' - t_2)^2 - \Delta^2) \right) \\ & \times \left((t'' - t_1)\delta((t'' - t_1)^2 - \Delta^2) + (t'' - t_2)\delta((t'' - t_2)^2 - \Delta^2) \right). \end{aligned}$$




Figure 5. Plot from mathematica describing the time evolution of $\hat{\mathcal{N}}$ from eq. (4.20) for $t \in [0, 4]$. The following parametrization was used $g = 0.6$, $\Delta = 0.8$, $t_0 = 0$, $t_1 = 0.5$, $t_2 = 2$.

Can we really iterate limits?

Iteration of limits is not guaranteed usually, as in the simple example below:

$$\lim_{y \rightarrow 0} \lim_{x \rightarrow 0} \frac{x^2}{x^2 + y^2} = \lim_{y \rightarrow 0} 0 = 0, \quad \lim_{x \rightarrow 0} \lim_{y \rightarrow 0} \frac{x^2}{x^2 + y^2} = \lim_{x \rightarrow 0} 1 = 1.$$

$$\lim_{n \rightarrow \infty} \int dx \frac{1}{\left(1 + \frac{x}{n}\right)^n - z} \neq \int dx \lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{x}{n}\right)^n - z} = \int dx \frac{1}{e^x - z}$$

the necessary condition to permit that is **Lebesgue dominated convergence**.

- When using complex deformations, one must be careful to perform the limits in the correct ordering:

$$\int d\Phi_{f,j,i} \lim_{\epsilon \rightarrow 0} \frac{1}{E_f - E_i - 2i\epsilon} \lim_{\epsilon \rightarrow 0} \frac{1}{E_j - E_i - i\epsilon} \neq \lim_{\epsilon \rightarrow 0} \int d\Phi_{f,j,i} \frac{1}{E_f - E_i - 2i\epsilon} \frac{1}{E_j - E_i - i\epsilon}$$

$$\int \frac{d^d \mathbf{p}}{(2\pi)^d} \lim_{\epsilon \rightarrow 0} f(\mathbf{p}, \epsilon) \neq \lim_{\epsilon \rightarrow 0} \int \frac{d^d \mathbf{p}}{(2\pi)^d} f(\mathbf{p}, \epsilon)$$

QFT = Functions + Distributions

For the asymptotic states, after deformation the order g term of U reads:

$$i \int_{-\infty}^{0-} dt' H(t', \mathbf{p}) = i \int_{-\infty}^{0-} dt' \int d\Phi_{f,i} \lim_{\epsilon \rightarrow 0} e^{i(E_f - E_i - i\epsilon)t'} |f\rangle \langle f| \hat{H}(\mathbf{p}) |i\rangle \langle i|$$

← **After plugging**
 $H(t, \mathbf{p}) = e^{i\hat{H}(\mathbf{p})t} \hat{H}(\mathbf{p}) e^{-i\hat{H}(\mathbf{p})t}$

- *The time integration commutes with the other operations:*

$$\left[\int_{-\infty}^{0-} dt', \lim_{\epsilon \rightarrow 0} \right] = \left[\int_{-\infty}^{0-} dt', \int d\Phi_{f,i} \right] = 0, \quad \left[\int d\Phi_{f,i}, \lim_{\epsilon \rightarrow 0} \right] \neq 0$$

Then:

$$i \int_{-\infty}^{0-} dt' H(t', \mathbf{p}) = i \int d\Phi_{f,i} \lim_{\epsilon \rightarrow 0} \int_{-\infty}^{0-} dt' e^{i(E_f - E_i - i\epsilon)t'} |f\rangle \langle f| \hat{H}(\mathbf{p}) |i\rangle \langle i|$$

Which leads to:

$$i \int_{-\infty}^{0-} dt' H(t', \mathbf{p}) = \int d\Phi_{f,i} \lim_{\epsilon \rightarrow 0} \frac{1}{(E_f - E_i - i\epsilon)} |f\rangle \langle f| \hat{H}(\mathbf{p}) |i\rangle \langle i|$$

By using Sokhotski-Plemelj theorem:

$$\int_{-\infty}^{0-} dt' H(t', \mathbf{p}) = \int d\Phi_{f,i} \left[P.v. \left(\frac{1}{E_f - E_i} \right) + i\pi\delta(E_f - E_i) \right] |f\rangle \langle f| \hat{H}(\mathbf{p}) |i\rangle \langle i|$$

Usually not important due


$$x \delta(x) = 0$$

Example in QED

The time ordered exponential for electron:

$$\hat{U}(0, -\infty) |e^-\rangle = \left(\hat{1} - i \int |e^-\gamma\rangle \frac{\langle e^-\gamma | \hat{H} | e^-\rangle}{E_{e^-\gamma} - E_{e^-} - i\epsilon} \langle e^- | - \int |\tilde{e}^-\rangle \frac{\langle \tilde{e}^- | \hat{H} | e^-\gamma\rangle \langle e^-\gamma | \hat{H} | e^-\rangle}{(E_{\tilde{e}^-} - E_{e^-} - i\epsilon)(E_{e^-\gamma} - E_{e^-} - i\epsilon)} \langle e^- | \right) |e^-\rangle$$

The electron self-energy has originally a ‘*badly defined*’ energy denominator. For no valid justification, removed due combining with its (c.c.) amplitude.



Afterwards, one should introduce the “**WF normalization**” or “**LSZ factor**”,

$$\hat{U}(0, -\infty) |e^-\rangle|_{corr.} \longrightarrow Z \hat{U}(0, -\infty) |e^-\rangle|_{corr.}$$

And extract **Z** by demanding unitarity,

$${}_{corr.} \langle e^- | |Z|^2 \hat{U}^\dagger(0, -\infty) \hat{U}(0, -\infty) |e^-\rangle|_{corr.} = 1$$

Reproducing via N

Keeping only the relevant terms, the following expansion for N is obtained:

$$\hat{\mathcal{N}}(0, -\infty) = \hat{\mathbf{1}} + \int |\tilde{e}^-\rangle \frac{\langle \tilde{e}^- | \hat{H} | e^-\gamma \rangle \langle e^-\gamma | \hat{H} | e^-\rangle}{(E_{\tilde{e}^-} - E_{e^-} - i\epsilon)(E_{e^-\gamma} - E_{e^-} - i\epsilon)} \langle e^- | \leftarrow \text{Removing the ill defined term of } U$$
$$- \frac{1}{2} \int |\tilde{e}^-\rangle \frac{\langle \tilde{e}^- | \hat{H} | e^-\gamma \rangle \langle e^-\gamma | \hat{H} | e^-\rangle}{(E_{e^-\gamma} - E_{\tilde{e}^-} + i\epsilon)(E_{e^-\gamma} - E_{e^-} - i\epsilon)} \langle e^- | \leftarrow \text{Introducing the term that we practically use}$$

- *We arrive at the very same familiar expression that we practically use, but this time from first principles!*

The WF for quark anti-quark

Assuming (wrongly) that normalization is conducted via $|q\bar{q}\rangle \rightarrow \sqrt{Z_q}\sqrt{Z_{\bar{q}}}|q\bar{q}\rangle$, at g^2 , the general structure of the WF for two partons based on U contains “self energy” and “gluon exchange” contributions:

$$|q\bar{q}\rangle_{g^2} = g^2 \int ((s.e.) + (branch - cut)) |q\bar{q}\rangle$$

The “gluon exchange” contribution is associated with a “branch cut integral”:

$$I = \int \frac{d^d \mathbf{p}}{(2\pi)^d} \lim_{\epsilon \rightarrow 0} \frac{\mathbf{p}}{\beta \mathbf{p}^2 - i\epsilon} \cdot \lim_{\epsilon \rightarrow 0} \frac{\mathbf{p} - \alpha \mathbf{k}}{\beta \mathbf{p}^2 - \gamma (\mathbf{p} - \alpha \mathbf{k})^2 - 2i\epsilon}$$

- Based on P , the structure will consist of “self-energy”, “2 Weizsäcker-Williams fields” and *anti-Hermitian combination* (due to the commutator) of “branch-cut” contributions:

$$|q\bar{q}\rangle_{g^2} = g^2 \int \left((s.e.) + (2w.w.) + (branch - cut) - (branch - cut)^\dagger \right) |q\bar{q}\rangle$$


The “gluon exchange” contribution is associated with the 2 WW fields:

$$I = \int \frac{d^d \mathbf{p}}{(2\pi)^d} \lim_{\epsilon \rightarrow 0} \frac{\mathbf{p}}{\beta \mathbf{p}^2 - i\epsilon} \cdot \lim_{\epsilon \rightarrow 0} \frac{\mathbf{p} - \alpha \mathbf{k}}{\gamma (\mathbf{p} - \alpha \mathbf{k})^2 + i\epsilon}$$



NLO cross sections: U vs P

The crucial (unbridgeable) difference between U and P emerges when computing the gluon exchange with shockwave prior to the gluon emission.



- **Based on U**: leads to extremely complicated integrals which contain a measure-0 of unregularizable delta-functions. See appendix J in “*Dijet impact factor in DIS at next-to-leading order in the Color Glass Condensate*”, F. Salazar.
- **Based on P**: trivial calculation (subtraction term):

$$\begin{aligned} \frac{d\sigma_{V \text{ ex.}}}{dk_1^+ d^2k_1 dk_2^+ d^2k_2} &= -\frac{2\alpha_{em} \alpha_s N_c}{(2\pi)^6 q^+} \left(\sum e_f^2 \right) \delta(q^+ - k_1^+ - k_2^+) \\ &\times \int_{\bar{x}, \bar{y}, x', y', z} \int_0^{1-\vartheta} d\xi e^{-ik_1 \cdot (x' - \bar{x}) - ik_2 \cdot (y' - \bar{y})} ((2\vartheta - 1)(2\vartheta + 2\xi - 1) + 1) \frac{4\vartheta(1 - \vartheta - \xi) + 2\xi}{\xi\vartheta(1 - \vartheta)} \\ &\times \frac{R' \cdot \bar{R}}{|R'| |\bar{R}|} \frac{X \cdot Y}{X^2 Y^2} \tilde{Q} \tilde{Q} K_1(\tilde{Q}|R'|) K_1(\tilde{Q}|\bar{R}) \mathcal{W}(x', y', \bar{y}, \bar{x}). \end{aligned}$$

Analytic properties of perturbative calculations

- For U, non-intuitive properties appears:

- 1) *The cross section can turn very large and negative.*
- 2) *The result involve contributions which are not Fourier transformable.*
- 3) *The NLO is bigger than LO and so on.*
- 4) *JIMWLK cannot be shown at the amplitude level.*

- For P:

None of these problems.


Positive definite energy denominator with Intuitive properties.

- Occam's razor: *“Entities must not be multiplied beyond necessity”.*

Singular Functions

- 1) $f(x)$ is continuous (but not absolutely continuous), and non-constant on $[a, b]$.
- 2) There exists a set N of measure 0 such that for all x outside of N the derivative $f'(x)$ exists and is zero, that is, the derivative of $f(x)$ vanishes almost everywhere.

A standard example of a singular function is the *Cantor ternary function* (or alternatively as *Lebesgue's singular function* or *Devil's staircase*). In practical terms, a singular function can be expressed as a **continuous sum of delta functions**.



Summary

- 1) The unitarity of the Dyson series is preserved for functions, but is broken by distributions.
- 2) The discrete part of the dynamics is contained in the normalization operator N .
- 3) The solution P is manifestly unitary, even when the terms of the series are conditionally convergent, or distributions are used.
- 4) Wider class of quantum field theories, in which unitarity is known to be broken, can now studied normally.
- 5) *Happy end for the “saga”*: the elegant structure have appeared again.

