

Luminosity measurement at CMS.

(Вимірювання світимості для експерименту CMS.)

Olena Karacheban

11 November 2022, Geneva, CERN

Дякую!



Дорогі вчителі, дякую вам за ваш час, за вашу зацікавленість і мотивацію розвиватися!

За те що хочете навчати нове покоління і оновлюєте програму навчання!


Завдякі вам буде більше зацікавлених фізикою молодих людей!



Outline




- **Ланцюг прискорювачів CERN і LHC.**
- **Експеримент CMS.**
- **Структура пучків частинок на LHC.**
- **Вимірювання світимості.**
- **Онлайн-детектори світимості.**
- **Калібрування світимості:**
 - **поправки;**
 - **стабільність і спеціальне сканування світимості.**
- **3D інтерактивна модель CMS детектора.**
- **CERN accelerators chain and LHC.**
- **CMS experiment.**
- **Particle beams structure at LHC.**
- **Luminosity measurements.**
- **Online luminosity detectors.**
- **Luminosity calibration**
 - **Corrections;**
 - **Stability and special emittance scans.**
- **3D interactive model of CMS detector.**

LHC — proton-proton collider, $\sqrt{s}_{\text{max}} = 14 \text{ TeV}$



CERN is accelerator complex!


CERN's Accelerator Complex



<https://youtu.be/RDdPuL-uOQc>

видео!

More about CERN and LHC



видео!

<https://www.youtube.com/watch?v=FLrEghnKncA>




**National geographic
видео с русской озвучкой!**

<https://youtu.be/ETWPnOOE1LQ>

What is LHC Run 1 - Run 2 - Run 3 ... and High Lumi LHC (HL-LHC)?

- LHC construction: 1998 - 2008, first collisions took place in 2010.
- First era of data taking -- LCH Run 1 -- started in 2011.
- Second era -- LCH Run 2: 2015 - 2018.
- Today we are at the beginning of Run 3! ★



4th July 2012
Higgs discovery


LHC Run 2, 2015-2018



- Energy per beam:
6.5 - 7 TeV
- The beam:
~2800 proton bunches
~ 10^{11} protons per bunch
- **Bunch spacing 25/50 ns**
- Beam width: ~20 μm




Each 10th RF bucket filled \rightarrow 25 ns




More about LHC beam structure


One LHC orbit (89 μ sec length, 89000 ns / 2.5 ns = 35600 RF buckets):







One bunch train example






One bunch profile, shape in XY plane:




How do we see collisions?





CMS detector



CMS pixel and tracker detectors



CMS Tracker showing silicon strips detectors in the barrel module.




Silicon pixel detector has 3 layers: 4cm, 7cm and 11cm from beam center. Each pixel is $100\mu\text{m}$ by $150\mu\text{m}$, about two hairs widths. 10 million particles per square centimetre per second.

The tracker silicon strip detector consists of four inner barrel (TIB) layers, six concentric layers of the outer barrel (TOB). Finally two endcaps (TEC) close off the tracker.

Each module consists of three elements: a set of sensors, its mechanical support structure and readout electronics.

CMS calorimeters and muon system



CMS = “Compact Muon Solenoid”: muons is one of CMS’s most important tasks.




One of the clearest "signatures" of the Higgs Boson is its decay into four muons.

In total there are 1400 muon chambers: 250 drift tubes (DTs) and 540 cathode strip chambers (CSCs) track the particles' positions.


The Electromagnetic Calorimeter (ECAL) measures the energy of electrons and photons by stopping them completely.

Hadrons, which are composite particles made up of quarks and gluons, fly through the ECAL and are stopped by the outer layer called the Hadron Calorimeter (HCAL).

Can CMS record all collisions?



- These CMS event displays shows just some examples of recorded collisions.
- Not all collisions are recorder due to processing time and time needed to readout detectors
- **Trigger system** based on interesting signals in the detectors decides which events to record.




3D interactive CMS detector


iSpy WebGL 4lepton.ig:Events/Run_178424/Event_666626491 [1 of 3]

DT Rec. Hits

Physics

- Vertices (Reco)
- Tracker Muons (Reco)
- Stand-alone Muons (Reco)
- Global Muons (Reco)
- Jets (PF)
- Jets (Reco)
- Missing Et (PF)
- Missing Et (Reco)

 CMS Experiment at the LHC, CERN
Data recorded: 2011-Oct-13 12:47:38.421105 GMT
Run / Event / LS: 178424 / 666626491 / 585






<https://opendata.cern.ch/visualise/events/cms>
<https://www.i2u2.org/elab/cms/event-display/#>

Спробуйте самі!

Pileup

Number of primary vertices
in one pp collision.

Is pileup 35 and 70 so
different???




Luminosity

What is luminosity \mathcal{L} ?



→ Proportionality factor between cross section σ_p and number of interactions per second $\frac{dR}{dt}$

$$\frac{dR}{dt} = \mathcal{L} \times \sigma_p \quad (\rightarrow \text{units: cm}^{-2}\text{s}^{-1})$$

Lower luminosity collision




Higher luminosity collision




Luminosity formula

For one collision:



For LHC collisions:



$$\mathcal{L} = \frac{N_1 N_2}{4\pi\sigma_x\sigma_y}$$

$$\mathcal{L} = \frac{N_1 N_2 f n_b}{4\pi\sigma_x\sigma_y}$$

Beam size in X and Y (about $15\mu\text{m}$): $\sigma_{x,y}$
 Number of protons in Beam 1,2: $N_{1,2}$

LHC revolution frequency: $f=11\text{kHz}$
 Number of bunches in LHC (about 2800):
 n_b

Example

Rare interactions and cross section

- Cross section σ measures how often a process occurs
- Characteristic for a given process
- Measured in: *barn* = $b = 10^{-28} m^2$ ($pb = 10^{-40} m^2$)

More common: *barn* = $b = 10^{-24} cm^2$ ($pb = 10^{-36} cm^2$)

We have for the LHC energy:

$$\sigma(pp \rightarrow X) \approx 0.1 b \text{ and}$$

$$\sigma(pp \rightarrow X + H) \approx 1 \cdot 10^{-11} b$$

$$\sigma(pp \rightarrow X + H \rightarrow \gamma\gamma) \approx 50 \cdot 10^{-15} b = 50 fb \text{ (femtobarn)}$$

- VERY rare (one in $2 \cdot 10^{12}$), need many collisions ...

Calculations for the LHC


- $N_1 = N_2 = 1.15 \times 10^{11}$ particles/bunch
- $n_b = 2808$ bunches/beam
- $f = 11.2455$ kHz, $\phi = 285 \mu rad$
- $\beta_x^* = \beta_y^* = 0.55$ m
- $\sigma_x^* = \sigma_y^* = 16.6 \mu m$, $\sigma_s = 7.7$ cm

$$\mathcal{L} = 1.200 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$$

$$\frac{dR}{dt} = \mathcal{L} \times \sigma_p$$

For small cross certain processes we need high luminosity to see some events!

Why do we want to collide at higher energy?



What really counts: Integrated luminosity

■ $\mathcal{L}_{\text{int}} = \int_0^T \mathcal{L}(t) dt$

■ $\mathcal{L}_{\text{int}} \cdot \sigma_p = \int_0^T \mathcal{L}(t) dt \cdot \sigma_p = \text{total number of events observed of process } p$

■ Unit is: cm^{-2} , i.e. inverse cross-section

■ Often expressed in **inverse barn**

■ **1 fb^{-1} (inverse femto-barn)** is $10^{39} cm^{-2}$

■ for 1 fb^{-1} : requires $10^6 s$ at $L = 10^{33} cm^{-2} s^{-1}$

■ Assume:



➤ You have accumulated 20 fb^{-1} (inverse femto-barn)

➤ You are interested in $\sigma(pp \rightarrow X + H \rightarrow \gamma\gamma) \approx 50 \text{ fb}$ (femtobarn)


➤ You have $20 \text{ fb}^{-1} \cdot 50 \text{ fb} = 1000$

➤ You have 1000 events of interest in your data !!



Full data 2015 - 2018:



Run 2 and Run 3 timelines



Run 2 luminosity



(see more [here](#))

LHC plans for Run 3

Run 3 will start in last week of **March, 2022**.

The energy will be 6.8 TeV per beam, so **13.6 TeV** centre-of-mass energy.

The current schedule foresees a 3 year run, but there will be very **likely an extra year** (2025).

Hope to get $\sim 30 \text{ fb}^{-1}$ in 2022, then 80 fb^{-1} per year. So 270 fb^{-1} at the end of Run 3. PU will be ~ 53 .
We will at least double and, with some luck, **maybe triple the Run 2 pp luminosity**.


| Detector | Target [fb ⁻¹] | Run 3 baseline [fb ⁻¹] | Run 3 + 1 year [fb ⁻¹] | | Leveled luminosity [10 ³⁴ cm ⁻² s ⁻¹] |
|-------------|----------------------------|------------------------------------|------------------------------------|--|---|
| ATLAS & CMS | 160 | 190 | 270 | | 2 |
| LHCb | 25 | 20 | 28 | | 0.15 - 0.2 |
| ALICE | 0.2 | 0.13 | 0.18 | | 0.0013 |

Preliminary numbers ... work in progress


High Lumi LHC (HL-LHC)

| Parameter | Nominal LHC (design report) | HL-LHC 25ns (standard) |
|---|--------------------------------|---------------------------|
| Beam energy in collision [TeV] | 7 | 7 |
| N_b | 1.15E+11 | 2.2E+11 |
| n_b^{12} | 2808 | 2760 |
| Beam current [A] | 0.58 | 1.1 |
| Half Crossing angle [μ rad] | 142.5 | 250 |
| Minimum β^* [m] | 0.55 | 0.15 |
| ϵ_n [μ m] | 3.75 | 2.50 |
| Total loss factor R0 without crab-cavity | 0.836 | 0.342 |
| Total loss factor R1 with crab-cavity | - | 0.716 |
| Virtual Luminosity with crab-cavity: $L_{\text{peak}} \cdot R1/R0$ [$\text{cm}^{-2} \text{s}^{-1}$] | - | 1.70E+35 |

- With more intense and more focused bunched LHC can get to the pileup of 400 in HL-LHC!
- Proton collisions will look like heavy ion collisions!
- New strategy for operations: luminosity leveling.




Level at $5 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
corresponds to ~ 140 pile up interactions



HL-LHC projection

Nominal HL-LHC






Luminosity detectors

- for integrated luminosity measurement
- but also for online luminosity measurement


Luminometers of the CMS experiment

- Any detector, which can provide particles hit rates can be used as a luminometer.
- A luminometer with a linear response produces a signal that is proportional to the instantaneous luminosity.
- In the CMS following Luminometers are used:
 - Pixel Detector
 - Forward calorimeter (HF)
 - Fast Beam Conditions Monitor (BCM1F)
 - Pixel Luminosity Telescope (PLT)




HF wedge


36 wedges in total




Unique inner carriage





4 PLT telescopes on titanium alloy SLM cassette



6 BCM1F sensors and dedicated ASICs mounted on PCB







BCM1L traces soldered onto PCB (C-shape)



- Specifications
- Sensors
- Electronics
- Readout
- Cooling
- Design
- Support structure and shape which fits with other CMS detectors
- Testing
- Installation
- Calibration
- Operations
- Data analysis
- Performance optimization

Detector assembly and tests in the laboratory



Happy BRIL team before installation 2015



Easter present 2015: beam splashes at LHC!




CMS Preliminary


Beam Splash 2015, 450 GeV



- Splash event: a single bunch is dumped in a collimator just upstream of the experiment.
- Since many particles cross the sensor simultaneously the signal amplitude saturates.
- First data recorded by the VME ADCs: the response of diamond sensors in splash events.



Before installation (2017)






New generation of the detectors




BCM1F for Run3

[Planning talk](#)

- Desired Si sensors, but no leakage current into FE-ASIC
 - AC coupling
- New sensor design: AC coupled double diodes, external bias resistor.
- First batch with recent HGCal sensor production.
- PCB features active cooling and various other improvements




Installation 2021




Part of the back-end

- 48 channels of BCM1F are served by CAEN v1721 ADCs:
 - 8 channels 8 bit 500 MS/s Digitizer (2 ns sampling time);
- Custom made RHU modules.
 - Histogramming units recording every collision for online luminosity measurement.



ONLINE Luminosity and Background measurements



PROTON PHYSICS: STABLE BEAMS

Energy:

6800 GeV

I B1:

1.31e+13

I B2:

1.30e+13

Beta* IP1:

19.20 m

Beta* IP2:

19.20 m

Beta* IP5:

19.20 m

Beta* IP8:

24.00 m

Inst. Lumi [(ub.s)⁻¹]

IP1: 19.00

IP2: 0.25

IP5: 17.55

IP8: 1.52


FBCT Intensity and Beam Energy

Updated: 11:41:04



Instantaneous Luminosity

Updated: 11:41:05



Comments (11-Nov-2022 11:22:05)

Fill #2 (146b) for VdM scans

CMS part 2 (until ~2pm)

next: LHCb (~2.5h), ATLAS (~4h)

BIS status and SMP flags

Link Status of Beam Permits

B1

B2

true true

Global Beam Permit

true true

Setup Beam

false false

Beam Presence

true true

Moveable Devices Allowed In



true true

Stable Beams

true true

Luminosity calibration


Calibration of the luminometers




$$L = N_1 N_2 f n_b / A_{eff},$$

where N_1 , N_2 - number of protons in beams 1 and 2, f - LHC orbit frequency, n_b - number of colliding bunches, A_{eff} - effective area.

- **Van der Meer scan method** proposed by Simon van der Meer: determine A_{eff} by measuring rates as a function of the beams displacement. Scans obtained in X and Y planes separately.



Visible cross section measurement



- Analysis framework is used to fit beam overlap and calculate visible cross section - the effective cross-section seen by the luminometer:


$$A_{\text{eff}} = 2\pi \Sigma_1 \Sigma_2,$$

$$\sigma_{\text{vis}} = \frac{2\pi \Sigma_x \Sigma_y}{N_1 N_2 f n_b} \left(\frac{\text{Peak}_x + \text{Peak}_y}{2} \right).$$

- The distribution is fitted with a Double Gaussian + Constant fit
- σ_{vis} is the important quantity, since it is used than for the data taking fills as the cross section from which the actual luminosity is measured.



Example of the Van der Meer scans program

- **Very complete pp VdM program, Fill 6868, July 2018:**
 - 7 pairs of VdM scans (3 pairs are Imaging scans with one beam fixed), 2 offset scans, 5 emittance scans;
 - length scale calibration with fixed and alternating separation;
 - 2 super separation measurements of background.






Special emittance scans for performance monitoring

- From 2017 CMS is using emittance scans – mini Van der Meer scans performed in 9 steps of beam separation.
- In 2017 it was demonstrated that emittance scans are powerful tools to monitor the detector performance.
- In 2018 from the first days with beams in the LHC emittance scans were used to cross check performance of the luminometers and correct calibrations for online purposes.
- Excellent online web-monitors allowed to display updating in real time luminosity in control room and accessible remotely.



Application of emittance scans

- Wide range of Single Bunch Instantaneous Luminosity (SBIL) is covered in each LHC fill -> Emittance scans early in the fill and towards the end of the fill allow for non-linearity measurement of online luminometers in each fill.
- Through the whole year emittance scans are employed to track the performance of the detectors and correct for efficiency drops.



Дякую за увагу!

Давайте подивимось 3D модель CMS!

<https://indico.cern.ch/event/1215773/timetable/>

2nd ONLINE Ukrainian Teacher Programme

Nov 7, 2022, 4:30 PM → Nov 11, 2022, 6:30 PM Europe/Kiev


Zoom (CERN)

Oksana Zhyvotova (CERN), Daria Ternova (CERN), Oleksandr Yurov, Jeff Wiener (CERN)

Description This is the second ONLINE Ukrainian Teacher Programme. It features talks and virtual visits delivered via Zoom that include ample opportunities for teachers to interact with the lecturers and discuss open questions.

Please note: The timetable is shown in the timezone of Ukraine, which is one hour ahead of Geneva. (This means a session that starts at 4:30 pm in Ukraine starts at 3:30 pm in Geneva.)

ONLINE Ukrainian Teacher Programme:
<https://indico.cern.ch/e/UATP22online-2>

Videoconference  2nd ONLINE Ukrainian Teacher Programme [Please log in and register](#)

Contact teacher.programmes@cern.ch

FRIDAY, NOVEMBER 11

4:30 PM → 6:30 PM Session 5





4:30 PM **CMS детектор та вимірювання світимості** 1h

Speaker: Olena Karacheban (Rutgers State Univ. of New Jersey (US))

 Nov2022_Teachers...


3D interactive CMS detector

Спробуйте самі!



<https://opendata.cern.ch/visualise/events/cms>
<https://www.i2u2.org/elab/cms/event-display/#>

| detid | pos_1 | pos_2 |
|-----------|---------------------------|---------------------------|
| 604017432 | 1.11126,-0.443558,6.07685 | 1.13947,-0.453736,6.22384 |
| 604017432 | 1.17623,-0.387179,6.07674 | 1.20609,-0.395927,6.22374 |



Спробуйте самі!

Text Documents

DETECT LANGUAGE ENGLISH UKRAINIAN RUSSIAN UKRAINIAN ENGLISH RUSSIAN

All scientific publications are there for you with translator google translate!


Using the golden decay channel to understand the production of the Higgs boson
The standard model of particle physics is currently the best way to describe interactions of fundamental particles that make up our Universe. It has been tested over many decades and many experiments, and the standard model manages to predict everything accurately. On 4th July 2012, the standard model achieved another great victory when CMS and ATLAS experiments announced the discovery of a new particle. The newly discovered particle turned out to be the Higgs boson, an essential piece in the standard model puzzle that explains the origin of the mass.

Усі наукові публікації для вас з перекладачем google translate!

Використання золотого каналу розпаду для розуміння утворення бозона Хіггса
Стандартна модель фізики елементарних частинок на даний момент є найкращим способом опису взаємодії фундаментальних частинок, які складають наш Всесвіт. Вона була перевірена протягом багатьох десятиліть і багатьох експериментів, і стандартна модель вдається передбачити все точно. 4 липня 2012 року стандартна модель здобула ще одну велику перемогу, коли експерименти CMS і ATLAS оголосили про відкриття нової частинки. Нещодавно відкрита частинка виявилася бозоном Хіггса, важливою частиною головоломки стандартної моделі, яка пояснює походження маси.

https://translate.google.com


By CMS Collaboration




Using the golden decay channel to understand the production of the Higgs boson

The standard model of particle physics is currently the best way to describe interactions of fundamental particles that make up our Universe. It has been tested over many decades and many experiments, and the standard model manages to predict everything accurately. On 4th July 2012, the standard model achieved another great victory when CMS and ATLAS experiments announced the discovery of a new particle. The newly discovered particle turned out to be the Higgs boson, an essential piece in the standard model puzzle that explains the origin of the mass.

Full Run 1 + Run 2



Run 1 alone



https://cms.cern/news/using-golden-decay-channel-understand-production-higgs-boson

Links and literature

LHC: <https://home.cern/science/accelerators/large-hadron-collider>

CERN Animation of CERN accelerator network: <https://youtu.be/RDdPuL-uOQc>

CMS News: https://cms.cern/cms-updates?field_article_type_target_id=382

Higgs boson <https://home.cern/science/physics/higgs-boson>

Higgs boson PDG <https://pdg.lbl.gov/2019/reviews/rpp2018-rev-higgs-boson.pdf>

Higgs article CMS: <https://cms.cern/news/using-golden-decay-channel-understand-production-higgs-boson>

Luminosity Werner Herr, Particle Colliders, CAS 2012, Granada

https://indico.cern.ch/event/173359/contributions/275970/attachments/218796/306434/luminosity_slides.pdf

CMS detector homepage: <https://cms.cern/detector>

HL-LHC by F.Moortgat <https://www.youtube.com/watch?v=s0toLfMFTIU>

Top quarks mass <https://home.cern/news/news/physics/run-top-quark-run>

Muons: <https://indico.cern.ch/event/426323/contributions/1048167/attachments/907573/1281134/muon.pdf>

LCH page 1: <https://op-webtools.web.cern.ch/vistar/vistars.php>

How to make your own cloud chamber: <https://home.cern/news/news/experiments/how-make-your-own-cloud-chamber>

Scoollab: <https://scoollab.web.cern.ch>

Cosmic Pi https://indico.cern.ch/event/555003/attachments/1305253/1951255/Cosmic_Pi_talk_ideasquare_070716.pdf