Contribution ID: 24

Type: Oral

## Impact of final-state de-excitation on modeling neutrino-nucleus interactions

Friday 25 August 2023 09:50 (20 minutes)

As neutrino oscillation physics enters the precision era, the modeling of neutrino-nucleus interactions constitutes an increasingly challenging source of systematic uncertainty for new measurements. To confront such uncertainties, a new generation of detectors is being developed to measure the exclusive final state of neutrino interaction. Precise simulations of the nuclear effects on the final-state nucleons are needed to fully benefit from the improved detector capabilities.

To address this problem, we focused on the re-interactions of final state nucleons with the nuclear medium (FSI) by comparing NuWro and INCL cascade models and considering the de-excitation model from ABLA. INCL is an evolved nuclear cascade code primarily designed to simulate nucleon-, pion- and light-ion-induced reactions on nuclei. ABLA is a de-excitation code coupled to INCL that can simulate all the particles emitted by de-excitation. This is the first detailed study using the INCL model in the framework of neutrino physics and the first study investigating the effect of de-excitation on the hadronic final state of neutrino interactions. The results feature various novelties, including the production of nuclear clusters (e.g., deuterons,  $\alpha$  particles) in the final state and the release of removal energy during the de-excitation stage.

We present a characterization of the hadronic final state after FSI and de-excitation, comparisons to available measurements of transverse kinematic imbalance, an assessment of the observability of nuclear clusters, and the impact of de-excitation on the energy released around the vertex.

Primary author: ERSHOVA, Anna (Université Paris-Saclay (FR))
Presenter: ERSHOVA, Anna (Université Paris-Saclay (FR))
Session Classification: parallel (room#301)

Track Classification: WG2: Neutrino Scattering Physics