



Contribution ID: 228

Type: not specified

Detecting Superradiant Dark Photon Strings in Gravitational Wave Experiments

Monday 8 May 2023 15:30 (15 minutes)

Black hole superradiance is a phenomenon in which, purely through gravitational interactions, light bosons are spontaneously produced outside spinning black holes. Through this effect, an exponentially growing Bose-Einstein condensate cloud forms around the black hole. In the case of dark photon superradiance, the cloud can form into strings of magnetic flux analogous to Abrikosov vortices in type II superconductors. Once formed, these strings are ejected from the black hole as loops of flux in a “stringy bosonova” event and travel across the galaxy at relativistic speeds. We discuss the prospects of detecting such a string here on earth. In particular, we consider the effect of a broken $U(1)_{B-L}$ gauge boson string passing through gravitational wave detectors such as LIGO and MAGIS. We show that the effect of the string on the detector is independent of the gauge coupling, g , and argue that, in some regions of the parameter space, these strings can produce a noticeable signal in the detector.

Primary author: RISTOW, Clayton (University of Maryland)

Co-authors: HOOK, Anson (University of Maryland); BRZEMINSKI, Dawid (University of Maryland - College Park); HUANG, Junwu (Perimeter Institute)

Presenter: RISTOW, Clayton (University of Maryland)

Session Classification: BSM XIII

Track Classification: BSM