



Contribution ID: 131

Type: not specified

# Multi-photon decays of the Higgs boson at the LHC

Tuesday 9 May 2023 15:00 (15 minutes)

Many new physics scenarios predict multi-photon Higgs resonances. One such scenario is the dark axion portal. The primary decay chain that we study is the Higgs to dark photon ( $\gamma_D$ ) pairs that subsequently decay into a photon and an axion-like particle ( $a$ ). The axion-like particles then decay into photon pairs. Hence, the signal is a six-photon Higgs decay:  $h \rightarrow \gamma_D \gamma_D \rightarrow 2 \gamma 2 a \rightarrow 6 \gamma$ . However, depending on the relevant kinematics, the photons can become well-collimated and appear as photon-jets (multiple photons that appear as a single photon in the detector) or  $\xi$ -jets (non-isolated multi-photon signals that do not pass the isolation criterion). These effects cause the true six-photon resonance to appear as other multi-photon signals, e.g., four or two photons. The four photon signal is particularly interesting. These events mainly occur when the photons from the axion are collimated into a photon jet. The apparent decay of the dark photon is then  $\gamma_D \rightarrow \gamma a \rightarrow \gamma + \gamma$ -jet. This decay seems to violate the Landau-Yang Theorem at the detector level since the dark photon appears to decay into a pair of massless photons. We explore and examine the multi-photon signals that could appear at the Large Hadron Collider (LHC). The mass regions where two, four, and six-photon resonances dominate are determined. Some additional signal categories involving  $\xi$ -jets are considered. All of these multi-photon signals provide excellent footing to explore new physics at the LHC and beyond.

**Primary authors:** LEE, Hye-Sung (KAIST); LEWIS, Ian; LANE, Samuel (KAIST)

**Presenter:** LANE, Samuel (KAIST)

**Session Classification:** SM III