


# Study of the $K^+ \rightarrow e^+ \nu_e e^+ e^-$ Decay with the NA62 Experiment<sup>1</sup>

**NA62**  
P326



**Zimányi School**  
5-9. 12. 2022.

Anna Fehérkuti  
CERN Summer Student 2022 (27. 6. - 23. 9.)  
Supervisors: Francesco Brizioli, Monica Pepe  
EP-UFT, Small Medium Expt



1 / 18

<sup>1</sup> <https://cds.cern.ch/record/2835579/>



# Motivation: K-physics

- $K^+ \rightarrow l^+ \nu$  described by Chiral Perturbation Theory (ChPT)  
→ test & inputs

- Decay amplitude includes:




- Inner Bremsstrahlung (IB)  
well predicted by  $K^+ \rightarrow l^+ \nu$

- Structure-Dependent components (SD): form factors ( $F_{A'}$ ,  $F_{V'}$ ,  $R$ )


- General  $K^+$ -decay sensitive to  $F_{A'}$ ,  $F_{V'}$

- $R$  contributes only to decays with  $e^+ e^-$  from  $\gamma^*$

- $K^+ \rightarrow e^+ \nu_e e^+ e^-$  (Ke2ee): SD > IB (←  $e^-$ -helicity suppression)



# About Branching Ratios



- Absolute measurement (of  $N^{\text{channel of interest}}/N^{\text{all}}$ ) impossible (interesting vs *everything*?)
- Normalization channel  $Br_2$  from PDG:  $Br_1/Br_2$ 
  - Likely process  $\rightarrow$  small external uncertainty (propagated, but negligible vs syst/stat)
  - Similar process  $\rightarrow$  small systematic error (many uncertainty factors fall out)
- Which one is better in this case?

# Uncertainties



$$\frac{Br^{signal}}{Br^{norm}} = \frac{N^{signal}}{N^{norm}} \cdot \frac{\varepsilon^{norm}}{\varepsilon^{signal}} = \frac{N^{signal}}{N^{norm}} \cdot \frac{Acc^{norm}}{Acc^{signal}} \cdot \frac{Trig^{norm}}{Trig^{signal}}$$

- *Br*: branching ratio
- *N*: actual measured counts
- $\varepsilon$ : selection efficiency
- *Acc*: acceptance, efficiency of the offline selection
  - From Monte Carlo (MC)
- *Trig*: efficiency of the online trigger selection
  - Different masks for signal and normalization
- Same cuts: *Acc* of signal vs normalization cancel
- Perfect MC: *N* & *Acc* balance each other






# Studied Processes



# The NA62 Experiment



- Kaon identification
- Tracking the beam
- Collimator vs *upstream*  
→ Fiducial Volume (FV): decay
- Photon vetoes (vs  $\pi^0$ )
- Spectrometry of charged ( $p$ )
- Cherenkov radius of charged ( $\beta$ )
- Electromagnetic & hadronic calorimeters
- Muon vetoes



# Event Selection I.



- General conditions:

- 1 single 3 track event
- Precise enough vertex ( $\chi^2 < 25$ ) of charge +1, in FV ( $z \in [105, 180]$  m) within 6 ns wrt trigger (vtx: CHOD vs trig: RICH)
- Opposit-charged particles in time wrt trigger:  
 $|t_{\text{NewCHOD}}^1 - t_{\text{NewCHOD}}^2| < 2 \text{ ns}, |t_{\text{NewCHOD}}^i - t_{\text{CHOD}}| < 2 \text{ ns}$
- Tracks in detector acceptance (STRAW, RICH, CHOD, NewCHOD, LKr)
- Reasonable track separation (15 mm in each STRAW chamber, 200 mm in LKr plane)
- Extra activity vetos:  $\gamma$ s,  $\mu$ s (reject event if activity within 2 ns wrt vertex)
- Good association between KTAG-GTK & RICH-CHOD:  
 $|t_{\text{GTK}} - t_{\text{KTAG}}| < 1.4 \text{ ns}, |t_{\text{vertex}} - t_{\text{RICH}}| < 2 \text{ ns}$
- Vertex-building from the three downstream tracks and the GTK track, where the GTK candidate gives the minimal  $\chi^2_{\text{vertex}}$
- Momentum of each track separately  $\in [8, 50]$  GeV
- 3-track momentum  $< 78$  GeV
- HLT (L1): KTAG was ok, no exotics in STRAW

# Event Selection II.



- Signal selection:
  - Particle identification (PID):
    - $e^-$  probability from calorimetric BDT  $> 0.5$  for the positive tracks (Boosted Decision Tree, BDT: neural algorithm)
    - $e^-$  RICH likelihood  $> 0.5$  for the positive (!) tracks
    - No EoP (from LKr) condition needed (EoP  $> 0.9$  [3])!
  - Kinematics:
    - Neutrino momentum (lower boundary):  $p_\nu > 200$  MeV
    - $p^T$  in GTK (lower boundary):  $p_{GTK}^T > 8$  MeV
    - Electron-positron invariant mass (lower boundary):  $m_{e^-,e^+} > 140$  MeV
      - Theory: vs divergence in the decay rate due to the small-energy  $\gamma$
      - Experimentally: vs  $K^+ \rightarrow e^+ \nu \pi^0$ ,  $\pi^0 \rightarrow e^- e^+ \nu$
    - Missing mass (upper boundary):  $m_{miss}^2 < 0.03$  GeV<sup>2</sup>
  - Trigger mask4 (“di-electron”), downscaling of 8:  
extra condition (over mask5\*) on LKr total energy (minimum 20 MeV)







# $m_{\text{miss}}^2$ vs $m_{e^-,e^+}$

- Well-peaked distribution → suitable for selection
- $\nu \rightarrow 0 = m_{\text{miss}}^2$

NA62 full data 2017-2018

NA62 MC v3.1.0-v3.1.3



2 way for  $m_{e^-,e^+}$  (2 positrons):

Choose the smaller

→ minimum cut on  $m_{e^-,e^+}$ : cut on both

# Event Selection III.




- Normalization selection:
  - Kinematics:
    - Kaon invariant mass:  $|m_{3\pi} - m_{K^+}| < 4 \text{ MeV}$   
(check if GTK was ok first)
  - No PID needed! (clean enough sample)
  - Separating data (events already identified as signal shall not be analyzed again):  
EoP < 0.9
  - Trigger mask5 (“multi-track”), downscaling of 100:
    - RICH was ok
    - Good newCHOD candidates



# Kaon invariant mass




NA62 full data 2017-18



- Well-peaked distribution  $\rightarrow$  suitable for selection
- Official value [4]:  $493.677 \pm 0.013$  MeV

# Stability studies: kaon invariant mass

Cut on kaon invariant mass



- Normed to the chosen cut
- Uncertainty of the central value: all stat + syst
- Uncertainty of the other values: relative to the central
- (On the following plots as well...) 12 / 18


# Stability studies: $m_{e^+e^-}$




- Differently zoomed:

- Too loose requirement  $\rightarrow$  misidentified signals
- Too strict  $\rightarrow$  hardly remaining events: big uncert.:


Cut on electron-positron invariant mass  $m_{e^+e^-}$



Cut on electron-positron invariant mass  $m_{e^+e^-}$




Cut on electron-positron invariant mass  $m_{e^+e^-}$



# Stability studies: $m_{\text{miss}}^2$



Cut on  $m_{\text{miss}}^2$



- Cut has to be where it is stable enough
- Not in the range of uncertainty: also almost different order of magnitude



# Trigger efficiency

- Wrt selection
  - From data: control (CTRL) data needed
    - Signal:  $\text{mask4}/\text{CTRL} = 708/11$  (too low stat)
    - Normalization:  $\text{mask5}/\text{CTRL} = 91.7\%$
  - From MC, emulating L0 triggers as well:
    - Signal (RICH, NewCHOD, LKr): 92.6%
    - Normalization (RICH, NewCHOD): 91.3%
    - ratio:  $(98.5 \pm 0.8)\%$
- /extra condition in mask4 and not in mask5 (LKr20): very small inefficiency/



# Summary: Results in Numbers

- Values from the literature:

- $Br_{normalization}^{PDG}: (5.583 \pm 0.024)\% [4]$

- $Br_{signal}^{theory}(m_{e^-,e^+} > 140 \text{ MeV}): 3.39 \cdot 10^{-8} [1]$

- $Br_{signal}^{measurement}(m_{e^-,e^+} > 140 \text{ MeV}): (2.91 \pm 0.34) \cdot 10^{-8} [2]$

- My analysis:

- $Br_{signal}(m_{e^-,e^+} > 140 \text{ MeV}): \mathbf{(3.13 \pm 0.12) \cdot 10^{-8}}$



	Signal	Normalization	Ratio
$N$	$708 \pm 26.61$	$230419472 \pm 15180$	$(3.073 \pm 0.116) \cdot 10^{-6}$
$Acc$	$0.02837 \pm 0.00015$	$0.06300 \pm 0.00008$	$2.221 \pm 0.012$
$\epsilon$	$0.9265 \pm 0.0069$	$0.9126 \pm 0.0017$	$0.9851 \pm 0.0076$


# Outlook

- Examining background contamination in signal case (cca. 20 events vs 778 [3])
- Analysis on bigger data





**Thank You for the attention!**



**Time for questions!**



# Backup slides



# Previous studies

Table 7: Theoretical values for the branching ratios for the decay  $K^+ \rightarrow e^+ \nu_e e^+ e^-$  for various cuts.

- Theory [1]:

- $z$ :  $m_{e^-,e^+}$

	tree level	form factors as given by CHPT
full phase space	$\approx 4 \cdot 10^{-9}$	$1.8 \cdot 10^{-7}$
$z, z_1 \geq 10^{-3}$	$3.0 \cdot 10^{-10}$	$1.22 \cdot 10^{-7}$
$z, z_1 \geq (50 \text{ MeV}/M_K)^2$	$5.2 \cdot 10^{-11}$	$8.88 \cdot 10^{-8}$
$z, z_1 \geq (140 \text{ MeV}/M_K)^2$	$2.1 \cdot 10^{-12}$	$3.39 \cdot 10^{-8}$

- Experimental results (BNL, 2002) [2]:  $N_{\text{signal}} = 410$   
(including 10% background contamination)

$$Br(m_{e^-,e^+} > 140 \text{ MeV}) = [291 \pm 16(\text{stat}) \pm 17(\text{syst}) \pm 0.7(\text{ext from model})] \cdot 10^{-10}$$

→ NA62: data collected 2016-2021

→ Full 2017-18 sample

+ v3.1.3 MC: kenuee, k3pi, k2pi.pi0d






# Stability studies






# Stability studies: BDT PID

Cut on calorimetric PID likelihood




# Stability studies: RICH PID

Cut on RICH PID likelihood




# Stability studies: $p_\nu$

Cut on  $p_\nu$





# Stability studies: $p_T^{\text{GTK}}$



# Parts of the NA62 Experiment





# KTAG

- Kaon-tagging vs proton
- PMTs
- Front-end readout
- Flashed with  $N_2$





# CEDAR

- Differential Cherenkov with KTAG
- Chromatic correctors + Mirrors
- 1.6 m
- $N_2$




# Safe Volume

- For emergency cases: leakage on beam pipe
- N<sub>2</sub> into CEDAR vs mechanical wave





# Magnets






# GTK

- GigaTracker: beam
- Between dipoles
  - 4 stations
- Si pixel



# Collimator




- Rainbow :)
- Vs upstream





# CHANTI

- Charged ANTIcounter
- Hodoscope: scintillators
- Veto vs upstream





# Beginning of the Fiducial Volume





# LAVs

- Vetoes against photons
- Leadglass scintillators





# STRAW

- Spectrometer
  - p of particle



- 4 stations
- 35 m



# Strong magnet for STRAW

- 0.9 Tm: horizontal momentum kick of 270 MeV/c
  - 75 GeV/c beam deflected too, by -3.6 mrad





# RICH

- Ring Imaging Cherenkov: Ne
- $\beta$  of the charged particles




- Mirror mosaic
- PM disk





# IRC


- Intermediate Ring Calorimeter
- Photon-veto
- Pb / scintillator Shashlyk





# (New)CHOD


- Hodoscope: scintis
- Time  $\rightarrow$  minimum bias trigger





# LKr

- EM calorimeter from NA48
- Accordeon Cu ribbons
- 9 m<sup>3</sup> liquid Kr



# MUV1,2

- Sampling hadronic calorimeters
- Fe / scintillators





# MUV3



- Extra iron before it
- Only muons
- Plastic scintillators






# HASC, SAC

- HASC:
  - Vs multitrack ( $\pi^+$  > 50 GeV)
  - Sampling calorimeter
- SAC ~ last LAV:
  - Small angle calorimeter
  - Photon veto
  - Shashlyk





# About the theoretical background



**BSM**



# Motivation of NA62: BSM Probes

- CKM (Wolfenstein) → unitarity triangle

$$\lambda = \frac{|V_{us}|}{\sqrt{|V_{ud}|^2 + |V_{us}|^2}}$$

- Area related to the amount of CPV

$$A = \frac{|V_{cb}|}{\lambda^2 \sqrt{|V_{ud}|^2 + |V_{us}|^2}}$$

- (Semi)leptonic kaon decays:  $|V_{cb}|$ ,  $|V_{us}|$ ,  $\gamma$


$$\rho = \frac{\text{Re}(V_{ub})}{A\lambda^3}$$

- $K^+ \rightarrow \pi^+ \nu \bar{\nu}$

$$\bar{\rho} = \rho \left( 1 - \frac{\lambda^2}{2} + \mathcal{O}(\lambda^4) \right)$$


$$\eta = \frac{\text{Im}(V_{ub})}{A\lambda^3}$$

- Inconsistency between channels of measurement:



$$\bar{\eta} = \eta \left( 1 - \frac{\lambda^2}{2} + \mathcal{O}(\lambda^4) \right)$$


- Hint to BSM



$$V_{CKM} = \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4)$$



# Unitarity triangle




- Restriction on matrix elements by unitarity:

$$\mathbb{1} \equiv \mathbf{V}_{\text{CKM}} \cdot \mathbf{V}_{\text{CKM}}^\dagger = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \cdot \begin{pmatrix} V_{ud}^* & V_{cd}^* & V_{td}^* \\ V_{us}^* & V_{cs}^* & V_{ts}^* \\ V_{ub}^* & V_{cb}^* & V_{tb}^* \end{pmatrix} =$$


$$\begin{pmatrix} V_{ud}V_{ud}^* + V_{us}V_{us}^* + V_{ub}V_{ub}^* & V_{ud}V_{cd}^* + V_{us}V_{cs}^* + V_{ub}V_{cb}^* & V_{ud}V_{td}^* + V_{us}V_{ts}^* + V_{ub}V_{tb}^* \\ V_{cd}V_{ud}^* + V_{cs}V_{us}^* + V_{cb}V_{ub}^* & V_{cd}V_{cd}^* + V_{cs}V_{cs}^* + V_{cb}V_{cb}^* & V_{cd}V_{td}^* + V_{cs}V_{ts}^* + V_{cb}V_{tb}^* \\ V_{td}V_{ud}^* + V_{ts}V_{us}^* + V_{tb}V_{ub}^* & V_{td}V_{cd}^* + V_{ts}V_{cs}^* + V_{tb}V_{cb}^* & V_{td}V_{td}^* + V_{ts}V_{ts}^* + V_{tb}V_{tb}^* \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- 3 angle per triangle
  - 6 triangles vs same: 1 CPV complex phase
- Area of the triangle  $\rightarrow$  CP violation
- Inconstistency in predictions in  $(\bar{\rho}, \bar{\eta})$  plane  $\rightarrow$  BSM

# Unitarity triangle



- Tree-level: semi-leptonic K-, B-decays
  - $|V_{us}|$ ,  $|V_{cb}|$ ,  $|V_{ub}|$  &  $R_b$  within SM
- Other measurement channels  $\rightarrow$  apex A on the plane
  - Loop-induced decays & transitions
  - CP violating B decays





# References

- [1] J. Bijnens, G. Ecker, and J. Gasser. “Radiative semileptonic kaon decays”. In: Nucl.Phys. B 396 (1993), pp. 81–118. doi: 10.1016/0550-3213(93)90259-R. arXiv:hep-ph/9209261 (cit. on p.2).
- [2] A. A. Poblaguev et al. “Experimental study of the radiative decays  $K^+ \rightarrow \mu^+ \nu e^+ e^-$  and  $K^+ \rightarrow e^+ \nu e^+ e^-$ ”. In: Phys. Rev. Lett. 89 (2002), p.061803. doi:10.1103/PhysRevLett.89.061803. arXiv: hep-ex/0204006 (cit. on p. 3).
- [3] G. Romolini, Study of the  $K^+ \rightarrow e^+ \nu e^+ e^-$  decay with the NA62 experiment at CERN.
- [4] M. Tanabashi et al. (Particle Data Group), Charged Kaon Mass, Phys. Rev. D, 2018, 98, 030001.
- <https://www.particlezoo.net/>
- <https://www.facebook.com/NA62Experiment>
- <https://phys.org/news/2020-07-cern-evidence-ultra-rare-physics.html>
- <https://cernbox.cern.ch/index.php/s/Q6l8onTbmJtS0ID>
- <https://www.mdpi.com/2218-1997/4/11/119/htm>
- <https://www.istockphoto.com/de/fotos/penguin-thinking>
- <https://www.pinterest.ch/pin/cute-penguin-jumping-cartoon-notepad-baby-gifts-child-new-born-gift-idea-diy-cyo-special-unique-design--733664595519053629/>
- [https://en.wikipedia.org/wiki/Penguin\\_diagram](https://en.wikipedia.org/wiki/Penguin_diagram)
- <https://www.nature.com/articles/nphys3464>
- <https://www.physik.uzh.ch/dam/jcr:8cb19cb1-d67a-4a44-9a73-c26871988dd8/chap05.pdf>
- <https://arxiv.org/abs/hep-ph/0304132v2>