

CAS SCHOOL, SEP 2023



INTRODUCTION TO SPECIAL RELATIVITY


Dr. Irina Shreyber, PhD



FORCES

| Force | Acts on | Strength | Range | Boson |
|-----------------|--|------------|-------------------|------------|
| Strong | Quarks and particles containing quarks | 10^4 | $\sim 10^{-14}$ m | g |
| Electromagnetic | Electrically charged particles | 10^2 | ∞ | γ |
| Weak | All particles | 10^{-2} | $\sim 10^{-17}$ m | W^\pm, Z |
| Gravitational | All particles | 10^{-34} | ∞ | ? |

Note: strength depends on distance or momentum transfer!



**Every day example:
plane, airport, bad weather**

The author consents to the photographic, audio and video recording of this lecture at the CERN Accelerator School. The term “lecture” includes any material incorporated therein including but not limited to text, images and references.

The author hereby grants CERN a royalty-free license to use his image and name as well as the recordings mentioned above, in order to post them on the CAS website.

The author hereby confirms that to his best knowledge the content of the lecture does not infringe the copyright, intellectual property or privacy rights of any third party. The author has cited and credited any third-party contribution in accordance with applicable professional standards and legislation in matters of attribution. Nevertheless the material represent entirely standard teaching material known for more than ten years. Naturally some figures will look alike those produced by other teachers.

CAS Website

These slides and the video will be available
the CAS school website

Books

- Jurgen Freund, "Special Relativity For Beginners"
- James H. Smith, "Introduction to Special Relativity"
- Mario Conte, William W. MacKay, "An Introduction to the Physics of Particle Accelerators"



**IN THIS LECTURE,
WE WILL LEARN...**

THE TRANSITION

in thinking that led from Galilean Relativity to the Special Theory of Relativity in 1905.

The postulates

of special relativity, which are the basis of the mathematics of the framework.


The consequences

of the postulates.

NEWTON'S PRINCIPLE OF RELATIVITY

GALILEO GALILEI IN 1632, AND LATER BY NEWTON


“The motions of bodies included in a given space are the same among themselves, whether that space is at rest or moves uniformly forward in a straight line.”




GALILEAN TRANSFORMATION

At the time of Newton the relation of the coordinates between two systems in motion with relative velocity v , was defined by the Galilean transformation of motion

Fixed frame



Moving frame



$$\begin{aligned}x' &= x - vt \\y' &= y \\z' &= z \\t' &= t\end{aligned}$$


$$\Rightarrow \begin{aligned}\mathbf{r}' &= \mathbf{r} - \mathbf{v}t \\t' &= t\end{aligned}$$

with $\mathbf{r} = (x, y, z)$.

MAXWELL EQUATIONS


DIFFERENTIAL FORM

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}$$




GAUSS'S LAW FOR E

$$\nabla \cdot \mathbf{B} = 0$$



GAUSS'S LAW FOR B

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$



FARADAY'S LAW
for time-varying
magnetic fields

$$\nabla \times \mathbf{B} = \mu_0 \left(\mathbf{J} + \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \right)$$



**AMPERE(-MAXWELL)
LAW**
for time-varying
electric fields

THE PROBLEM WITH GALILEAN TRANSFORMATION

▶
$$\left(\frac{\partial^2}{\partial x'^2} + \frac{\partial^2}{\partial y'^2} + \frac{\partial^2}{\partial z'^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t'^2} \right) \Psi = 0$$

Maxwell wave equation

▶
$$x = x' - vt, \quad y' = y, \quad z' = z, \quad t' = t$$

Galilean transformation

▶
$$\left(\left[1 - \frac{v^2}{c^2} \right] \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} + \frac{2v}{c^2} \frac{\partial^2}{\partial x \partial t} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \Psi = 0$$

Maxwell wave equation

Maxwell Equations. CONSEQUENCES.



IF THERE IS A DISTURBANCE IN THE FIELD SUCH THAT LIGHT IS GENERATED, THESE ELECTROMAGNETIC WAVES GO OUT IN ALL DIRECTIONS EQUALLY AND AT THE SAME SPEED:


$c = 299\,792\,458$ m/s in vacuum
("celeritas" = speed)

IF THE SOURCE OF THE DISTURBANCE IS MOVING, THE LIGHT EMITTED GOES THROUGH SPACE AT THE SAME SPEED **C**.

This is analogous to the case of sound, the speed of sound waves being likewise independent of the motion of the source.

MICHELSON-MORLEY EXPERIMENT (1887)

The goal was to determine the absolute velocity of the earth through this hypothetical “ether”:



$$B \rightarrow E : ct_1 = L + ut_1 \Rightarrow t_1 = L / (c - u)$$

$$E \rightarrow B : ct_2 = L - ut_2 \Rightarrow t_2 = L / (c + u)$$

$$t_1 + t_2 = \frac{2L}{(1 - u^2/c^2)}$$

$$B \rightarrow C : (ct_3)^2 = L^2 + (ut_3)^2 \Rightarrow t_3 = L / \sqrt{c^2 - u^2}$$

$$C \rightarrow B : t_4 = t_3$$


If there is an “aether drift” then

$$t_1 + t_2 \neq t_3 + t_4$$

The apparatus was amply sensitive to observe such an effect, but no time difference was found — the velocity of the earth through the aether could not be detected. The result of the experiment was null.

MICHELSON-MORLEY EXPERIMENT (1887)

The goal was to determine the absolute velocity of the earth through this hypothetical “ether”:




Light travels at a fixed and constant speed in any medium, regardless of the relative velocity of the light-source and the light-observer →

This is unlike any other phenomenon described in mechanics, and implies that Newton's Mechanics is incomplete.

MICHELSON-MORLEY EXPERIMENT (1887)


The goal was to determine the absolute velocity of the earth through this hypothetical “ether”:



No medium is required for light to propagate; unlike a mechanical oscillatory phenomenon (wave), to exist light requires no medium to be distorted →
this implies Maxwell's Equations are complete

MICHELSON-MORLEY EXPERIMENT (1887)

The goal was to determine the absolute velocity of the earth through this hypothetical “ether”:



These lessons would not be absorbed fully until 1905, when Albert Einstein published the definitive papers explaining how to reconcile mechanics, electricity and magnetism, and the Michelson-Morley experiment

"If the Michelson–Morley experiment had not brought us into serious embarrassment, no one would have regarded the relativity theory as a (halfway) redemption." Einstein


"COMPRESSION OF BODIES IN THE AETHER"

HENDRIK LORENTZ

replacement for the Galilean Relativity equations

THE EFFECTS OF THE AETHER ON BODIES IN MOTION

- Mechanical bodies would compress along the direction of motion in the aether, with a precise mathematical description for the process
- In transforming observations from the aether frame to other frames of reference, he would conceive of an alteration of time that also had a mathematical description



LORENTZ TRANSFORMATION


$$L_{\parallel} = L_0 \sqrt{1 - v^2/c^2}$$


$$x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}}$$

$$y' = y$$

$$z' = z$$

$$t' = \frac{t - vx/c^2}{\sqrt{1 - v^2/c^2}}$$





**EINSTEIN'S
PRINCIPLE OF
RELATIVITY**

IN ALBERT EINSTEIN'S ORIGINAL TREATMENT, IN 1905,
THE PRINCIPLE OF RELATIVITY IS BASED ON
TWO POSTULATES:

1. SPECIAL PRINCIPLE OF RELATIVITY:

The laws of physics are invariant (i.e. identical) in all inertial frames of reference (i.e. non-accelerating frames of reference).

2. INVARIANCE OF C:

The speed of light in a vacuum is the same for all observers, regardless of the motion of the light source or observer.

1905, Albert Einstein, "On the Electrodynamics of Moving Bodies".




BREAKDOWN THE POSTULATES

- **"EVENT"**
- **"FRAME OF REFERENCE"**
- **"SIMULTANEITY"**
- **"SPEED OF LIGHT"**



"EVENT"

**IS ANYTHING WITH A LOCATION IN SPACE AND
TIME**




"EVENT"

**IS ANYTHING WITH A LOCATION IN SPACE AND
TIME**



"EVENT"


**IS ANYTHING WITH A LOCATION IN SPACE AND
TIME**



The dot is at position $x=0$ m at time $t=0$ s


"EVENT"

**IS ANYTHING WITH A LOCATION IN SPACE AND
TIME**




"EVENT"

**IS ANYTHING WITH A LOCATION IN SPACE AND
TIME**




The dot is at position $x=2\text{m}$ at time $t=2\text{s}$




"FRAME OF REFERENCE"

***IS ANY OBJECT OR SYSTEM ALL OF WHOSE PARTS
MOVE AT THE SAME VELOCITY WITH RESPECT TO AN
AGREED-UPON REFERENCE POINT IN SPACE.***



Do the red dot and blue dot share the same or different frames of reference?



"FRAME OF REFERENCE"

***IS ANY OBJECT OR SYSTEM ALL OF WHOSE PARTS
MOVE AT THE SAME VELOCITY WITH RESPECT TO AN
AGREED-UPON REFERENCE POINT IN SPACE.***

Black
•

Blue
• →

← Red
•

Do the red dot and blue dot share the
same or different frames of reference
NOW?

"SIMULTANEITY"

***TWO EVENTS (OR MORE) ARE SAID TO BE SIMULTANEOUS
(THAT IS, TO POSSESS OF SIMULTANEITY), IF THEY ARE
OBSERVED TO OCCUR AT THE SAME MOMENT IN TIME***

Think really hard about
whether events are simultaneous,
and for whom
(which observers in which frames of
reference) they are simultaneous.



Modern Speed of Light

The speed of light, based on modern definitions of the meter and the second, is defined to be exactly 299,792,458m/s. Light travels roughly one foot in one billionth of a second (1ft/ns).

"THE SPEED OF LIGHT"

IT IS THE NUMBER OF METERS LIGHT CAN TRAVEL, ONCE EMITTED BY A SOURCE, IN A CERTAIN AMOUNT OF TIME.

Galileo Galilei: attempted to measure this by uncovering a lantern, having an assistant on a distant hill who uncovers their lantern upon seeing his, and upon seeing the assistant's lantern light he records the time for the round trip, taking into account human reaction time



Planck's constant \hbar and speed of light c :

$$\hbar \equiv \frac{h}{2\pi} \approx 1.055 \times 10^{-34} \text{ J s}$$

$$c \approx 2.998 \times 10^8 \text{ m/s}$$

Units in the international system:

$$[\hbar] = \frac{ML^2}{T} = \frac{\text{kg m}^2}{\text{s}}$$

$$[c] = \frac{L}{T} = \frac{\text{m}}{\text{s}}$$

"Natural" units:

$$\hbar = c \equiv 1 \quad ; \quad [\hbar] = [c] = 1$$

$$[e] = [\sqrt{\hbar c}] = [1] \quad ; \quad \alpha = \frac{1}{4\pi} \frac{e^2}{\hbar mc} = \frac{1}{4\pi} \frac{e^2}{mc^2} =$$

NATURAL UNITS

Basic unit: electronvolt (eV) \equiv energy gained by an electron in a potential difference of 1V:

$$[E, M, p] = \frac{ML^2}{T^2} = \text{eV}$$

$$1 \text{ GeV} = 10^9 \text{ eV} \approx M_p$$

| | I | II | III | |
|---------|-------------------|-------------------|-------------------|---------------|
| Quarks | 2.4 MeV u | 1.3 GeV c | 170 GeV t | 0 γ |
| | 4.8 MeV d | 104 MeV s | 4.2 GeV b | 0 g |
| | < 2 eV ν_L | < 2 eV ν_M | < 2 eV ν_H | 91 GeV Z |
| Leptons | 0.5 MeV e | 16 MeV μ | 1.8 GeV τ | 80 GeV W |
| | | | | 126 GeV H |
| | | | | Bosons |

| Quantity | Conversion factor | Natural units | IS units |
|----------|---|-------------------|----------------------|
| Mass | 1 kg = 5.61×10^{26} GeV | GeV | GeV/c ² |
| Length | 1 m = 5.07×10^{15} GeV ⁻¹ | GeV ⁻¹ | $\hbar c/\text{GeV}$ |
| Time | 1 s = 1.52×10^{24} GeV ⁻¹ | GeV ⁻¹ | \hbar/GeV |

LORENTZ TRANSFORMATION

VELOCITY:

$$\beta = \frac{v}{c} \in [0, 1]$$

LORENTZ FACTOR:

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}} \in [1, \infty)$$



LORENTZ TRANSFORMATION

$$x' = \gamma (x - \beta ct)$$

$$y' = y$$

$$z' = z$$

$$ct' = \gamma (ct - \beta x)$$

$$x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}}$$

$$y' = y$$

$$z' = z$$


$$t' = \frac{t - vx/c^2}{\sqrt{1 - v^2/c^2}}$$

$$\mathbf{r}'_{\parallel} = \gamma (\mathbf{r}_{\parallel} - \beta ct)$$

$$\mathbf{r}'_{\perp} = \mathbf{r}_{\perp}$$

$$ct' = \gamma (ct - \mathbf{r}_{\parallel} \cdot \boldsymbol{\beta})$$

where \mathbf{r}_{\parallel} and \mathbf{r}_{\perp} are the components of \mathbf{r} w.r.t. $\boldsymbol{\beta}$ (or \mathbf{v})



SPECIAL RELATIVITY & FOUR-VECTORS

The space-time 4-vector

$$\vec{R} = \begin{bmatrix} ct \\ x \\ y \\ z \end{bmatrix} = \begin{bmatrix} ct \\ \vec{r} \end{bmatrix}$$

The energy-momentum 4-vector

$$\vec{P} = \begin{bmatrix} E \\ p_x c \\ p_y c \\ p_z c \end{bmatrix} = \begin{bmatrix} E \\ \vec{p}c \end{bmatrix}$$

Scalar product of two 4-vectors

$$\vec{R}_a = \begin{bmatrix} ct \\ \vec{r}_a \end{bmatrix} \quad \vec{R}_b = \begin{bmatrix} ct \\ \vec{r}_b \end{bmatrix} \quad \vec{R}_a \cdot \vec{R}_b = ct_a ct_b - \vec{r}_a \cdot \vec{r}_b$$

$$\vec{P}_a = \begin{bmatrix} E_a \\ \vec{p}_a c \end{bmatrix} \quad \vec{P}_b = \begin{bmatrix} E_b \\ \vec{p}_b c \end{bmatrix} \quad \vec{P}_a \cdot \vec{P}_b = E_a E_b - \vec{p}_a \cdot \vec{p}_b c^2$$

Length of the 4-vector squared

$$\vec{R} \cdot \vec{R} = (ct)^2 - (x^2 + y^2 + z^2)$$

$$\sqrt{P \cdot P} = \sqrt{E^2 - (pc)^2} = m_0 c^2$$

- Four-vector of electromagnetic current density:

$$j^\mu = (\rho, \vec{j})$$

- ρ : charge density
- \vec{j} : current density

- Four-vector of electromagnetic potential:

$$A^\mu = (V, \vec{A})$$

the electrical potential energy per charge is the electric potential.

- V : scalar electric potential
- \vec{A} : magnetic vector potential $\mathbf{B} = \nabla \times \mathbf{A}$

SPECIAL RELATIVITY & FOUR-VECTORS


A ray of light connects two events, $x_1^\mu = (ct_1, \vec{x}_1)$ et $x_2^\mu = (ct_2, \vec{x}_2)$:

$$\left((x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2 \right)^{\frac{1}{2}} = c (t_1 - t_2)$$

Consequence: the length of a four-vector is independent of the inertial frame:

$$s = c^2 t^2 - x^2 - y^2 - z^2 = \text{invariant}$$

Lorentz transformations = rotations and translations in space-time which leave s invariant.



$$x' = \gamma (x - \beta ct)$$

$$y' = y$$

$$z' = z$$

$$ct' = \gamma (ct - \beta x)$$


$$x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}}$$

$$y' = y$$

$$z' = z$$


$$t' = \frac{t - vx/c^2}{\sqrt{1 - v^2/c^2}}$$

$$\begin{bmatrix} ct' \\ x' \\ y' \\ z' \end{bmatrix} = \begin{bmatrix} \gamma & -\beta\gamma & 0 & 0 \\ -\beta\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} ct \\ x \\ y \\ z \end{bmatrix}$$



$$\begin{bmatrix} E' \\ p_x' c \\ p_y' c \\ p_z' c \end{bmatrix} = \begin{bmatrix} \gamma E - \beta \gamma p_x c \\ -\beta \gamma E + \gamma p_x c \\ p_y c \\ p_z c \end{bmatrix}$$

$$\begin{bmatrix} E' \\ p_x' c \\ p_y' c \\ p_z' c \end{bmatrix} = \begin{bmatrix} \gamma & -\beta \gamma & 0 & 0 \\ -\beta \gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} E \\ p_x c \\ p_y c \\ p_z c \end{bmatrix}$$



SPECIAL RELATIVITY & FOUR-VECTORS

- **Scalar product** between four-vectors:

$$x_{\mu}x^{\mu} = c^2t^2 - x^2 - y^2 - z^2 = c^2t^2 - \vec{x}^2$$

$$p_{\mu}p^{\mu} = \frac{E^2}{c^2} - p_x^2 - p_y^2 - p_z^2 = E^2/c^2 - \vec{p}^2 = m^2c^2$$

$$p_{\mu}x^{\mu} = Et - p_x x - p_y y - p_z z = Et - \vec{p}\vec{x}$$

- **Scalars** under Lorentz transformations, invariant when changing between inertial frames

- System of "natural" units:

$$x^{\mu} = (x^0, x^1, x^2, x^3) = (t, x, y, z) \quad ; \quad p^{\mu} = (p^0, p^1, p^2, p^3) = (E, p_x, p_y, p_z)$$
$$x_{\mu}x^{\mu} = t^2 - x^2 - y^2 - z^2 = t^2 - \vec{x}^2 \quad ; \quad p_{\mu}p^{\mu} = E^2 - p_x^2 - p_y^2 - p_z^2 = E^2 - \vec{p}^2 = m^2$$

LHC Energy LHC

The Einstein relationship for energy includes both the kinetic energy and rest mass energy for a particle.

$$E = mc^2$$

The kinetic energy of a high-speed particle can be calculated from $E_K = mc^2 - m_0c^2$

The mass of proton is 938,3 MeV/c²

- 1) What would be “relativistic mass”, m ?
- 2) Calculate Lorenz factor, γ ?
- 3) Calculate velocity, v ?
- 4) Calculate the energy of the rest of proton, E_0 ?

Let's take a look at γ (gamma) when the proton reaches LHC energy (7TeV per beam).

$$E_k = \gamma \cdot m_0 \cdot c^2 - m_0 \cdot c^2 \quad \text{Kinetic energy}$$

$$E_k = m_0 \cdot c^2 (\gamma - 1)$$

$$M_p = 938.3 \text{ MeV}/c^2 \quad m_0 \cdot c^2 = 9,383 \cdot 10^{-4} \text{ TeV}$$

$$7 \text{ TeV} = 9,383 \cdot 10^{-4} (\gamma - 1)$$

$$\gamma \sim 7461$$

$\gamma \gg 1$, therefore we are nearing Special Relativity

EXAMPLE

We can now verify the of the proton with that energy comes close to that of the speed of light.

$$\gamma = 1/[1 - (v/c)^2]^{1/2}$$

$$\gamma = 7461 \quad \Rightarrow \quad v = 0,99999999991 \cdot c$$

$$v \sim c$$

LORENTZ TRANSFORMATION

$$x' = \gamma (x - \beta ct)$$

$$y' = y$$

$$z' = z$$

$$ct' = \gamma (ct - \beta x)$$

$$x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}}$$

$$y' = y$$

$$z' = z$$


$$t' = \frac{t - vx/c^2}{\sqrt{1 - v^2/c^2}}$$

$$\mathbf{r}'_{\parallel} = \gamma (\mathbf{r}_{\parallel} - \beta ct)$$

$$\mathbf{r}'_{\perp} = \mathbf{r}_{\perp}$$

$$ct' = \gamma (ct - \mathbf{r}_{\parallel} \cdot \boldsymbol{\beta})$$

where \mathbf{r}_{\parallel} and \mathbf{r}_{\perp} are the components of \mathbf{r} w.r.t. $\boldsymbol{\beta}$ (or \mathbf{v})



LENGTH CONTRACTION

▶ or **Lorentz contraction**, is the solution that Lorentz proposed to solve the Michelson-Morley experiment:

is the phenomenon that a moving object's length is measured to be shorter than its proper length, which is the length as measured in the object's own rest frame

$$\Delta x' = \frac{\Delta x}{\gamma}$$


TIME DILATION

is a difference in the elapsed time measured by two clocks, due to them having a velocity relative to each other

$$\Delta t' = \gamma \Delta t$$

TIME DILATION

Reflection of light between 2 mirrors seen inside moving frame and from outside



Frame moving with velocity v

Seen from outside the path is longer, but c must be the same ..

MUON DECAY IN THE ATMOSPHERE


MUONS ARE FORMED IN COLLISIONS OF COSMIC RAYS WITH NUCLEI OF ATMOSPHERE'S ATOMS, AT HEIGHTS OF ABOUT 12000

- The half-life of a muon is 2.2 microseconds and so even moving at 0.994 c they would only expect to travel about 660 m before half of them decayed.
- As they are formed at 12000 m altitude it would take 40 μ s, or about 20 half lives, to reach the ground.. So, they almost would not reach the ground
- But they do! This means that the muons are living longer???

- Their relativistic factor is:

$$\gamma = \frac{1}{\sqrt{1 - 0.994^2}} = 9.1424$$

Their time slows down, and 2.2 μ s become about γ times longer, or Lengths contract and the 12000m become 12000/ γ m.



SOME CLARIFICATION

- **Lorentz Contraction:**
 - It is not the matter that is compressed (what Lorentz thought)
 - It is the space that is modified (Einstein)
- **Time Dilation**
 - It is not the clock that is changed (what Lorentz and others thought)
 - It is the time that is modified (Einstein)
- **EINSTEIN'S MAIN CONTRIBUTION: TO BELIEVE IT!**

PROPER DEFINITIONS

PROPER MASS:

mass of a body at rest


PROPER TIME:

time as measured
in its own frame

PROPER LENGTH:

length as measured in its own frame





**EINSTEIN'S
PRINCIPLE OF
RELATIVITY**

IN ALBERT EINSTEIN'S ORIGINAL TREATMENT, IN 1905,
THE PRINCIPLE OF RELATIVITY IS BASED ON
TWO POSTULATES:

1. SPECIAL PRINCIPLE OF RELATIVITY:

The laws of physics are invariant (i.e. identical) in all inertial frames of reference (i.e. non-accelerating frames of reference).

2. INVARIANCE OF C:

The speed of light in a vacuum is the same for all observers, regardless of the motion of the light source or observer.

1905, Albert Einstein , "On the Electrodynamics of Moving Bodies".

EINSTEIN POSTULATES CONSEQUENCES: SPECIAL PRINCIPLE OF RELATIVITY

- All physical laws (e.g. Newton's Laws or Maxwell's Equations) all have the same observed form in all inertial reference frames. This is "helpful" in that the basic laws of physics are not dependent on your state of motion.
- But as a consequence of this, it is impossible to tell from the laws of physics in your frame whether you are in motion or not.

There is no such thing as an absolute state of rest or motion - all motion is relative.

EINSTEIN POSTULATES CONSEQUENCES: INVARIANCE OF C

- All observers agree that light moves at a fixed speed - this is the singular invariant independent of states of relative motion;
- But as a consequence of this, the belief that time or space or both are experienced in the same way by observers in different states of motion must be abandoned.

There is no such thing as an absolute measure of time or space; measurements in one frame of reference need not agree with those in another, but all observers will agree that light signals travel at a fixed speed.

EINSTEIN POSTULATES CONSEQUENCES

- Space and time are NOT independent quantities
- Relativistic phenomena (with relevance for accelerators):
 - No speed of moving objects can exceed speed of light
 - (Non-) Simultaneity of events in independent frames
 - Lorentz Contraction and Time Dilation
 - Relativistic Doppler effect – change in frequency (and wavelength) of light, caused by the relative motion of the source and the observer
- There are no absolute time and space, no absolute motion

Inertial system: It is not possible to know whether one is moving or not

DEFINITIONS

| | | |
|---|-----------------------|------------|
| m | rest mass | MeV/ c^2 |
| $E_0 = mc^2$ | rest energy | MeV |
| $E = \gamma mc^2$ | total energy | MeV |
| $K = E - mc^2$ | kinetic energy | MeV |
| \mathbf{v} | velocity | m/s |
| $\boldsymbol{\beta} = \mathbf{v}/c$ | relativistic velocity | — |
| $\gamma = 1/\sqrt{1 - \boldsymbol{\beta} \cdot \boldsymbol{\beta}}$ | lorentz factor | — |
| $\mathbf{P} = \boldsymbol{\beta}\gamma mc$ | momentum | MeV/ c |
| $E^2 = (Pc)^2 + (mc^2)^2$ | total energy | MeV |

DEFINITIONS AND PRACTICAL UNITS

$$\beta = \frac{v}{c} = \frac{1}{\gamma}$$

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{1}{\sqrt{1 - \beta_r^2}}$$

β_r relativistic speed: $\beta_r = [0, 1]$

LHC: $\beta_r \approx 0.9999999991$

γ Lorentz factor: $\gamma = [1, \infty)$

LHC: $\gamma \approx 7461$

USEFUL RELATIONS AND QUANTITIES

$$E^2 = P^2 c^2 + m^2 c^4$$

total energy

MeV

$$\mathbf{P}c = E\boldsymbol{\beta}$$

total momentum times c

$$m_e = 0.510999$$

rest mass of the electron

MeV/ c^2

$$m_p = 938.272$$

rest mass of the proton

MeV/ c^2

$$m_\mu = 105.66$$

rest mass of the muon

MeV/ c^2

Frequent subdivisions

$$\gamma \simeq 1$$

non-relativistic

$$\gamma > 1$$

relativistic

$$\gamma \gg 1$$



ultra-relativistic

EXAMPLES

total
energy

$$\gamma = \frac{E}{E_0} = \frac{m}{m_0} = \frac{1}{\sqrt{1 - v^2/c^2}} = \frac{1}{\sqrt{1 - \beta^2}}$$

rest energy



$$\beta = \frac{v}{c} = \sqrt{1 - \frac{1}{\gamma^2}}$$

Normalized velocity

TRANSFORMATIONS OF ELECTROMAGNETIC FIELDS

$$\begin{cases} E_x = \gamma (E'_x + vB'_y) \\ E_y = \gamma (E'_y - vB'_x) \\ E_z = E'_z \end{cases}$$

Unprimed quantities are in the lab frame, primed quantities in the a frame moving with velocity v along the z axis

$$\begin{cases} B_x = \gamma (B'_x - vE'_y/c) \\ B_y = \gamma (B'_y + vE'_x/c) \\ B_z = B'_z \end{cases}$$

$$\mathbf{E} = \gamma (\mathbf{E}' - \mathbf{v} \times \mathbf{B}') - \frac{\gamma^2}{1 + \gamma} (\mathbf{v} \cdot \mathbf{E}') \mathbf{v}$$

$$\gamma = 1/\sqrt{1 - v^2/c^2}$$

In compact 3d vector form for a frame moving with arbitrary velocity \mathbf{v}

"For a charge moving in an electromagnetic field, the force experienced by the charge is equal to the electric force, transformed into the rest frame of the charge" A. Einstein

EVERY DAY EXAMPLE: GPS SATELLITE



- 20'000 km above ground, (unlike popular believe: not on geostationary orbits)
- Orbital speed 14'000 km/h (i.e. relative to observer on earth)
- On-board clock accuracy 50 ns
- Navigation accuracy 15 meters

Do we correct for relativistic effects ?

EVERY DAY EXAMPLE: GPS SATELLITE

Orbital speed 14000 km/h ≈ 3.9 km/s

$\rightarrow \beta \approx 1.3 \times 10^{-5}, \gamma \approx 1.00000000000084$

Small, but accumulates $7 \mu\text{s}$ during one day compared to reference time on earth!

After one day: your position wrong by ≈ 2 km !! (including general relativity error is 10 km per day)

Special relativity: $7 \mu\text{s}$ slower, general relativity: $45 \mu\text{s}$ faster

Countermeasures:

- (1) Minimum **4 satellites** (avoid reference time on earth)
- (2) Detune data transmission frequency from **1.023 MHz** to **1.022999999543 MHz** prior to launch



THE END



Thank you for your attention!



CONTACT INFORMATION

irina.shreyber@cern.ch

<https://www.linkedin.com/in/ishreyber/>

