

Measurement of the Energy Response of the ATLAS Calorimeter to Charged Pions from tau-lepton Decays in Run 2 Data

Monday, 25 September 2023 16:30 (13 minutes)

The measurement of the ATLAS calorimeter response E/p_T is performed for single charged pions with a transverse momentum (p_T) ranging from 10 to 300 GeV using 139 fb⁻¹ of proton-proton collision data at $\sqrt{s}=13$ TeV taken in RUN 2 by the ATLAS detector of the Large Hadron Collider (LHC) [1]. The use of charged pions originating from τ -lepton decays allows the measurement of the response of the calorimeter in the high p_T regime. In the ATLAS simulation, this response is found to be approximately 2% overestimated in the central region and 4% underestimated in the endcaps. The uncertainties in the measurements in the central region are less than 1% for p_T ranging from 15 to 185 GeV and reach 0.6% in the most precise region. A brief introduction about the ATLAS calorimeter will be presented, followed by a description of the event selection. The energy response as well as the uncertainties in the measurements will also be evaluated [2].

References

[1] ATLAS Collaboration, The ATLAS Experiment at the CERN Large Hadron Collider

URL: <https://cds.cern.ch/record/1129811>

[2] Measurement of the Energy Response of the ATLAS Calorimeter to Charged Pions from $W^\pm \rightarrow \tau^\pm(\rightarrow \pi^\pm \nu_\tau) \nu_\tau$ Events in Run 2 data : CERN-EP-2021-147

URL: <https://cds.cern.ch/record/2778857>

Abstract Category

Instrumentations & Detectors

Primary author: NOURY, Meryem (Universite Hassan II, Ain Chock (MA))

Presenter: NOURY, Meryem (Universite Hassan II, Ain Chock (MA))

Session Classification: Parallel Session 1

Track Classification: Physics Research