Contribution ID: 23 Type: not specified

## TMD factorisation for diffractive jets in photon-nucleus interactions

Tuesday 23 May 2023 11:50 (20 minutes)

Using the colour dipole picture and the colour glass condensate (CGC) framework, we systematically study the diffractive production of two (quark-antiquark) and three (quark, antiquark, and gluon) jets via coherent photon-nucleus interactions at high energy. We focus on the hard scattering regime where at least one of the transverse momentum scales in the problem (the photon virtuality and/or the relative transverse momentum of two of the produced jets) is much larger than the saturation momentum  $Q_s$  of the nuclear target. We argue that, despite this hardness, the cross-sections for all these processes are controlled by large dipole configurations, with transverse size  $R \sim 1/Q_s$ , which undergo strong scattering. We demonstrate that these dominant contributions admit factorisations in terms of the quark or gluon diffractive transverse-momentum dependent distributions (TMDs) of the Pomeron. These diffractive TMDs have support at transverse momenta  $k_{\perp} \leq Q_s$  and vanish as a power of 1-x when the parton splitting fraction x approaches to unity. Their dependence upon the longitudinal momentum fraction of the Pomeron  $x_{\mathbb{P}}$  is controlled by the BK/JIMWLK equations of the CGC. By integrating over  $k_{\perp}$  up to the hard scale, we deduce explicit results for the respective diffractive PDFs and structure functions.

Authors: IANCU, Edmond (Université Paris-Saclay (FR)); MUELLER, Alfred (Columbia University); TRI-

ANTAFYLLOPOULOS, Dionysios (ECT\*/FBK); Dr WEI, Shu-yi (Shandong University)

Presenter: IANCU, Edmond (Université Paris-Saclay (FR))

**Session Classification:** TMDs