

# Investigating the exclusive photoproduction of a photon meson pair as a promising channel to probe GPDs

REVESTRUCTURE workshop

Saad Nabeebaccus  
IJCLab



**Gluodynamics**

July 10, 2023

Based on 2212.00655, 2302.12026 and work in progress with S. Wallon, L. Szymanowski, B. Pire, G. Duplančić, K. Passek-Kumerički, N. Crnković

# Introduction

Quark GPDs: twist 2 Chiral-even

Quark GPDs at twist 2 [Diehl: hep-ph/0307382]

without helicity flip (chiral-even  $\Gamma$  matrices): 4 chiral-even GPDs:  
(Note:  $\Delta = p' - p$ )

$$\begin{aligned} F^q &= \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}(-\frac{1}{2}z) \gamma^+ q(\frac{1}{2}z) | p \rangle \Big|_{z^+=0, z_\perp=0} \\ &= \frac{1}{2P^+} \left[ H^q(x, \xi, t) \bar{u}(p') \gamma^+ u(p) + E^q(x, \xi, t) \bar{u}(p') \frac{i \sigma^{+\alpha} \Delta_\alpha}{2m} u(p) \right], \end{aligned}$$

# Introduction

Quark GPDs: twist 2 Chiral-even

Quark GPDs at twist 2 [Diehl: hep-ph/0307382]

without helicity flip (chiral-even  $\Gamma$  matrices): 4 chiral-even GPDs:  
(Note:  $\Delta = p' - p$ )

$$\begin{aligned} F^q &= \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}(-\frac{1}{2}z) \gamma^+ q(\frac{1}{2}z) | p \rangle \Big|_{z^+=0, z_\perp=0} \\ &= \frac{1}{2P^+} \left[ H^q(x, \xi, t) \bar{u}(p') \gamma^+ u(p) + E^q(x, \xi, t) \bar{u}(p') \frac{i \sigma^{+\alpha} \Delta_\alpha}{2m} u(p) \right], \end{aligned}$$

$$\begin{aligned} \tilde{F}^q &= \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}(-\frac{1}{2}z) \gamma^+ \gamma_5 q(\frac{1}{2}z) | p \rangle \Big|_{z^+=0, z_\perp=0} \\ &= \frac{1}{2P^+} \left[ \tilde{H}^q(x, \xi, t) \bar{u}(p') \gamma^+ \gamma_5 u(p) + \tilde{E}^q(x, \xi, t) \bar{u}(p') \frac{\gamma_5 \Delta^+}{2m} u(p) \right]. \end{aligned}$$

# Introduction

Quark GPDs: twist 2 Chiral-even

Quark GPDs at twist 2 [Diehl: hep-ph/0307382]

without helicity flip (chiral-even  $\Gamma$  matrices): 4 chiral-even GPDs:  
(Note:  $\Delta = p' - p$ )

$$\begin{aligned} F^q &= \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}(-\frac{1}{2}z) \gamma^+ q(\frac{1}{2}z) | p \rangle \Big|_{z^+=0, z_\perp=0} \\ &= \frac{1}{2P^+} \left[ H^q(x, \xi, t) \bar{u}(p') \gamma^+ u(p) + E^q(x, \xi, t) \bar{u}(p') \frac{i\sigma^{+\alpha} \Delta_\alpha}{2m} u(p) \right], \end{aligned}$$

$$\begin{aligned} \tilde{F}^q &= \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}(-\frac{1}{2}z) \gamma^+ \gamma_5 q(\frac{1}{2}z) | p \rangle \Big|_{z^+=0, z_\perp=0} \\ &= \frac{1}{2P^+} \left[ \tilde{H}^q(x, \xi, t) \bar{u}(p') \gamma^+ \gamma_5 u(p) + \tilde{E}^q(x, \xi, t) \bar{u}(p') \frac{\gamma_5 \Delta^+}{2m} u(p) \right]. \end{aligned}$$

$$H^q \xrightarrow{\xi=0, t=0} \text{PDF } q \qquad \tilde{H}^q \xrightarrow{\xi=0, t=0} \text{polarised PDF } \Delta q$$

with helicity flip (chiral-odd  $\Gamma$  matrices): 4 chiral-odd GPDs:

$$\begin{aligned} & \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}(-\frac{1}{2}z) i\sigma^{+i} q(\frac{1}{2}z) | p \rangle \Big|_{z^+=0, z_\perp=0} \\ &= \frac{1}{2P^+} \bar{u}(p') \left[ H_T^q i\sigma^{+i} + \tilde{H}_T^q \frac{P^+ \Delta^i - \Delta^+ P^i}{m^2} \right. \\ & \quad \left. + E_T^q \frac{\gamma^+ \Delta^i - \Delta^+ \gamma^i}{2m} + \tilde{E}_T^q \frac{\gamma^+ P^i - P^+ \gamma^i}{m} \right] u(p), \end{aligned}$$

# Introduction

## Quark GPDs: twist 2 Chiral-odd

with helicity flip (chiral-odd  $\Gamma$  matrices): 4 chiral-odd GPDs:

$$\begin{aligned} & \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}(-\frac{1}{2}z) i\sigma^{+i} q(\frac{1}{2}z) | p \rangle \Big|_{z^+=0, z_\perp=0} \\ &= \frac{1}{2P^+} \bar{u}(p') \left[ H_T^q i\sigma^{+i} + \tilde{H}_T^q \frac{P^+\Delta^i - \Delta^+P^i}{m^2} \right. \\ & \quad \left. + E_T^q \frac{\gamma^+\Delta^i - \Delta^+\gamma^i}{2m} + \tilde{E}_T^q \frac{\gamma^+P^i - P^+\gamma^i}{m} \right] u(p), \end{aligned}$$

$H_T^q \xrightarrow{\xi=0, t=0}$  quark transversity PDFs  $\delta q$

**Note:**  $\tilde{E}_T^q(x, -\xi, t) = -\tilde{E}_T^q(x, \xi, t)$

# Why consider a gamma-meson pair?

Understanding quark transversity

- ▶ Transverse spin content of the proton:

$$\begin{array}{lcl} |\uparrow\rangle_{(x)} & \sim & |\rightarrow\rangle + |\leftarrow\rangle \\ |\downarrow\rangle_{(x)} & \sim & |\rightarrow\rangle - |\leftarrow\rangle \\ \text{spin along } x & & \text{helicity states} \end{array}$$

- ▶ Observables which are sensitive to helicity flip thus give access to transversity PDFs. Poorly known.

# Why consider a gamma-meson pair?

## Understanding quark transversity

- ▶ Transverse spin content of the proton:

$$\begin{array}{l} |\uparrow\rangle(x) \quad \sim \quad |\rightarrow\rangle + |\leftarrow\rangle \\ |\downarrow\rangle(x) \quad \sim \quad |\rightarrow\rangle - |\leftarrow\rangle \\ \text{spin along } x \quad \quad \quad \text{helicity states} \end{array}$$

- ▶ Observables which are sensitive to helicity flip thus give access to transversity PDFs. Poorly known.
- ▶ Transversity GPDs are completely unknown experimentally.



# Why consider a gamma-meson pair?

## Understanding quark transversity

- ▶ Transverse spin content of the proton:

$$\begin{array}{l} |\uparrow\rangle(x) \quad \sim \quad |\rightarrow\rangle + |\leftarrow\rangle \\ |\downarrow\rangle(x) \quad \sim \quad |\rightarrow\rangle - |\leftarrow\rangle \\ \text{spin along } x \quad \quad \quad \text{helicity states} \end{array}$$

- ▶ Observables which are sensitive to helicity flip thus give access to transversity PDFs. Poorly known.
- ▶ Transversity GPDs are completely unknown experimentally.
- ▶ For massless (anti)particles, chirality = (-)helicity
- ▶ Transversity GPDs can thus be accessed through **chiral-odd  $\Gamma$**  matrices.

# Why consider a gamma-meson pair?

## Understanding quark transversity

- ▶ Transverse spin content of the proton:

$$\begin{array}{l} |\uparrow\rangle(x) \quad \sim \quad |\rightarrow\rangle + |\leftarrow\rangle \\ |\downarrow\rangle(x) \quad \sim \quad |\rightarrow\rangle - |\leftarrow\rangle \\ \text{spin along } x \quad \quad \quad \text{helicity states} \end{array}$$

- ▶ Observables which are sensitive to helicity flip thus give access to transversity PDFs. Poorly known.
- ▶ Transversity GPDs are completely unknown experimentally.
- ▶ For massless (anti)particles, chirality = (-)helicity
- ▶ Transversity GPDs can thus be accessed through **chiral-odd**  $\Gamma$  matrices.
- ▶ Since (in the massless limit) QCD and QED are chiral-even ( $\gamma^\mu, \gamma^\mu\gamma^5$ ), **the chiral-odd quantities** ( $1, \gamma^5, [\gamma^\mu, \gamma^\nu]$ ) **which one wants to measure should appear in pairs.**

# Why consider a gamma-meson pair?

Can we probe quark transversity GPDs in DVMP?

- ▶ the leading DA (twist 2) of  $\rho_T$  is **chiral-odd** ( $\sigma^{\mu\nu}$  coupling)

# Why consider a gamma-meson pair?


Can we probe quark transversity GPDs in DVMP?

- ▶ the leading DA (twist 2) of  $\rho_T$  is **chiral-odd** ( $\sigma^{\mu\nu}$  coupling)
- ▶ unfortunately  $\gamma^* N \rightarrow \rho_T N' = 0$ , since such a process would require a helicity transfer of 2 from a photon. [Diehl, Gousset, Pire: hep-ph/9808479], [Collins, Diehl: hep-ph/9907498]

# Why consider a gamma-meson pair?

Can we probe quark transversity GPDs in DVMP?

- ▶ the leading DA (twist 2) of  $\rho_T$  is **chiral-odd** ( $\sigma^{\mu\nu}$  coupling)
- ▶ unfortunately  $\gamma^* N \rightarrow \rho_T N' = 0$ , since such a process would require a helicity transfer of 2 from a photon. [Diehl, Gousset, Pire: hep-ph/9808479], [Collins, Diehl: hep-ph/9907498]
- ▶ lowest order diagrammatic argument:



$$\gamma^\alpha [\gamma^\mu, \gamma^\nu] \gamma_\alpha = 0$$

# Why consider a gamma-meson pair?

Go to higher twist?

- ▶ This vanishing only occurs at [twist 2](#)
- ▶ At twist 3 this process does not vanish [[Ahmad, Goldstein, Liuti: 0805.3568](#)], [[Goloskokov, Kroll: 1106.4897, 1310.1472](#)]

# Why consider a gamma-meson pair?

Go to higher twist?

- ▶ This vanishing only occurs at **twist 2**
- ▶ At twist 3 this process does not vanish [Ahmad, Goldstein, Liuti: 0805.3568], [Goloskokov, Kroll: 1106.4897, 1310.1472]
- ▶ However processes involving **twist 3 DAs** may face problems with factorisation (end-point singularities)

⇒ can be made safe in the high-energy  $k_T$ -factorisation approach

[Anikin, Ivanov, Pire, Szymanowski, Wallon: 0909.4090]

# Why consider a gamma-meson pair?

A convenient alternative solution

Circumvent this using 3-body final states:

▶  $\gamma N \rightarrow MMN'$ :

El Beiyad, Enberg, Ivanov, Pire, Segond, Szymanowski, Teryaev, Wallon:  
[hep-ph/0209300, hep-ph/0601138, 1001.4491]

▶  $\gamma N \rightarrow \gamma MN'$ :

Boussarie, Duplančić, Nabeebaccus, Passek-Kumerički Pire, Szymanowski,  
Wallon: [1609.03830, 1809.08104, 2212.00655, 2302.12026]



# Why consider a gamma-meson pair?

A convenient alternative solution

Circumvent this using 3-body final states:

▶  $\gamma N \rightarrow MMN'$ :

El Beiyad, Enberg, Ivanov, Pire, Segond, Szymanowski, Teryaev, Wallon:  
[hep-ph/0209300, hep-ph/0601138, 1001.4491]

▶  $\gamma N \rightarrow \gamma MN'$ :

Boussarie, Duplančić, Nabeebaccus, Passek-Kumerički Pire, Szymanowski,  
Wallon: [1609.03830, 1809.08104, 2212.00655, 2302.12026]

Also many others that are not sensitive to chiral-odd GPDs:

▶  $\gamma N \rightarrow \gamma\gamma N'$ : *See Jakub's talk!*

Grocholski, Pedrak, Pire, Sznajder, Szymanowski, Wagner: [1708.01043,  
2003.03263, 2110.00048, 2204.00396]

▶  $\pi N \rightarrow \gamma\gamma N'$ :

Qiu, Yu: [2205.07846]

# Why consider a gamma-meson pair?

A convenient alternative solution

Circumvent this using 3-body final states:

▶  $\gamma N \rightarrow MMN'$ :

El Beiyad, Enberg, Ivanov, Pire, Segond, Szymanowski, Teryaev, Wallon:  
[hep-ph/0209300, hep-ph/0601138, 1001.4491]

▶  $\gamma N \rightarrow \gamma MN'$ :

Boussarie, Duplančić, Nabeebaccus, Passek-Kumerički Pire, Szymanowski,  
Wallon: [1609.03830, 1809.08104, 2212.00655, 2302.12026]

Also many others that are not sensitive to chiral-odd GPDs:

▶  $\gamma N \rightarrow \gamma\gamma N'$ : *See Jakub's talk!*

Grocholski, Pedrak, Pire, Sznajder, Szymanowski, Wagner: [1708.01043,  
2003.03263, 2110.00048, 2204.00396]

▶  $\pi N \rightarrow \gamma\gamma N'$ :


Qiu, Yu: [2205.07846]

In all of the above cases, the richer kinematics of the process allows the sensitivity of GPDs wrt  $x$  to be probed (unlike moment-type dependence, e.g. in DVCS) Qiu, Yu: [2305.15397]

# Why consider a gamma-meson pair?

A convenient alternative solution

- ▶ Consider the process  $\gamma N \rightarrow \gamma MN'$ ,  $M$  = meson. Collinear factorisation of the amplitude at large  $M_{\gamma M}^2$ ,  $t'$ , and small  $t$ .




large angle factorisation  
à la Brodsky Lepage

# Why consider a gamma-meson pair?

A convenient alternative solution

- ▶ Consider the process  $\gamma N \rightarrow \gamma M N'$ ,  $M$  = meson. Collinear factorisation of the amplitude at large  $M_{\gamma M}^2$ ,  $t'$ , and small  $t$ .




large angle factorisation  
à la Brodsky Lepage

- ▶ Mesons considered in the final state:  $\pi^\pm$ ,  $\rho_{L,T}^\pm, 0$ .

# Why consider a gamma-meson pair?

A convenient alternative solution

- ▶ Consider the process  $\gamma N \rightarrow \gamma M N'$ ,  $M$  = meson. Collinear factorisation of the amplitude at large  $M_{\gamma M}^2$ ,  $t'$ , and small  $t$ .



large angle factorisation


à la Brodsky Lepage

- ▶ Mesons considered in the final state:  $\pi^\pm$ ,  $\rho_{L,T}^{\pm,0}$ .
- ▶ Leading order and leading twist

# Why consider a gamma-meson pair?

Chiral-odd GPDs using  $\rho_T \gamma$  production

How does it work (at LO)?



Typical non-zero diagram for a **transverse**  $\rho$  meson

the  $\sigma$  matrices (from either the DA or the GPD) do not kill it anymore!

# Is QCD factorisation really justified?

- ▶ Recently, factorisation has been proved for the process  $\pi^\pm N \rightarrow \gamma\gamma N'$  by Qiu, Yu [2205.07846].
- ▶ This was extended to a wide range of  $2 \rightarrow 3$  exclusive processes by Qiu, Yu [2210.07995]

# Is QCD factorisation really justified?


- ▶ Recently, factorisation has been proved for the process  $\pi^\pm N \rightarrow \gamma\gamma N'$  by Qiu, Yu [2205.07846].
- ▶ This was extended to a wide range of  $2 \rightarrow 3$  exclusive processes by Qiu, Yu [2210.07995]
- ▶ The proof relies on having **large  $p_T$** , rather than large invariant mass (e.g. photon-meson pair).



# Is QCD factorisation really justified?

- ▶ Recently, factorisation has been proved for the process  $\pi^\pm N \rightarrow \gamma\gamma N'$  by Qiu, Yu [2205.07846].
- ▶ This was extended to a wide range of  $2 \rightarrow 3$  exclusive processes by Qiu, Yu [2210.07995]
- ▶ The proof relies on having large  $p_T$ , rather than large invariant mass (e.g. photon-meson pair).
- ▶ In fact, NLO computation has been performed for  $\gamma N \rightarrow \gamma\gamma N'$  by Grocholski, Pire, Sznajder, Szymanowski, Wagner [2110.00048]: See Jakub's talk!
- ▶ Also, NLO computation for  $\gamma\gamma \rightarrow \pi^+\pi^-$  by crossing symmetry Duplancic, Nizic: [hep-ph/0607069].

$$\gamma(q) + N(p_1) \rightarrow \gamma(k) + M(p_M, \varepsilon_M) + N'(p_2)$$



Useful Mandelstam variables:

$$t = (p_2 - p_1)^2$$

$$u' = (p_M - q)^2$$


$$t' = (k - q)^2$$

► Factorisation requires:

$$-u' > 1 \text{ GeV}^2, \quad -t' > 1 \text{ GeV}^2 \quad \text{and} \quad (-t)_{\min} \leq -t \leq .5 \text{ GeV}^2$$

⇒ sufficient to ensure **large**  $p_T$ .

$$\gamma(q) + N(p_1) \rightarrow \gamma(k) + M(p_M, \varepsilon_M) + N'(p_2)$$



Useful Mandelstam variables:

$$t = (p_2 - p_1)^2$$

$$u' = (p_M - q)^2$$

$$t' = (k - q)^2$$

- ▶ Factorisation requires:

$$-u' > 1 \text{ GeV}^2, \quad -t' > 1 \text{ GeV}^2 \quad \text{and} \quad (-t)_{\min} \leq -t \leq .5 \text{ GeV}^2$$

$\implies$  sufficient to ensure **large**  $p_{T\gamma}$ .


- ▶ Cross-section differential in  $(-u')$  and  $M^2_{\gamma M}$ , and evaluated at  $(-t) = (-t)_{\min}$ .

$$\mathcal{A} = \int_{-1}^1 dx \int_0^1 dz T(x, \xi, z) H(x, \xi, t) \Phi_M(z)$$

$$\mathcal{A} = \int_{-1}^1 dx \int_0^1 dz T(x, \xi, z) H(x, \xi, t) \Phi_M(z)$$

► Differential cross section:

$$\left. \frac{d\sigma}{dt du' dM_{\gamma M}^2} \right|_{-t=(-t)_{min}} = \frac{|\overline{\mathcal{A}}|^2}{32S_{\gamma N}^2 M_{\gamma M}^2 (2\pi)^3}.$$



$$\mathcal{A} = \int_{-1}^1 dx \int_0^1 dz T(x, \xi, z) H(x, \xi, t) \Phi_M(z)$$


- Differential cross section:

$$\left. \frac{d\sigma}{dt du' dM_{\gamma M}^2} \right|_{-t=(-t)_{min}} = \frac{|\overline{\mathcal{A}}|^2}{32 S_{\gamma N}^2 M_{\gamma M}^2 (2\pi)^3}.$$

- Kinematic parameters:  $S_{\gamma N}$ ,  $M_{\gamma M}^2$ ,  $-t$ ,  $-u'$
- Useful dimensionless variables (hard part):

$$\alpha = \frac{-u'}{M_{\gamma M}^2},$$

$$\xi = \frac{M_{\gamma M}^2}{2(S_{\gamma N} - m_N^2) - M_{\gamma M}^2}.$$



# Computation

Parametrising the GPDs: 2 scenarios for polarised and transversity PDFs

Quark GPDs are parametrised in terms of **Double Distributions**  
[Radyushkin: [hep-ph/9805342](https://arxiv.org/abs/hep-ph/9805342)]

Quark GPDs are parametrised in terms of **Double Distributions**  
[Radyushkin: [hep-ph/9805342](https://arxiv.org/abs/hep-ph/9805342)]

For **polarised** PDFs (and hence **transversity** PDFs), two scenarios are proposed for the parameterization:

- ▶ “**standard**” scenario, with flavor-symmetric light sea quark and antiquark distributions.
- ▶ “**valence**” scenario with a completely flavor-asymmetric light sea quark densities.



- ▶ We take the simplistic **asymptotic** form of the DAs

$$\phi_{\text{as}}(z) = 6z(1 - z).$$

- ▶ We take the simplistic **asymptotic** form of the DAs

$$\phi_{\text{as}}(z) = 6z(1 - z).$$

- ▶ We also investigate the effect of using a **holographic** DA:

$$\phi_{\text{hol}}(z) = \frac{8}{\pi} \sqrt{z(1 - z)}.$$

Suggested by

- ▶ AdS/QCD correspondence [[Brodsky, de Teramond: hep-ph/0602252](#)],
- ▶ dynamical chiral symmetry breaking on the light-front [[Shi, Chen, Chang, Roberts, Schmidt, Zong: 1504.00689](#)],
- ▶ recent lattice results. [[Gao, Hanlon, Karthik, Mukherjee, Petreczky, Scior, Syritsyn, Zhao: 2206.04084](#)]

# Exclusive photoproduction of $\pi^0\gamma$

## Gluonic GPD contributions

- ▶ Because of the quantum numbers of  $\pi^0$  ( $J^{PC} = 0^{-+}$ ), the exclusive photoproduction of  $\pi^0\gamma$  is also sensitive to gluonic GPD contributions.

# Exclusive photoproduction of $\pi^0\gamma$



## Gluonic GPD contributions

- ▶ Because of the quantum numbers of  $\pi^0$  ( $J^{PC} = 0^{-+}$ ), the exclusive photoproduction of  $\pi^0\gamma$  is also sensitive to gluonic GPD contributions.
- ▶ A total of 24 diagrams contribute in this case (compared to 20 diagrams from quark GPD contributions), with 6 groups of 4 related by symmetries ( $x \rightarrow -x$  and  $z \rightarrow 1 - z$  separately).

# Exclusive photoproduction of $\pi^0\gamma$


## Gluonic GPD contributions

- ▶ Because of the quantum numbers of  $\pi^0$  ( $J^{PC} = 0^{-+}$ ), the exclusive photoproduction of  $\pi^0\gamma$  is also sensitive to gluonic GPD contributions.
- ▶ A total of 24 diagrams contribute in this case (compared to 20 diagrams from quark GPD contributions), with 6 groups of 4 related by symmetries ( $x \rightarrow -x$  and  $z \rightarrow 1 - z$  separately).
- ▶ Diagrams amount to connecting photons to the following two topologies.



# Exclusive photoproduction of $\pi^0\gamma$

Gluonic GPD contributions



$$D_a = ((x - \xi)p + \bar{z}p_M)^2 + i\epsilon$$

$$= s\bar{\alpha}\bar{z} \left[ x - \xi + i\epsilon \right],$$

$$D_b = (k + zp_M - (x + \xi)p)^2 + i\epsilon$$


$$= -s \left[ z \left( x - \xi - i\epsilon \right) + \alpha\bar{z} \left( x + \xi - i\epsilon \right) \right],$$

$$D_c = (k + zp_M)^2 + i\epsilon$$

$$= 2s\xi z + i\epsilon$$

# Exclusive photoproduction of $\pi^0\gamma$

Gluonic GPD contributions



$$D_a = ((x - \xi)p + \bar{z}p_M)^2 + i\epsilon$$

$$= s\bar{\alpha}\bar{z} \left[ x - \xi + i\epsilon \right],$$

$$D_b = (k + zp_M - (x + \xi)p)^2 + i\epsilon$$

$$= -s \left[ z \left( x - \xi - i\epsilon \right) + \alpha\bar{z} (x + \xi - i\epsilon) \right],$$


$$D_c = (k + zp_M)^2 + i\epsilon$$

$$= 2s\xi z + i\epsilon$$

$\implies$  pinching of poles in the propagators in the limit of  $z \rightarrow 1$

# Exclusive photoproduction of $\pi^0\gamma$

Gluonic GPD contributions



$$D_a = ((x - \xi)p + \bar{z}p_M)^2 + i\epsilon$$

$$= s\bar{\alpha}\bar{z} \left[ \boxed{x - \xi + i\epsilon} \right],$$

$$D_b = (k + zp_M - (x + \xi)p)^2 + i\epsilon$$

$$= -s \left[ z \left( \boxed{x - \xi - i\epsilon} \right) + \alpha\bar{z}(x + \xi - i\epsilon) \right],$$

$$D_c = (k + zp_M)^2 + i\epsilon$$

$$= 2s\xi z + i\epsilon$$

$\implies$  *pinching of poles in the propagators in the limit of  $z \rightarrow 1$*


Assuming an asymptotic form of the DA, they manifest themselves *as a purely imaginary part*, in terms of

- ▶  $\int_0^1 \frac{dz}{z\bar{z}}$  contributions, when the  $x$ -integration is performed first,
- ▶  $\int_1^1 dx \frac{\ln(x-\xi-i\epsilon)}{(x-\xi+i\epsilon)}$  contributions, when the  $z$ -integration is performed first.



# Exclusive photoproduction of $\pi^0\gamma$


Glauonic GPD contributions: Singularity structure of the full amplitude



- Unfortunately, no cancellations between the 4 corners.

# Exclusive photoproduction of $\pi^0\gamma$


Gluonic GPD contributions: Singularity structure of the full amplitude



- ▶ Unfortunately, no cancellations between the 4 corners.
- ▶ Problem also shows up in  $\pi^0 N \rightarrow \gamma\gamma N$ .

# Exclusive photoproduction of $\pi^0\gamma$


Gluonic GPD contributions: Singularity structure of the full amplitude



- ▶ Unfortunately, no cancellations between the 4 corners.
- ▶ Problem also shows up in  $\pi^0 N \rightarrow \gamma\gamma N$ .
- ▶ In  $\gamma\gamma \rightarrow \pi^+\pi^-$ , only ERBL region exists, no poles are crossed, and endpoint contributions are suppressed by DAs.

# Exclusive photoproduction of $\pi^0\gamma$

Gluonic GPD contributions: Singularity structure of the full amplitude




- ▶ Unfortunately, no cancellations between the 4 corners.
- ▶ Problem also shows up in  $\pi^0 N \rightarrow \gamma\gamma N$ .
- ▶ In  $\gamma\gamma \rightarrow \pi^+\pi^-$ , only ERBL region exists, no poles are crossed, and endpoint contributions are suppressed by DAs.
- ▶ Problem with factorisation?  
At twist-2??


# Results

Single differential cross-section:  $\gamma\rho_p^+$  vs  $\gamma\pi_p^+$

$$\frac{d\sigma_{\gamma\rho_p^+}^{\text{even}}}{dM_{\gamma\rho_p^+}^2} \text{ (pb} \cdot \text{GeV}^{-2}\text{)}$$



$$\frac{d\sigma_{\gamma\pi_p^+}^{\text{even}}}{dM_{\gamma\pi_p^+}^2} \text{ (pb} \cdot \text{GeV}^{-2}\text{)}$$



$$S_{\gamma N} = 8, 14, 20 \text{ GeV}^2$$


Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario


# Results

Single differential cross-section:  $\gamma\rho_p^+$  vs  $\gamma\pi_p^+$

$$\frac{d\sigma^{\text{even}}}{dM^2} \frac{\gamma\rho_p^+}{\gamma\rho_p^+} \text{ (pb} \cdot \text{GeV}^{-2}\text{)}$$



$$\frac{d\sigma^{\text{even}}}{dM^2} \frac{\gamma\pi_p^+}{\gamma\pi_p^+} \text{ (pb} \cdot \text{GeV}^{-2}\text{)}$$



$$S_{\gamma N} = 8, 14, 20 \text{ GeV}^2$$

Dashed: Holographic DA      non-dashed: Asymptotical DA


Dotted: standard scenario      non-dotted: valence scenario

⇒ Effect of GPD model more important on  $\pi_p^+$  than on  $\rho_p^+$


# Results

Single differential cross-section:  $\gamma\rho_{pL}^+$  vs  $\gamma\rho_{pT}^+$

$$\frac{d\sigma^{\text{even}}}{dM^2} \frac{\gamma\rho_{pL}^+}{\gamma\rho_p^+} \text{ (pb} \cdot \text{GeV}^{-2}\text{)}$$



$$\frac{d\sigma^{\text{odd}}}{dM^2} \frac{\gamma\rho_{pT}^+}{\gamma\rho_p^+} \text{ (pb} \cdot \text{GeV}^{-2}\text{)}$$





$$S_{\gamma N} = 80, 140, 200 \text{ GeV}^2$$

Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario

# Results

Single differential cross-section:  $\gamma\rho_{P,L}^+$  vs  $\gamma\rho_{P,T}^+$



$$S_{\gamma N} = 80, 140, 200 \text{ GeV}^2$$

Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario


⇒ CO cross-section is suppressed by a factor of  $\xi^2$  ( $\xi \approx \frac{M^2_{\gamma\rho}}{2S_{\gamma N}}$ ):  
Measurable at small  $S_{\gamma N}$ , but drops rapidly with increasing  $S_{\gamma N}$ .




# Results

Integrated cross-section:  $\gamma\pi_p^+$  vs  $\gamma\pi_n^-$

$\sigma_{\gamma\pi_p^+}^{even}$  (pb)



$\sigma_{\gamma\pi_n^-}^{even}$  (pb)



Dashed: Holographic DA

non-dashed: Asymptotical DA


Dotted: standard scenario

non-dotted: valence scenario


# Results

Integrated cross-section:  $\gamma\pi_p^+$  vs  $\gamma\pi_n^-$

$\sigma_{\gamma\pi_p^+}^{even}$  (pb)



$\sigma_{\gamma\pi_n^-}^{even}$  (pb)





Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario

⇒ Huge effect from GPD model in  $\pi_p^+$  case.

# Results

Integrated cross-section:  $\gamma\rho_{pL}^+$  vs  $\gamma\rho_{pT}^+$



Dashed: Holographic DA



non-dashed: Asymptotical DA

Dotted: standard scenario

non-dotted: valence scenario

# Results

Integrated cross-section:  $\gamma\rho_{pL}^+$  vs  $\gamma\rho_{pT}^+$



Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario

$\implies \xi^2$  suppression in the chiral-odd case causes the cross-section to drop rapidly with  $S_{\gamma N}$ .

# Results

## Polarisation Asymmetries wrt incoming photon

We consider an **unpolarised target**, and determine polarisation asymmetries wrt the incoming photon.

We consider an **unpolarised target**, and determine polarisation asymmetries wrt the incoming photon.

- ▶ Circular polarisation asymmetry = 0.
- ▶ Linear polarisation asymmetry,  $LPA = \frac{d\sigma_x - d\sigma_y}{d\sigma_x + d\sigma_y}$ , where  $x$  is the direction defined by  $p_{\perp}$  (direction of outgoing photon in the transverse plane).

We consider an **unpolarised target**, and determine polarisation asymmetries wrt the incoming photon.

- ▶ Circular polarisation asymmetry = 0.
- ▶ Linear polarisation asymmetry,  $LPA = \frac{d\sigma_x - d\sigma_y}{d\sigma_x + d\sigma_y}$ , where  $x$  is the direction defined by  $p_{\perp}$  (direction of outgoing photon in the transverse plane).

- ▶ In fact,

$$LPA_{\text{Lab}} = LPA \cos(2\theta) ,$$

where  $\theta$  is the angle between the lab frame  $x$ -direction and  $p_{\perp}$ .

We consider an **unpolarised target**, and determine polarisation asymmetries wrt the incoming photon.

- ▶ Circular polarisation asymmetry = 0.
- ▶ Linear polarisation asymmetry,  $LPA = \frac{d\sigma_x - d\sigma_y}{d\sigma_x + d\sigma_y}$ , where  $x$  is the direction defined by  $p_{\perp}$  (direction of outgoing photon in the transverse plane).

- ▶ In fact,

$$LPA_{\text{Lab}} = LPA \cos(2\theta),$$



where  $\theta$  is the angle between the lab frame  $x$ -direction and  $p_{\perp}$ .

- ▶ **Kleiss-Sterling** spinor techniques used to obtain expressions.
- ▶ **Both asymmetries zero in chiral-odd case!**



# Results

LPA wrt incoming photon: Single-differential level:  $\gamma\rho_p^+$  vs  $\gamma\pi_p^+$





$$S_{\gamma N} = 8, 14, 20 \text{ GeV}^2$$

Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario

# Results

LPA wrt incoming photon: Single-differential level:  $\gamma\rho_p^+$  vs  $\gamma\pi_p^+$



$$S_{\gamma N} = 8, 14, 20 \text{ GeV}^2$$



Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario

⇒ GPD model changes the behaviour of the LPA completely in the  $\pi_p^+$  case!

# Results

LPA wrt incoming photon: Integrated level:  $\gamma\rho_n^-$  vs  $\gamma\pi_n^-$



Dashed: Holographic DA



non-dashed: Asymptotical DA

Dotted: standard scenario

non-dotted: valence scenario

# Results

LPA wrt incoming photon: Integrated level:  $\gamma\rho_n^-$  vs  $\gamma\pi_n^-$



Dashed: Holographic DA

non-dashed: Asymptotical DA

Dotted: standard scenario

non-dotted: valence scenario

$\Rightarrow$  LPAs are sizeable!

# Prospects at experiments

Counting rates: JLab

Good statistics: For example, at [JLab Hall B](#):

- ▶ untagged incoming  $\gamma \Rightarrow$  **Weizsäcker-Williams** distribution

Good statistics: For example, at [JLab Hall B](#):

- ▶ untagged incoming  $\gamma \Rightarrow$  **Weizsäcker-Williams** distribution
- ▶ with an expected luminosity of  $\mathcal{L} = 100 \text{ nb}^{-1}\text{s}^{-1}$ , for 100 days of run:
  - $\rho_L^0$  (on  $p$ ) :  $\approx 2.4 \times 10^5$
  - $\rho_T^0$  (on  $p$ ) :  $\approx 4.2 \times 10^4$  (Chiral-odd)
  - $\rho_L^+$  :  $\approx 1.4 \times 10^5$
  - $\rho_T^+$  :  $\approx 6.7 \times 10^4$  (Chiral-odd)
  - $\pi^+$  :  $\approx 1.8 \times 10^5$

Good statistics: For example, at [JLab Hall B](#):

- ▶ untagged incoming  $\gamma \Rightarrow$  **Weizsäcker-Williams** distribution
- ▶ with an expected luminosity of  $\mathcal{L} = 100 \text{ nb}^{-1}\text{s}^{-1}$ , for 100 days of run:
  - $\rho_L^0$  (on  $p$ ) :  $\approx 2.4 \times 10^5$
  - $\rho_T^0$  (on  $p$ ) :  $\approx 4.2 \times 10^4$  (Chiral-odd)
  - $\rho_L^+$  :  $\approx 1.4 \times 10^5$
  - $\rho_T^+$  :  $\approx 6.7 \times 10^4$  (Chiral-odd)
  - $\pi^+$  :  $\approx 1.8 \times 10^5$
- ▶ No problem in detecting outgoing photon at JLab.

At COMPASS:

- ▶ Taking a luminosity of  $\mathcal{L} = 0.1 \text{ nb}^{-1}\text{s}^{-1}$ , and 300 days of run,
  - $\rho_L^0$  (on  $p$ ) :  $\approx 1.2 \times 10^3$
  - $\rho_T^0$  (on  $p$ ) :  $\approx 1.5 \times 10^2$  (Chiral-odd)
  - $\rho_L^+$  :  $\approx 7.4 \times 10^2$
  - $\rho_T^+$  :  $\approx 2.6 \times 10^2$  (Chiral-odd)
  - $\pi^+$  :  $\approx 7.4 \times 10^2$



At COMPASS:

- ▶ Taking a luminosity of  $\mathcal{L} = 0.1 \text{ nb}^{-1}\text{s}^{-1}$ , and 300 days of run,
  - $\rho_L^0$  (on  $p$ ) :  $\approx 1.2 \times 10^3$
  - $\rho_T^0$  (on  $p$ ) :  $\approx 1.5 \times 10^2$  (Chiral-odd)
  - $\rho_L^+$  :  $\approx 7.4 \times 10^2$
  - $\rho_T^+$  :  $\approx 2.6 \times 10^2$  (Chiral-odd)
  - $\pi^+$  :  $\approx 7.4 \times 10^2$
- ▶ Lower numbers due to low luminosity (factor of  $10^3$  less than JLab!)

# Prospects at experiments

Counting rates: EIC

- ▶ At the future EIC, with an expected integrated luminosity of  $10 \text{ fb}^{-1}$  (about 100 times smaller than JLab):
  - $\rho_L^0$  (on  $\rho$ ) :  $\approx 2.4 \times 10^4$
  - $\rho_T^0$  (on  $\rho$ ) :  $\approx 2.4 \times 10^3$  (Chiral-odd)
  - $\rho_L^+$  :  $\approx 1.5 \times 10^4$
  - $\rho_T^+$  :  $\approx 4.2 \times 10^3$  (Chiral-odd)
  - $\pi^+$  :  $\approx 1.3 \times 10^4$

# Prospects at experiments

Counting rates: EIC

- ▶ At the future EIC, with an expected integrated luminosity of  $10 \text{ fb}^{-1}$  (about 100 times smaller than JLab):
  - $\rho_L^0$  (on  $p$ ) :  $\approx 2.4 \times 10^4$
  - $\rho_T^0$  (on  $p$ ) :  $\approx 2.4 \times 10^3$  (Chiral-odd)
  - $\rho_L^+$  :  $\approx 1.5 \times 10^4$
  - $\rho_T^+$  :  $\approx 4.2 \times 10^3$  (Chiral-odd)
  - $\pi^+$  :  $\approx 1.3 \times 10^4$
- ▶ **Small  $\xi$  study:**  
 $300 < S_{\gamma N} / \text{GeV}^2 < 20000$  ( $5 \cdot 10^{-5} < \xi < 5 \cdot 10^{-3}$ ):
  - $\rho_L^0$  (on  $p$ ) :  $\approx 1.2 \times 10^3$
  - $\rho_T^0$  (on  $p$ ) :  $\approx 6.5$  (Chiral-odd) (tiny)
  - $\rho_L^+$  :  $\approx 9.3 \times 10^2$
  - $\pi^+$  :  $\approx 5.0 \times 10^2$

For p-Pb UPCs at LHC (integrated luminosity of  $1200 \text{ nb}^{-1}$ ):

► With future data from runs 3 and 4,


- $\rho_L^0 : \approx 1.6 \times 10^4$
- $\rho_T^0 : \approx 1.7 \times 10^3$  (Chiral-odd)
- $\rho_L^+ : \approx 1.1 \times 10^4$
- $\rho_T^+ : \approx 2.9 \times 10^3$  (Chiral-odd)
- $\pi^+ : \approx 9.3 \times 10^3$

For p-Pb UPCs at LHC (integrated luminosity of  $1200 \text{ nb}^{-1}$ ):

- ▶ With future data from runs 3 and 4,
  - $\rho_L^0 : \approx 1.6 \times 10^4$
  - $\rho_T^0 : \approx 1.7 \times 10^3$  (Chiral-odd)
  - $\rho_L^+ : \approx 1.1 \times 10^4$
  - $\rho_T^+ : \approx 2.9 \times 10^3$  (Chiral-odd)
  - $\pi^+ : \approx 9.3 \times 10^3$
- ▶  $300 < S_{\gamma N} / \text{GeV}^2 < 20000$  ( $5 \cdot 10^{-5} < \xi < 5 \cdot 10^{-3}$ ):
  - $\rho_L^0 : \approx 8.1 \times 10^2$
  - $\rho_L^+ : \approx 6.4 \times 10^2$
  - $\pi^+ : \approx 3.4 \times 10^2$

# Prospects at experiments

Why counting rates not as high for UPCs at LHC?



- ▶ Photon flux enhanced by a factor of  $Z^2$ , but drops rapidly with  $S_{\gamma N} \implies$  *Low luminosity not compensated by larger photon flux.*
- ▶ LHC great for high energy, but JLab better in terms of luminosity.
- ▶ Still, LHC gives us access to the small  $\xi$  region of GPDs!

# Conclusion

- ▶ Exclusive photoproduction of photon-meson pair provides additional channel for **extracting GPDs**.

# Conclusion

- ▶ Exclusive photoproduction of photon-meson pair provides additional channel for **extracting GPDs**.
- ▶ Especially interesting since it can probe **chiral-odd GPDs** at the leading twist, and provides **better sensitivity to  $x$ -dependence of GPDs**.



# Conclusion

- ▶ Exclusive photoproduction of photon-meson pair provides additional channel for **extracting GPDs**.
- ▶ Especially interesting since it can probe **chiral-odd GPDs** at the leading twist, and provides **better sensitivity to  $x$ -dependence of GPDs**.
- ▶ **Proof of factorisation** for this family of processes now available, but intriguing **\*indication\*** of violation of collinear factorisation at twist-2 with **gluonic contributions to  $\pi^0\gamma$  photoproduction**.

# Conclusion

- ▶ Exclusive photoproduction of photon-meson pair provides additional channel for **extracting GPDs**.
- ▶ Especially interesting since it can probe **chiral-odd GPDs** at the leading twist, and provides **better sensitivity to  $x$ -dependence of GPDs**.
- ▶ **Proof of factorisation** for this family of processes now available, but intriguing **\*indication\*** of violation of collinear factorisation at twist-2 with **gluonic contributions to  $\pi^0\gamma$  photoproduction**.
- ▶ **Good statistics** in various experiments, particularly at JLab.

# Conclusion

- ▶ Exclusive photoproduction of photon-meson pair provides additional channel for **extracting GPDs**.
- ▶ Especially interesting since it can probe **chiral-odd GPDs** at the leading twist, and provides **better sensitivity to  $x$ -dependence of GPDs**.
- ▶ **Proof of factorisation** for this family of processes now available, but intriguing **\*indication\*** of violation of collinear factorisation at twist-2 with **gluonic contributions to  $\pi^0\gamma$  photoproduction**.
- ▶ **Good statistics** in various experiments, particularly at JLab.
- ▶ **Small  $\xi$  limit** of GPDs can be investigated by exploiting high energies available in collider mode such as EIC and UPCs at LHC.

- ▶ Resolve **gluonic GPD** contributions to  $\gamma N \rightarrow \gamma \pi^0 N$  [ongoing]

- ▶ Resolve **gluonic GPD** contributions to  $\gamma N \rightarrow \gamma \pi^0 N$  [ongoing]
- ▶ Compute **NLO** corrections (422 NLO diagrams, vs 20 LO diagrams!) [ongoing]


- ▶ Resolve **gluonic GPD** contributions to  $\gamma N \rightarrow \gamma \pi^0 N$  [ongoing]
- ▶ Compute **NLO** corrections (422 NLO diagrams, vs 20 LO diagrams!) [ongoing]
- ▶ Generalise to **electroproduction** ( $Q^2 \neq 0$ ).
- ▶ Add **Bethe-Heitler** component (photon emitted from incoming lepton)
  - zero in chiral-odd case.
  - suppressed in chiral-even case.

## BACKUP SLIDES

# Computation

## Hard Part: Diagrams

A total of 20 diagrams to compute



- ▶ Need to compute 10 diagrams: Other half related by  $q \leftrightarrow \bar{q}$  (anti)symmetry.
- ▶ In fact, by choosing the **right gauge**, **only 4 diagrams** can be used to generate all the others by various symmetries (eg. photon exchange).
- ▶ **Red** diagrams **cancel** in the chiral-odd case



# Computation

Parametrising the GPDs:  $\rho_L$  and  $\pi$  case, Chiral-even

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p_2, \lambda_2 | \bar{\psi}_q \left( -\frac{1}{2}z^- \right) \gamma^+ \psi \left( \frac{1}{2}z^- \right) | p_1, \lambda_1 \rangle$$
$$= \frac{1}{2P^+} \bar{u}(p_2, \lambda_2) \left[ H^q(x, \xi, t) \gamma^+ + E^q(x, \xi, t) \frac{i\sigma^{\alpha+} \Delta_\alpha}{2m} \right] u(p_1, \lambda_1)$$

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p_2, \lambda_2 | \bar{\psi}_q \left( -\frac{1}{2}z^- \right) \gamma^+ \gamma^5 \psi \left( \frac{1}{2}z^- \right) | p_1, \lambda_1 \rangle$$
$$= \frac{1}{2P^+} \bar{u}(p_2, \lambda_2) \left[ \tilde{H}^q(x, \xi, t) \gamma^+ \gamma^5 + \tilde{E}^q(x, \xi, t) \frac{\gamma^5 \Delta^+}{2m} \right] u(p_1, \lambda_1)$$

- ▶ Take the limit  $\Delta_\perp = 0$ .
- ▶ In that case and for small  $\xi$ , the dominant contributions come from  $H^q$  and  $\tilde{H}^q$ .

# Computation

Parametrising the GPDs:  $\rho_T$  case, Chiral-odd

$$\begin{aligned} & \int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p_2, \lambda_2 | \bar{\psi}_q \left( -\frac{1}{2}z^- \right) i\sigma^{+i} \psi \left( \frac{1}{2}z^- \right) | p_1, \lambda_1 \rangle \\ &= \frac{1}{2P^+} \bar{u}(p_2, \lambda_2) \left[ H_T^q(x, \xi, t) i\sigma^{+i} + \tilde{H}_T^q(x, \xi, t) \frac{P^+ \Delta^i - \Delta^+ P^i}{m_N^2} \right. \\ &+ \left. E_T^q(x, \xi, t) \frac{\gamma^+ \Delta^i - \Delta^+ \gamma^i}{2m_N} + \tilde{E}_T^q(x, \xi, t) \frac{\gamma^+ P^i - P^+ \gamma^i}{m_N} \right] u(p_1, \lambda_1) \end{aligned}$$

- ▶ Take the limit  $\Delta_\perp = 0$ .
- ▶ In that case and for small  $\xi$ , the dominant contributions come from  $H_T^q$ .

- ▶ GPDs can be represented in terms of **Double Distributions**  
[Radyushkin: hep-ph/9805342]

$$H^q(x, \xi, t = 0) = \int_{-1}^1 d\beta \int_{-1+|\beta|}^{1-|\beta|} d\alpha \delta(\beta + \xi\alpha - x) f^q(\beta, \alpha)$$

- ▶ ansatz for these Double Distributions:

- ▶ chiral-even sector:

$$\begin{aligned} f^q(\beta, \alpha, t = 0) &= \Pi(\beta, \alpha) q(\beta) \Theta(\beta) - \Pi(-\beta, \alpha) \bar{q}(-\beta) \Theta(-\beta), \\ \tilde{f}^q(\beta, \alpha, t = 0) &= \Pi(\beta, \alpha) \Delta q(\beta) \Theta(\beta) + \Pi(-\beta, \alpha) \Delta \bar{q}(-\beta) \Theta(-\beta). \end{aligned}$$

- ▶ chiral-odd sector:

$$f_T^q(\beta, \alpha, t = 0) = \Pi(\beta, \alpha) \delta q(\beta) \Theta(\beta) - \Pi(-\beta, \alpha) \delta \bar{q}(-\beta) \Theta(-\beta).$$

- ▶  $\Pi(\beta, \alpha) = \frac{3}{4} \frac{(1-\beta)^2 - \alpha^2}{(1-\beta)^3}$  : profile function

- ▶ simplistic factorised ansatz for the  $t$ -dependence:

$$H^q(x, \xi, t) = H^q(x, \xi, t = t_{\min}) \times F_H(t)$$

with  $F_H(t) = \frac{(t_{\min} - C)^2}{(t - C)^2}$  a standard **dipole form factor**  
( $C = 0.71 \text{ GeV}^2$ )

# Sets of PDFs used to model GPDs

- ▶  $q(x)$  : unpolarised PDF:
  - GRV-98 [Glück, Reya, Vogt: hep-ph/9806404]
  - MSTW2008lo [Martin, Stirling, Thorne, Watt: 0901.0002]
  - MSTW2008nnlo [Martin, Stirling, Thorne, Watt: 0901.0002]
  - ABM11nnlo [Alekhin, Blumlein, Moch: 1202.2281]
  - CT10nnlo [Gao, Guzzi, Huston, Lai, Li, Nadolsky, Pumplin, Stump, Yuan: 1302.6246]
- ▶  $\Delta q(x)$  polarised PDF
  - GRSV-2000 [Glück, Reya, Stratmann, Vogelsang: hep-ph/0011215]
- ▶  $\delta q(x)$  : transversity PDF:
  - Based on parameterisation for TMDs from which transversity PDFs obtained as limiting case [Anselmino, Boglione, D'Alesio, Melis, Murgia, Prokudin: 1303.3822]

Effects are not significant! But relevant for NLO corrections!

- Helicity conserving (vector) DA at twist 2:  $\rho_L$

$$\langle 0 | \bar{u}(0) \gamma^\mu u(x) | \rho_L^0(p) \rangle = \frac{p^\mu}{\sqrt{2}} f_\rho \int_0^1 du e^{-iup \cdot x} \phi_\rho(u)$$

- Helicity flip (tensor) DA at twist 2:  $\rho_T$

$$\langle 0 | \bar{u}(0) \sigma^{\mu\nu} u(x) | \rho_T^0(p, s) \rangle = \frac{i}{\sqrt{2}} (\epsilon_\rho^\mu p^\nu - \epsilon_\rho^\nu p^\mu) f_\rho^\perp \int_0^1 du e^{-iup \cdot x} \phi_\rho(u)$$

- Helicity conserving (axial) DA at twist 2:  $\pi^\pm$

$$\langle 0 | \bar{u}(0) \gamma^\mu \gamma^5 d(x) | \pi(p) \rangle = ip^\mu f_\pi \int_0^1 du e^{-iup \cdot x} \phi_\pi(u)$$


# Computation

Valence vs Standard scenarios in  $\tilde{H}$  (Chiral-even, Axial)

Typical kinematic point (for JLab kinematics):

$$\xi = .1 \leftrightarrow S_{\gamma N} = 20 \text{ GeV}^2 \text{ and } M_{\gamma\rho}^2 = 3.5 \text{ GeV}^2$$

$$\tilde{H}^{q(-)}(x, \xi, t) = \tilde{H}^q(x, \xi, t) - \tilde{H}^q(-x, \xi, t) \quad [C = -1]$$



“valence” and “standard”: two GRSV Ansätze for  $\Delta q(x)$


# Computation

Valence vs Standard scenarios in  $H_T$  (Chiral-odd)

Typical kinematic point (for JLab kinematics):

$$\xi = .1 \leftrightarrow S_{\gamma N} = 20 \text{ GeV}^2 \text{ and } M_{\gamma\rho}^2 = 3.5 \text{ GeV}^2$$

$$H_T^{q(-)}(x, \xi, t) = H_T^q(x, \xi, t) + H_T^q(-x, \xi, t) \quad [C = -1]$$



“valence” and “standard”: two GRSV Ansätze for  $\Delta q(x)$

$\Rightarrow$  two Ansätze for  $\delta q(x)$



# Computation

## Kinematics

- ▶ Work in the limit of:

- $\Delta_{\perp} \ll p_{\perp}$
- $m_N^2, m_M^2 \ll M_{\gamma M}^2$

- ▶ initial state particle momenta:

$$q^{\mu} = n^{\mu},$$


$$p_1^{\mu} = (1 + \xi) p^{\mu} + \frac{m_N^2}{s(1+\xi)} n^{\mu}$$

- ▶ final state particle momenta:

$$p_2^{\mu} = (1 - \xi) p^{\mu} + \frac{m_N^2 + \vec{p}_t^2}{s(1 - \xi)} n^{\mu} + \Delta_{\perp}^{\mu}$$

$$k^{\mu} = \alpha n^{\mu} + \frac{(\vec{p}_t - \vec{\Delta}_t/2)^2}{\alpha s} p^{\mu} + p_{\perp}^{\mu} - \frac{\Delta_{\perp}^{\mu}}{2}, \quad \Delta \downarrow$$


$$p_M^{\mu} = \alpha_M n^{\mu} + \frac{(\vec{p}_t + \vec{\Delta}_t/2)^2 + m_M^2}{\alpha_M s} p^{\mu} - p_{\perp}^{\mu} - \frac{\Delta_{\perp}^{\mu}}{2},$$



} hard scale  
 $M_{\gamma M}^2 \propto p_{\perp}^2$

# Exclusive photoproduction of $\pi^0\gamma$

Gluonic GPD contributions



$$D_a = ((x - \xi)p + \bar{z}p_M)^2 + i\epsilon$$

$$= s\bar{\alpha}\bar{z}[x - \xi + i\epsilon] ,$$

$$D_b = (k + zp_M - (x + \xi)p)^2 + i\epsilon$$

$$= -s[z(x - \xi - i\epsilon) + \alpha\bar{z}(x + \xi - i\epsilon)] ,$$



$$D_c = (zp_M - (x + \xi)p)^2 + i\epsilon$$

$$= -s\bar{\alpha}z[x + \xi - i\epsilon]$$

$\implies$  pinching of poles in the propagators ( $D_a$  and  $D_b$ ) in the limit of  $z \rightarrow 1$

# Results

Fully-differential cross-sections:  $\gamma\rho_{\rho L}^+$  vs  $\gamma\rho_{\rho L}^-$



$$S_{\gamma N} = 20 \text{ GeV}^2, \quad -t = (-t)_{\min}, \quad M_{\gamma\rho}^2 = 3, 4, 5 \text{ GeV}^2$$

Dashed: Holographic DA      non-dashed: Asymptotical DA


Dotted: standard scenario      non-dotted: valence scenario

# Results


Phase space integration: Evolution in  $(-t, -u')$  plane

large angle scattering:  $M_{\gamma\rho}^2 \sim -u' \sim -t'$  ( $S_{\gamma N} = 20 \text{ GeV}^2$ )


$\implies -u' > 1 \text{ GeV}^2$  and  $-t' > 1 \text{ GeV}^2$  and  $(-t)_{\min} \leq -t \leq .5 \text{ GeV}^2$




$M_{\gamma\rho} = 2.2 \text{ GeV}^2$




$M_{\gamma\rho} = 2.5 \text{ GeV}^2$




$M_{\gamma\rho} = 3 \text{ GeV}^2$



$M_{\gamma\rho} = 5 \text{ GeV}^2$




$M_{\gamma\rho} = 8 \text{ GeV}^2$



$M_{\gamma\rho} = 9 \text{ GeV}^2$

# Results

## Necessity for Importance Sampling



- ▶ Need enough points at boundaries for distribution in  $(-u')$
- ▶ Need enough points to resolve peak (at low  $M_{\gamma\rho_L^0}^2$ ) for distribution in  $M_{\gamma\rho_L^0}^2$

# Results

Explaining the difference between chiral-even and chiral-odd plots

▶  $\xi = \frac{M_{\gamma M}^2}{2S_{\gamma N} - M_N^2} \approx \frac{M_{\gamma M}^2}{2S_{\gamma N}}$  for  $M_{\gamma M}^2 \ll S_{\gamma N}$

▶ Chiral-even (unpolarised) cross-section:

$$|\overline{\mathcal{M}}_{\text{CE}}|^2 = \frac{2}{s^2} (1 - \xi^2) C_{\text{CE}}^2 \left\{ 2 |N_A|^2 + \frac{p_{\perp}^4}{s^2} |N_B|^2 + \frac{p_{\perp}^2}{s} (N_A N_B^* + \text{c.c.}) + \frac{p_{\perp}^4}{4s^2} |N_{A_5}|^2 + \frac{p_{\perp}^4}{4s^2} |N_{B_5}|^2 \right\}.$$

▶ Chiral-odd (unpolarised) cross-section:

$$|\overline{\mathcal{M}}_{\text{CO}}|^2 = \frac{2048}{s^2} \xi^2 (1 - \xi^2) C_{\text{CO}}^2 \left\{ \alpha^4 |N_{TA}|^2 + |N_{TB}|^2 \right\}.$$

▶ Note:  $\alpha = \frac{-u'}{M_{\gamma M}^2}$ .

# Results

Integrated cross-section: Mapping procedure for different values of  $S_{\gamma N}$

To obtain distribution in  $S_{\gamma N}$ , we exploit non-trivial mapping between 1 set of data at a fixed  $S_{\gamma N}$  to other values  $\tilde{S}_{\gamma N}$  *lower* than it.

$$\tilde{M}_{\gamma M}^2 = M_{\gamma M}^2 \frac{\tilde{S}_{\gamma N} - m_N^2}{S_{\gamma N} - m_N^2},$$
$$- \tilde{u}' = \frac{\tilde{M}_{\gamma M}^2}{M_{\gamma M}^2} (-u').$$

Implementing **importance sampling**  $\implies$  careful consideration of the various limits involved are needed.

Mapping possible since **different** sets of  $(S_{\gamma N}, M_{\gamma M}^2, -u')$  correspond to the **same**  $(\alpha, \xi)$ .

$$\alpha = \frac{-u'}{M_{\gamma M}^2}, \quad \xi = \frac{M_{\gamma M}^2}{2(S_{\gamma N} - m_N^2) - M_{\gamma M}^2}.$$

# Results

Why does the circular asymmetry vanish for unpolarised target?

Consider

$$\gamma(q, \lambda_q) + N(p_1, \lambda_1) \rightarrow \gamma(k, \lambda_k) + \pi^\pm(p_\pi) + N'(p_2, \lambda_2),$$

where  $\lambda_i$  represent the helicities of the particles.

QED/QCD **invariance under parity** implies that [Bourenly, Soffer, Leader: Phys.Rept. 59 (1980) 95-297]

$$\mathcal{A}_{\lambda_2\lambda_k; \lambda_1\lambda_q} = \eta(-1)^{\lambda_1 - \lambda_q - (\lambda_2 - \lambda_k)} \mathcal{A}_{-\lambda_2 - \lambda_k; -\lambda_1 - \lambda_q},$$

where  $\eta$  represents phase factors related to intrinsic spin.



Thus, at the cross-section level, it is clear that circular asymmetry will vanish, since

$$\sum_{\lambda_i, i \neq q} |\mathcal{A}_{\lambda_2\lambda_k; \lambda_1+}|^2 = \sum_{\lambda_i, i \neq q} |\mathcal{A}_{\lambda_2\lambda_k; \lambda_1-}|^2$$



# Results

LPA wrt incoming photon: Fully-differential level:  $\gamma\rho_{PL}^0$  vs  $\gamma\rho_{PL}^+$




$$S_{\gamma N} = 20 \text{ GeV}^2, -t = (-t)_{\min}, M_{\gamma\rho}^2 = 3, 4, 5 \text{ GeV}^2$$

Dashed: Holographic DA      non-dashed: Asymptotical DA

Dotted: standard scenario      non-dotted: valence scenario

# Angular cuts on outgoing photon at JLab


Angular distribution:  $\rho_p^0 \gamma$  photoproduction at  $S_{\gamma N} = 20 \text{ GeV}^2$



- ▶  $M_{\gamma\rho_p^0}^2 = 4 \text{ GeV}^2$  (solid blue)
- ▶  $M_{\gamma\rho_p^0}^2 = 6 \text{ GeV}^2$  (dotted red)
- ▶  $M_{\gamma\rho_p^0}^2 = 8 \text{ GeV}^2$  (dashed green)

# Angular cuts on outgoing photon at JLab

Single differential cross-section:  $\rho_p^0 \gamma$  photoproduction at  $S_{\gamma N} = 20 \text{ GeV}^2$



- ▶ no angular cut (solid red)
- ▶  $\theta \leq 35^\circ$  (dashed blue)
- ▶  $\theta \leq 30^\circ$  (dotted green)
- ▶  $\theta \leq 25^\circ$  (dashed-dotted brown)
- ▶  $\theta \leq 20^\circ$  (long-dashed magenta)
- ▶  $\theta \leq 15^\circ$  (short-dashed purple)
- ▶  $\theta \leq 10^\circ$  (dotted black)