

Generalized Parton Distributions through Universal Moment Parameterization (GUMP): Towards global analysis at non zero skewness

1

M Gabriel Santiago

With Yuxun Guo, Xiangdong Ji, Kyle Shiells and Jinghong Yang

JHEP 05 (2023) 150, [arXiv:2302.07279](https://arxiv.org/abs/2302.07279)

Outline


- ▶ GPD Review
- ▶ GUMP Program
 - ▶ Conformal moment parameterization
- ▶ First Step Towards Global Analysis: u and d quarks
 - ▶ Simplified GPD moment ansatz
 - ▶ Experimental and lattice input
- ▶ Non-zero Skewness Global Fit
- ▶ GPD Extraction
 - ▶ Ambiguity in the ERBL/DA-like region
 - ▶ D-terms vs DA-like terms
- ▶ Moving Forward
 - ▶ Gluons from J/ψ
 - ▶ NLO corrections
 - ▶ Simultaneous DVCS and DVMP
 - ▶ Full global fitting
- ▶ Conclusions

GPDs

- ▶ GPDs generalize the well known PDFs to encode full 3 dimensional information on the quarks and gluons within hadrons

$$f(x) \rightarrow F(x, \xi, t)$$

$x \sim$ parton momentum fraction, $\xi \sim$ longitudinal momentum transfer,
 $t = \Delta^2 \sim$ momentum transfer squared




GPDs

- Polarization of the hadron and its parton constituents connects GPDs to the distribution of angular momentum within hadrons (X. Ji 1997)

- Ji sum rule
$$J_i = \frac{1}{2} \int_0^1 dx x [H_i(x, \xi) + E_i(x, \xi)]$$

- Related via a Fourier transform to the impact parameter distribution of partons (M. Burkardt 2003)

$$\rho(x, r_{\perp}) = \int \frac{d^2 \Delta_{\perp}}{(2\pi)^2} e^{-i\Delta_{\perp} \cdot r_{\perp}} H(x, 0, \Delta_{\perp}^2)$$



- Related to bulk properties of hadron states encoded in form factors

$$\int dx x H_i(x, \xi, t) = A_i(t) + (2\xi)^2 C_i(t), \quad \int dx x E_i(x, \xi, t) = B_i(t) - (2\xi)^2 C_i(t)$$

GUMP program: Moment Parameterization

- Parameterize GPDs by directly parameterizing their conformal moments

$$F_i(x, \xi, t) = \sum_{j=0}^{\infty} (-1)^j p_{i,j}(x, \xi) \mathcal{F}_{i,j}(\xi, t) \quad (\text{D. Mueller and A. Schafer 2006})$$

- Expansion based on eigenfunctions of evolution – Gegenbauer polynomials

$$(-1)^j p_j(x, \xi) = \xi^{-j-1} \frac{2^j \Gamma(\frac{5}{2} + j)}{\Gamma(\frac{3}{2}) \Gamma(j+3)} \left[1 - \left(\frac{x}{\xi} \right)^2 \right] C_j^{3/2} \left(\frac{x}{\xi} \right)$$

conformal wave
function

$$\int_{-1}^1 \frac{dx'}{|\xi|} \mathcal{K} \left(\frac{x}{\xi}, \frac{x'}{\xi} \right) C_j^{3/2} \left(\frac{x}{\xi} \right) = \gamma_j C_j^{3/2} \left(\frac{x}{\xi} \right)$$

GPD evolution
kernel

GUMP program: Moment Parameterization

- Conformal moment parameterization has nice features for fitting GPDs
- Simple evolution implementation – conformal moments are multiplicatively renormalized at LO
 - Follows from using eigenfunctions of evolution kernel
- Polynomiality condition (X. Ji 1998) automatically enforced on conformal moments

$$F_{i,n}(\xi, t) = \int_{-1}^1 dx x^{n-1} F(x, \xi, t) = \sum_{k=0, \text{ even}}^n \xi^k F_{i,n,k}(t)$$

$$\mathcal{F}_{i,j}(\xi, t) = \sum_{k=0, \text{ even}}^{j+1} \xi^k \mathcal{F}_{i,j,k}(t)$$

First Step Toward Global GPD Analysis

- Apply in GUMP program for global analysis of u and d quark GPDs at non-zero skewness with LO scale evolution
- Parameterize each GPD moment with five parameters

$$F_{i,j,0} = N_i B(j+1-\alpha_i, 1+\beta_i) \frac{j+1-\alpha_i}{j+1-\alpha_i(t)} \beta(t)$$

↑
↑
 Euler Beta Regge trajectory
 Function

$$\beta(t) = e^{-b|t|}$$

$$\alpha(t) = \alpha + \alpha' t$$

- Take each moment to be a power series in skewness – polynomiality condition

$$F_{i,j} = F_{i,j,0}(t) + \xi^2 R_{\xi^2} F_{i,j,0}(t) + \xi^4 R_{\xi^4} F_{i,j,0}(t) \dots$$

First Step Toward Global GPD Analysis

- The number of parameters needed for modelling all the species of GPD grows very quickly
- We impose extra constraints for simplicity

GPDs species and flavors	Fully parameterized	GPDs linked to	Proportional constants
H_{u_V} and \tilde{H}_{u_V}	✓	-	-
E_{u_V} and \tilde{E}_{u_V}	✓	-	-
H_{d_V} and \tilde{H}_{d_V}	✓	-	-
E_{d_V} and \tilde{E}_{d_V}	✗	E_{u_V} and \tilde{E}_{u_V}	$R_{d_V}^{E/\tilde{E}}$
$H_{\bar{u}}$ and $\tilde{H}_{\bar{u}}$	✓	-	-
$E_{\bar{u}}$ and $\tilde{E}_{\bar{u}}$	✗	$H_{\bar{u}}$ and $\tilde{H}_{\bar{u}}$	$R_{\text{sea}}^{E/\tilde{E}}$
$H_{\bar{d}}$ and $\tilde{H}_{\bar{d}}$	✓	-	-
$E_{\bar{d}}$ and $\tilde{E}_{\bar{d}}$	✗	$H_{\bar{d}}$ and $\tilde{H}_{\bar{d}}$	$R_{\text{sea}}^{E/\tilde{E}}$
H_g and \tilde{H}_g	✓	-	-
E_g and \tilde{E}_g	✗	H_g and \tilde{H}_g	$R_{\text{sea}}^{E/\tilde{E}}$

Non-zero Skewness Global Fit

- ▶ Even with constraints, lots of parameters!
 - ▶ Very high dimensional space to navigate for best fit
 - ▶ Very computationally demanding to do error propagation
- ▶ We employ a sequential fit, starting with forward (PDF, t-dependent PDF) constraints for each GPD species then apply the off-forward constraints from DVCS data



Semi-Forward Inputs


- ▶ JAM (2022) PDF global analysis results
 - ▶ Full global analysis should in principle fit to PDF sensitive data directly, but here we fit to JAM results
 - ▶ Limited number of points taken to avoid need for more sophisticated forward limit
- ▶ Globally extracted electromagnetic form factors (*Z. Ye et al 2018*)
- ▶ Lattice GPDs (*Alexandrou et al 2020*) and form factors (*Alexandrou et al 2022*)
 - ▶ x, t -dependent GPDs (semi-forward limit)

Off-Forward Inputs

- DVCS measurements from JLab (*CLAS* 2019 & 2021, *Hall A* 2018 & 2022) and HERA (*H1* 2010)
- Only using t -dependent cross sections due to practical limitations
- Far more points from JLab data than from HERA from φ -dependence and both UU and LU polarization channels
- Off-forward lattice GPDs not used in fitting, but can supply crucial constraints for future work!


Non-zero Skewness Global Fit

- Total χ^2/dof is approximately 1.4
- Some agreement with both JLAB and H1 data
- Gluon GPDs not well constrained at non-zero skewness
 - Only contribute to DVCS through evolution at LO
- Error propagation is not yet implemented
 - Very computationally expensive with so many parameters!




Non-zero Skewness Global Fit

- CFFs from fit are mostly consistent with local extraction from JLAB Hall A data as well as KM15 extractions
- Some inconsistencies can be expected from degeneracies in CFF contribution to cross sections – need more polarization configurations!



Extracted GPDs

- ▶ GPDs are mostly constrained on the $\xi = x$ line and in the DGLAP region $|\xi| < |x|$
- ▶ ERBL region shows large oscillations which are characteristic of the Gegenbauer polynomials used in the moment expansion



Ambiguity in ERBL Region

- We can add terms in the moment expansion which only contribute to the ERBL region

$$(-1)^j p_j(x, \xi) = \xi^{-j-1} \frac{2^j \Gamma(\frac{5}{2} + j)}{\Gamma(\frac{3}{2}) \Gamma(j+3)} \left[1 - \left(\frac{x}{\xi} \right)^2 \right] C_j^{3/2} \left(\frac{x}{\xi} \right), \quad |x| < |\xi|$$

- This suggests an interpretation of the GPDs in terms of quark and antiquark pieces as well as a ERBL region distribution amplitude (DA) piece

$$F_q(x, \xi, t) = F_{\hat{q}}(x, \xi, t) \mp F_{\bar{q}}(-x, \xi, t) + F_{q\bar{q}}$$

quark
 $x > -\xi$

antiquark
 $-x > -\xi$

DA
 $\xi > x > -\xi$

Connection to D-term

- These DA terms don't have a large affect on CFFs, but they do contain information related to the various D-terms in QCD, ex.

- Gravitational form factor C/D

$$\int_{-1}^1 dx x H_q(x, \xi, t) = A_q(t) + (2\xi)^2 C_q(t)$$


- Dispersion relation subtraction term

$$F(\xi, t, Q^2) = \frac{1}{\pi} \int_0^1 d\xi' \left(\frac{1}{\xi - \xi'} \mp \frac{1}{\xi + \xi'} \right) \text{Im} [F(\xi' - i0, t, Q^2)] + C(t, Q^2)$$

- By constraining the DA terms with further experimental data and lattice calculations, we can access the mechanical properties of hadrons contained in these D-terms!


Constraining DA Terms

- ▶ Adding in lattice GPD calculations can give us constraints directly in the ERBL region
- ▶ Adding just a few terms to the moment expansion can remove the unphysical oscillations



Moving Forward: Adding in Gluons!

- ▶ DVCS at LO is only sensitive to gluon GPDs through scale evolution
- ▶ Using Deeply Virtual Meson Production (DVMP) gives a direct probe of gluons at LO



- ▶ Light vector mesons have similar sensitivity to quarks and gluons
 - ▶ KM framework applied to produce simultaneous fits of DVCS and DVMP for ρ^0 and ϕ meson production with data from HERA (M. Meskauskas and D. Muller 2011)
- ▶ Add heavy vector meson to obtain better constraints on gluon GPDs – use J/ψ production!

Deeply Virtual J/ψ Production

- ▶ Charm quark contribution for nucleon target is negligible – direct probe of gluons
- ▶ Complementary with GUMP work on quark GPDs, but mostly sensitive to small- x_B region whereas JLab data combined with HERA gives better constraint at moderate x_B
- ▶ Caveat: mass of the J/ψ is too large for usual collinear factorization

$$M_{J/\psi}^2 / Q_{\text{max bin}}^2 \approx 9/20 \rightarrow \text{corrections of order } 1/2$$

- ▶ Need to take heavy mass corrections into account – non-relativistic (NR) QCD!

Non-relativistic model approach

- Encoding the J/ψ formation into NR matrix elements

$$\Gamma[J/\psi \rightarrow l^+l^-] \approx \frac{2e_c^2 \pi \alpha_{EM}^2}{3m_c^2} \langle \mathcal{O}_1 \rangle_{J/\psi} \left(1 - \frac{8\alpha_s}{3\pi}\right)^2$$

- Maintain the form of the factorization theorem for the process – still sensitive to leading twist GPDs (D. Y. Ivanov et al 2004)

$$\mathcal{M} = \left(\frac{\langle \mathcal{O}_1 \rangle_V}{m_c}\right)^{1/2} \sum_i F_i(x, \xi, t) \otimes_x H_i(x, \xi)$$

- Systematically improvable with relativistic, higher twist, and NLO QCD corrections
- Bridge between electroproduction and photoproduction regimes

Implementing NR J/ψ Production in GUMP

- ▶ LO framework used for previous global fit does not match data in HERA kinematics
- ▶ NLO evolution is known in moment space (*Mueller et al 2013*)
- ▶ Finite mass corrections are only known in momentum fraction space
- ▶ Numerical complex integral to construct GPD from moments is computationally expensive

Future Improvements/Additions

- Implement J/ψ electroproduction fits with NLO
- Add threshold J/ψ production – potentially constrain D-term/DA-terms
- Add quark flavors and implement ρ^0 and ϕ electroproduction
- Full simultaneous global analysis with DVCS and DVMP contributions
- Implement t -integrated cross sections – speed up for NLO could make t -integrated cross sections practical

Conclusions

- Global fit combining experimental data and lattice calculations to constrain GPDs at non-zero skewness
- Developing the GUMP program to include gluon GPDs in global analysis through J/ψ production data
- Implementing NLO corrections
- Several directions for future improvements available

Backup Slides

Best Fit χ^2 Breakdown

Sub-fits	χ^2	N_{data}	$\chi^2_{\nu} \equiv \chi^2/\nu$
Semi-forward			
<i>t</i> PDF H	281.7	217	1.41
<i>t</i> PDF E	59.7	50	1.36
<i>t</i> PDF \tilde{H}	159.3	206	0.84
<i>t</i> PDF \tilde{E}	63.8	58	1.23
Off-forward			
JLab DVCS	1413.7	926	~ 1.53
H1 DVCS	19.7	24	~ 0.82
Off-forward total	1433	950	1.53
Total	2042	1481	1.40

Best Fit Parameters

Vector GPDs H and E		Axial-vector GPDs \tilde{H} and \tilde{E}	
Parameter	Value (uncertainty)	Parameter	Value (uncertainty)
N_{uV}^H	4.923 (89)	$N_{uV}^{\tilde{H}}$	4.833 (429)
α_{uV}^H	0.216 (7)	$\alpha_{uV}^{\tilde{H}}$	-0.264 (34)
β_{uV}^H	3.229 (23)	$\beta_{uV}^{\tilde{H}}$	3.186 (122)
α'_{uV}^H	2.347 (51)	$\alpha'_{uV}^{\tilde{H}}$	2.182 (175)
$N_{\bar{u}}^H$	0.163 (8)	$N_{\bar{u}}^{\tilde{H}}$	0.070 (33)
$\alpha_{\bar{u}}^H$	1.136 (10)	$\alpha_{\bar{u}}^{\tilde{H}}$	0.538 (112)
$\beta_{\bar{u}}^H$	6.894 (207)	$\beta_{\bar{u}}^{\tilde{H}}$	4.229 (1320)
N_{dV}^H	3.359 (170)	$N_{dV}^{\tilde{H}}$	-0.664 (170)
α_{dV}^H	0.184 (18)	$\alpha_{dV}^{\tilde{H}}$	0.248 (76)
β_{dV}^H	4.418 (77)	$\beta_{dV}^{\tilde{H}}$	3.572 (477)
α'_{dV}^H	3.482 (171)	$\alpha'_{dV}^{\tilde{H}}$	0.542 (103)
$N_{\bar{d}}^H$	0.249 (12)	$N_{\bar{d}}^{\tilde{H}}$	-0.086 (42)
$\alpha_{\bar{d}}^H$	1.052 (10)	$\alpha_{\bar{d}}^{\tilde{H}}$	0.495 (137)
$\beta_{\bar{d}}^H$	6.554 (216)	$\beta_{\bar{d}}^{\tilde{H}}$	2.554 (897)
N_g^H	2.864 (108)	$N_g^{\tilde{H}}$	0.243 (304)
α_g^H	1.052 (8)	$\alpha_g^{\tilde{H}}$	0.631 (330)
β_g^H	7.413 (165)	$\beta_g^{\tilde{H}}$	2.717 (2865)
N_{uV}^E	0.181 (38)	$N_{uV}^{\tilde{E}}$	7.993 (3480)
α_{uV}^E	0.907 (17)	$\alpha_{uV}^{\tilde{E}}$	0.800 (116)
β_{uV}^E	1.102 (245)	$\beta_{uV}^{\tilde{E}}$	6.415 (1577)
α'_{uV}^E	0.461 (86)	$\alpha'_{uV}^{\tilde{E}}$	2.076 (933)
N_{dV}^E	-0.223 (47)	$N_{dV}^{\tilde{E}}$	-2.407 (1239)
R_{sea}^E	0.768 (169)	$R_{sea}^{\tilde{E}}$	38 (8)
$R_{u,2}^H$	0.229 (0.032)	$R_{u,2}^{\tilde{H}}$	0.246 (81)
$R_{d,2}^H$	-2.639 (202)	$R_{d,2}^{\tilde{H}}$	1.656 (375)
$R_{u,2}^E$	0.799 (285)	$R_{u,2}^{\tilde{E}}$	2.684 (171)
$R_{d,2}^E$	3.404 (1157)	$R_{d,2}^{\tilde{E}}$	38 (2)
υ_{sea}^H	3.448 (133)	$\upsilon_{sea}^{\tilde{H}}$	9.852 (1330)