

# *Mediator Particle Decay at the Threshold*

*Shigeki Matsumoto*

*(Kavli IPMU, U. Tokyo)*

*Collaborators: Graham White, Yu Watanabe, Yuki Watanabe  
(arXiv:2212.10739)*

- ✓ *Light mediator is often predicted in the dark sector scenario.*
- ✓ *The mediator particle weakly interacts with SM particles, and a small coupling controls the strength of the interaction.*
- ✓ *The strength of the interaction is influenced by the threshold singularity when the mediator decays at a threshold region.*
- ✓ *We discuss how this influence is quantitatively incorporated and how much modification we have on the interaction strength.*

# Dark sector scenario

*A scenario addressing big questions in PP (DM,  $\nu$  mass, BAU, etc.)*

*< EW scale*

*Standard Model*

*Weak Connection*

*Dark sector*

- ✓ *Via higher-dim. operator (Or very heavy particles).*
- ✓ *Hidden valley scenario.*
- ✓ *Via light mediator particle*

*Origin of*

- ✓ *the dark matter,*
- ✓ *the neutrino masses/mixings,*
- ✓ *the baryon asymmetry of U., etc.*

*Ex.) Fermionic singlet (thermal) dark matter in the dark sector*

*Suppose that the dark matter is stabilized by the  $Z_2$  symmetry, and interacts with the SM particles via renormalizable interactions,*

*Any renormalizable interaction between DM and SM particles can not be written in an SM+DM system due to the  $Z_2$  and SM sym.*

*A mediator particle must be introduced in theory, and its mass must be light enough to explain the dark matter abundance observed today!*

# Mediator particle

*Various mediator particles are now being considered in the literature.*

*To guarantee weak interactions among the light mediator particle & SM particles, the mediator should be singlet under the SM gauge group.*

*The following three mediator particles are being intensively studied, as the simplest mediator candidates with the spin of 0,  $\frac{1}{2}$ , and 1.*

## Dark Higgs $\Phi$

✓ Spin-0 mediator

✓ Interaction(s)

$$\Phi|H|^2, \Phi^2|H|^2$$

**H: Higgs doublet**

*It causes the mixing between  $\phi$  and H, so the mediator tends to interact with heavy SM particles.*

## Dark photon $V$

✓ Spin-1 mediator

✓ Interaction

$$\text{e.g., } V_{\mu\nu} B^{\mu\nu}$$

**B: Hyper-charge gauge boson**

*It causes the mixing between  $V$  &  $\gamma$ , so it couples to charged SM particles.*

## Sterile neutrino $N$

✓ Spin- $\frac{1}{2}$  mediator

✓ Interaction

$$LHN + \text{h.c.}$$

**L: Lepton doublet**

*It causes the mixing between  $N$  and SM neutrinos  $\nu$ , so  $N$  interacts mainly with SM leptons.*

# Implications to collider physics & cosmology

***:: Light mediator particle is light and weakly interacting with the SM!***

## ***○ Typical production modes***



***Direct prod., e.g.,  $ee \rightarrow Med + X$ .***

***Boson decay, e.g.,  $B \rightarrow Med + X$ .***

## ***○ Typical decay modes***

***Decays into an SM particle pair, such as  $Med \rightarrow \mu\mu, KK, BB$ , etc.***

***It could be a long-lived particle at various collider experiments!***



***It could affect the evolution of the universe as a long-lived light relic.***

***✓ Spoiling the successful BBN.***

***✓ Modifying CMB via its decay.***

***✓ Pulling energy out from stars.***

***✓ Modifying CMB via  $V \leftrightarrow \gamma$ .***

***✓ May play a role of dark matter.***

# Vector mediator particle

- ✓ *Weak interaction of the mediator is guaranteed by a small coupling.*
- ✓ *It is not always valid when the mediator decays at a threshold.*
- ✓ *The mediator may have sizable interactions with a small coupling.*

*Let us consider such a case using the dark photon mediator scenario!*

## ✓ Starting Lagrangian is

$$\mathcal{L}_V = \mathcal{L}_{\text{SM}} - \frac{1}{4}(V_{\mu\nu})^2 + \frac{1}{2}M_V^2(V_\mu)^2 - \frac{\xi}{2}V_{\mu\nu}B^{\mu\nu} + \dots$$

*The mediator mass term is assumed to be from a Higgs mechanism.*

## ✓ Making Kinetic terms canonical and diagonalizing mass matrix gives

$$\mathcal{L}_V = \frac{1}{2}A'_\mu [(\square + m_{A'}^2)g^{\mu\nu} - \partial^\mu\partial^\nu]A'_\nu + \frac{\epsilon e m_Z^2}{m_Z^2 - m_{A'}^2} [A'_\mu J_{\text{EM}}^\mu + F_{A'W^\dagger W}] - \frac{\epsilon (g'/c_W) m_{A'}^2}{m_Z^2 - m_{A'}^2} A'_\mu J_Y^\mu + \dots$$


*where  $J_{\text{EM}}$  and  $J_Y$  are EM and hyper-charge currents,  $w/\epsilon = \xi \cos\theta_W$ .*

*The vector mediator particle interacts with the SM particles through the EM currents when its mass is smaller enough than the EW scale!*

# Decay width of the mediator particle

Let us consider the decay width of the mediator particle, for it gives its lifetime (relevant to collider physics) and the interaction strength between the mediator and the SM particles (relevant to cosmology).

- The mediator perturbatively decays into lepton & quark pairs at LO:




$$\Gamma_0(A' \rightarrow \ell_i^- \ell_i^+) \simeq \epsilon^2 \frac{\alpha m_{A'}}{3} \left( 1 + 2 \frac{m_{\ell_i}^2}{m_{A'}^2} \right) \sqrt{1 - \frac{4m_{\ell_i}^2}{m_{A'}^2}}$$

$$\Gamma_0(A' \rightarrow \text{all hadrons}) \simeq \epsilon^2 \frac{\alpha m_{A'}}{3} \left( 1 + 2 \frac{m_{\mu}^2}{m_{A'}^2} \right) \sqrt{1 - \frac{4m_{\mu}^2}{m_{A'}^2}} R$$

*Naive parton level calculation or the use of so-called the R-ratio are usually adopted.*

- Non-perturbative effect becomes sizable at the threshold region.



Positronium, muonic atom,  $J/\psi$ ,  $Y$ , etc.

**Threshold singularities (Sommerfeld effect, bound states) caused by long-range forces. Singularities appear not only for the quark final states (via QCD force) but also for the leptonic final states (via Coulomb force).**

# Quantitative description

Let us consider how such an effect is quantitatively described using an example that  $A'$  decays into a bottom quark pair at the threshold.

*Recipe ... The use of the potential non-relativistic (pNR) lagrangian:*

1. Integrating out long-range force fields from the Lagrangian.
2. NR-expand the (nearly) on-shell part of the bottom quark field.
3. Introducing auxiliary fields describing the two-body states of  $b\bar{b}$ .
4. Integrating out the NR-expanded bottom quark field.

$$\mathcal{L}_V^{(\text{pNR})} \simeq -\frac{1}{2}\vec{A}'(\square + m_{A'}^2)\vec{A}' + \int d^3r \bar{\chi}^\dagger(\vec{r}, x) \left( i\partial_{x^0} + \frac{\nabla_x^2}{4m_b} + \frac{\nabla_r^2}{m_b} - V(\vec{r}) \right) \chi(\vec{r}, x) + \sqrt{2/3} \epsilon \vec{A}' \left[ e^{2im_b x^0} \bar{\chi}^\dagger(\vec{0}, x) + e^{-2im_b x^0} \chi(\vec{0}, x) \right] + \dots$$

Here,  $\chi(r, x)$  is the field describing the two-body states of  $b\bar{b}$ .

5. Decomposing the two-body field in terms of the partial waves.

$$\mathcal{L}_V^{(\text{pNR})} \simeq -\frac{1}{2}\vec{A}'(\square + m_{A'}^2)\vec{A}' + \sum_i \bar{f}_i (i\not{D} - m_{f_i}) f_i - \sum_n \frac{1}{2} \bar{\varphi}_n (\square + m_{\varphi_n}^2) \varphi_n + e \sqrt{\frac{2m_b}{3\pi}} \left[ \epsilon \vec{A}' - ie \sum_i \int d^4y Q_i \mathcal{G}^{(\gamma)}(x-y) \bar{f}_i(y) \not{\gamma} f_i(y) \right] \cdot \sum_n R_{n00}(0) \varphi_n$$

**Mixing!**

**Radial wave fn.**

$\varphi_n$  is a partial wave component w/  $1^-$  quantum number ( $Y$  meson).

Note that no continuum states exist due to the color consignment.


# Mixing & Decays

When  $m_{\varphi_n}$  becomes very close to  $m_{A'}$ , the perturbative calculation of the width breaks down, and we have to diagonalize the mass matrix:

$$\begin{pmatrix} \vec{A}' \\ \vec{\varphi}_n \end{pmatrix} = \begin{pmatrix} \cos \vartheta_n & -\sin \vartheta_n \\ \sin \vartheta_n & \cos \vartheta_n \end{pmatrix} \begin{pmatrix} \vec{A}' \\ \vec{\varphi}_n \end{pmatrix}$$


Then the interaction of the mediator particle becomes

$$\mathcal{L}_V^{(\text{PNR})} \simeq -e\vec{A}' \cdot \sum_i Q_i \int d^4y \left[ \epsilon \cos \vartheta_n \delta(x-y) - ie \sin \vartheta_n \sqrt{\frac{2m_b}{3\pi}} R_{n00}(0) \mathcal{G}^{(\gamma)}(x-y) \right] \bar{f}_i(y) \vec{\gamma} f_i(y)$$



Originating in the  $A'$  int.

+



Originating in the  $Y$  int.

Other interactions originating in those of the  $Y$  mesons are

$Y \rightarrow l^- l^+$  (lepton pairs)

$Y \rightarrow qq$  (light hadrons)

$Y \rightarrow ggg, gg\gamma$ , (**3** gauge bosons)


$Y(nS) \rightarrow Y(mS) + X$  (with  $n > m$ )

[De-excitation process]

$Y \rightarrow B$  meson pairs ( $m_Y > 2m_B$ )



# Result & Summary



*We have quantitatively estimated **the threshold singularity effect** on the interaction strength (total decay width) of the vector mediator particle when its mass is close to those of **upsilon mesons**. We have given a formalism to quantitatively describe this effect based on **pNR** and found that the effect **could be sizable**. The same effect will appear for another mass and another type of mediator particle whenever it decays at a threshold with a final state feeling a long-range force.*