



Exploring Right Handed Neutrinos at ILC500

Work in progress

Jurina Nakajima, Daniel Jeans^A, Arindam Das^B, Keisuke Fujii^A,
Nobutaka Okada^C, Satomi Okada^C, Ryo Yonamine^A


SOKENDAI, KEK^A, Hokkaido Univ.^B, Alabama Univ.^C



S O K E N D A I

Introduction of ILC

International Linear Colider



- e+e- linear collider
- $\sqrt{s} = 250 \text{ GeV}$
(Extend to 500, 1000GeV)
- Polarized beams
 - e⁻: 80%, e⁺: 30%
 - “eLpR” = (e⁻ -80, e⁺ +30)
 - “eRpL” = (e⁻ +80, e⁺ -30)
- Higgs factory
- Measuring Higgs precisely

+

New physics search
beyond SM

Motivation and introduction

The Right Handed Neutrino (RHN) can address the following big questions

- Why does matter dominate anti-matter in our universe?
- Do quarks and leptons unify?
- Why is neutrino mass so small?


RHN is assumed to be

a Majorana particle ($N = \bar{N}$) gauge boson : Z'

minimal $U(1)_{B-L}$ model

$$G_{B-L} \equiv SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$$

→ RHN pair production
[arXiv\[1812.11931\]](https://arxiv.org/abs/1812.11931)



Benchmark points with $M_N = 100, 150, 200, 225$ GeV

Benchmark points


- $\text{Pol}(e^-, e^+) = (-0.8, +0.3), (+0.8, -0.3): \mathcal{L} = 1600 [\text{fb}^{-1}]$
- $\text{Pol}(e^-, e^+) = (-0.8, -0.3), (+0.8, +0.3): \mathcal{L} = 400 [\text{fb}^{-1}]$

M_N [GeV] RHN mass	$M_{Z'}$ [TeV] Z' mass	g_1' $U(1)_{B-L}$ coupling	$ V_{eN} ^2$ mixing angle	BR ($N \rightarrow eW$)	$\sigma(e_L^- e_R^+ \rightarrow NN)$ 100% polarization [fb]	Event # at ILC500 [4000fb $^{-1}$]
100	7	1	0,0009	0,44	0,71	1261
150	7	1	0,0009	0,33	0,45	229
200	7	1	0,0009	0,30	0,16	131
225	7	1	0,0009	0,29	0,052	18

- ▶ minimal $U(1)_{B-L}$ model
- ▶ ILC 500 with initial state radiation (ISR) and beamstrahlung (BS)


Current limits - Z' mass

SM like Z' coupling



ATLAS-TDR-LHCC2017-2018

RHN pair production crosssection at ILC250
for expected HL-LHC limits on $M_{Z'}/g'$




arXiv[1812.11931]

The heavier Z' mass less constrained by LHC


Analysis tool and backgrounds

ILC500

Signal process:



6f and 4f major background processes:



UFO model files

WHIZARD ver 2.8.5
Make Events



ILD Full Simulation
& (Geant4)
Reconstruction

miniDST
Events format

Cut conditions to select signal events


- Isolated e # = 2 && Isolated γ, μ # = 0
- Same sign isolated electrons
- Isolated electron energies $E_{\text{iso}} < 200$ [GeV]
- Isolated electron angles $|\cos\theta_{\text{isoel}}| < 0.95$
- IsolatedLepTagging(min) > 0.9
- Jet clustering with Durham $\log_{10}(y_{12}) > -1$
- $P_{\text{miss}} < 100$ [GeV]
&& ($P_{\text{miss}} < 40$ [GeV] II $|\cos\theta_{P_{\text{miss}}}| > 0.95$)

Signal efficiency $\sim 20\%$
Remaining backgrounds events
 ~ 150 (eLpR), 20 (eRpL)



Reconstruction methods

After removing isolated electrons force into 4 jets (Durham)



Jet pair 1 $\rightarrow M_{jj1}$, Jet pair 2 $\rightarrow M_{jj2}$

$$F_1 = (M_{jj1} - M_w)^2 + (M_{jj2} - M_w)^2$$

Best jet pair 1 + iso e $\rightarrow M_{jje1}$
Best jet pair 2 + iso e $\rightarrow M_{jje2}$
We expect for " $M_{jje1} = M_{jje2}$ "


$$F_2 = (M_{jje1} - M_{jje2})^2$$

Choose combination with minimum F_1, F_2

Signal mass cut

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- $\mathcal{L} = 1600 [\text{fb}^{-1}]$


For each M_N , mass window $M_N-10, M_N+15 [\text{GeV}]$



Assume background distribution is flat
20 (eLpR) background events remain in mass window

Signal mass cut

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- $\mathcal{L} = 1600 [\text{fb}^{-1}]$




Assume background distribution is flat

20 (eLpR) and 3 (eRpL) background events remain in mass window

Less backgrounds thanks to beam polarization
Reduce W contribution

Exclusion plot on cross-section



Exclude benchmark points and cross-sections up to 10x smaller

Conclusion:

Can use same sign lepton signature to set powerful limits on RHN at ILC!

Current Activity & Future Plan

- ❑ ILC250 case (on going)
→ Try to improve signal efficiency

- ❑ Same sign muons
→ Expect smaller backgrounds

Part1: RHN

Model : minimal $U(1)_{B-L}$

Gauged B-L extension of Standard Model (SM)

The unique anomaly free global symmetry in the SM

$$G_{B-L} \equiv SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$$

- ▶ Anomaly free requirement → **RHNs**
- ▶ **Seesaw mechanism** ← automatically included

Gauge boson : Z'

If B-L symmetry breaks spontaneously $\rightarrow Z'$ becomes **massive**



minimal $U(1)_{B-L}$ model : charge					
	SU(3) _C	SU(2) _L	U(1) _Y	U(1) _{B-L}	
RHN	N_R^i	1	1	0	-1
New Higgs field	Φ	1	1	0	2

i=1,2,3


[arXiv\[1812.11931\]](#)

Arindam Das, Nobuchika Okada, Satomi Okada, Digesh Raut

Signal




Destructive interference



Isolated e,γ,μ


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$

Number of isolated e




- Isolated e # = 2 && Isolated γ, μ = 0

Number of isolated γ




Number of isolated μ




Distribution of Isolated electron energy

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- Isolated e # = 2 && Isolated γ # = 0 && Isolated μ # = 0
- Isolated e is same sign ($e_1 \times e_2 = 1$)



Distribution of $\cos\theta_{\text{iso}}$


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- Isolated e # = 2 && Isolated γ # = 0 && Isolated μ # = 0
- Isolated e is same sign ($e_1 \times e_2 = 1$)



4 fermions semi leptonic processes in t-channel \rightarrow distributed in $|\cos\theta_{\text{iso}}| \sim 1$


Distribution of y_{12} (Durham)

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- Isolated $e \# = 2 \&\& \text{Isolated } \gamma \# = 0 \&\& \text{Isolated } \mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)



$\text{Cos}\theta_{\text{Pmiss}}$ vs Magnitude of missing momentum P_{miss}

- ILC 500 with ISR / BS



$P_{\text{miss}} < 100 \ \&\& (P_{\text{miss}} < 40 \ \text{II} \ |\text{cos}\theta_{\text{Pmiss}}| > 0.95)$

Cut flow (eRpL)

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$ $\mathcal{L} = 1600 [\text{fb}^{-1}]$

ILD work in progress


	Expected signal				Expected background					
	$M_N=100$	$M_N=150$	$M_N=200$	$M_N=225$	$eeqqqq$	$4f_{\text{singleW}}_{\text{semileptonic}}$	$4f_{\text{singleZee}}_{\text{semileptonic}}$	$6f_{\text{ttbar}}_{2\text{electrons}}$	$6f_{\text{ttbar}}_{1\text{electron}}$	$6f_{\text{ttbar}}_{0\text{electron}}$
No cut	558	394	143	45	3925	258648	612455	7100	56233	4894
$e_{\text{iso}} \# == 2 \& \& \gamma_{\text{iso}} \# == 0 \& \&$	420	343	126	40	1935	9426	249000	6142	1295	127
Same sign ($e_{\text{iso1}} \times e_{\text{iso2}} = 1$)	346	115	81	12	1231	7210	140176	3911	870	94
$E_{\text{iso}} < 200$ [GeV]	171	114	41	12	14	3741	3294	2	177	19
$-0.95 < \cos\theta_{\text{iso}} < 0.95$	158	103	37	11	3	1324	475	1	113	12
IsolatedLepTaugging _{min} > 0.9	96	91	32	10	0	198	101	0	15	1
$\log_{10}(y_{12}) > -1$	88	90	30	9	0	199	86	0	6	0
$P_{\text{miss}} < 100 \& \& (P_{\text{miss}} < 40 \text{ } \cos\theta_{\text{miss}} > 0.95)$	86	84	29	9	0	4	15	0	2	0

Results

	M_N [GeV]	# of Signal After cut	# of BG After cut	Signal Significance	σ_0 [fb] Initial benchmark	σ^{95} [fb] 95% exclusion limit	$\frac{\sigma^{95}}{\sigma_0}$	α^{95} [TeV ⁻⁴]
LR 80,30	100	53,64	20,12	6,25	0,55	0,12	0,21	3,83E-05
	150	52,73		6,18	0,36	0,076	0,21	2,96E-05
	200	18,30		2,95	0,14	0,086	0,61	7,57E-05
	225	5,51		1,18	0,046	0,085	1,8	2,21E-04
RL 80,30	100	66,75	3,24	7,98	0,71	0,065	0,092	1,69E-05
	150	63,41		7,77	0,45	0,043	0,097	1,35E-05
	200	21,23		4,29	0,16	0,047	0,29	3,57E-05
	225	6,077		1,99	0,052	0,052	1	1,21E-04

Exclusion plot on U(1)_{B-L} parameters

Translate to the U(1)_{B-L} model parameters



The benchmark points isn't excluded only at $M_N = 225 \text{ GeV}$

eLpR case

Cut flow (eLpR)



- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$ $\mathcal{L} = 1600 [\text{fb}^{-1}]$

ILD work in progress

	Signal Entries				Background Entries					
	$M_N=100$	$M_N=150$	$M_N=200$	$M_N=225$	$e\bar{e}qqqq$	$4f_{\text{singleW semileptonic}}$	$4f_{\text{singleZee semileptonic}}$	$6f_{\text{ttbar 2electrons}}$	$6f_{\text{ttbar 1electron}}$	$6f_{\text{ttbar 0electron}}$
No cut	554	394	143	45	11898	2825010	699475	16425	129283	11028
$e_{\text{iso}} \# == 2 \&\& \gamma_{\text{iso}} \# == 0 \&\&$	347	343	79	40	4721	90818	162774	9422	2271	201
Same sign ($e_{\text{iso1}} \times e_{\text{iso2}} = 1$)	176	115	39	12	39	46138	3800	8	439	25
$E_{\text{iso}} < 200 [\text{GeV}]$	175	114	39	12	39	41319	3557	8	439	25
$-0.95 < \cos\theta_{\text{isoe}} < 0.95$	156	103	36	11	13	17506	623	4	266	15
IsolatedLepTagging _{min} > 0.9	94	91	31	10	2	2632	128	1	50	0
$\log_{10}(y_{12}) > -1$	94	90	31	9	2	2632	128	1	50	0
$P_{\text{miss}} < 100 \&\& (P_{\text{miss}} < 40 \text{ II } \cos\theta_{\text{miss}} > 0.95)$	84	84	28	9	1	79	30	0	9	0

Electron Charge


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e^- \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$



We use only same sign samples $e_1 \times e_2 = 1$

Distribution of IsolatedLepTagging

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e \# = 2 \ \&\& \text{Isolated } \gamma \# = 0 \ \&\& \text{Isolated } \mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)




IsolatedLepTagging_{min} > 0.9

Isolated e,γ,μ


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$

Number of isolated e




- Isolated e # = 2 && Isolated γ, μ = 0

Number of isolated γ




Number of isolated μ




Distribution of Isolated electron energy

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated e # = 2 && Isolated γ # = 0 && Isolated μ # = 0
- Isolated e is same sign ($e_1 \times e_2 = 1$)



Distribution of $\cos\theta_{\text{iso}e}$


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e \# = 2 \ \&\& \text{ Isolated } \gamma \# = 0 \ \&\& \text{ Isolated } \mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)



4 fermions semi leptonic processes in t-channel \rightarrow distributed in $|\cos\theta_{\text{iso}e}| \sim 1$

Distribution of y_{12} (Durham)

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e \# = 2 \ \&\& \text{ Isolated } \gamma \# = 0 \ \&\& \text{ Isolated } \mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)



4f and 6f background information


Cross section — BG

- ILC 500 with ISR / BS

(100%,100%)	eeqqqq			4f_singleW _semileptonic	4f_singleZee _semileptonic
eLpR	eexyyx	xxxxee	yyyyee	4f_sw_sl	4f_sze_sl
1,64E+01	8,71E-02	1,45E-01	7,81E+03	1,96E+03	
3,64	4,62E-02	5,31E-02	2,28E+01	1,73E+03	
6,63	3,38E-02	2,20E-02	7,53E+02	1,78E+03	
6,61	3,30E-02	1,97E-02	7,50E+02	1,78E+03	

Cross section – BG


- ILC 500 with ISR / BS



Information associated
with $U(1)_{B-L}$ model


Current limits - Z' mass

SM like Z' coupling



ATLAS-TDR-LHCC2017-2018

HL-LHC prospects limit for $U(1)_{B-L}$ model




[arXiv\[1812.11931\]](#)

The heavier Z' mass less constrained by LHC

Current limits $|V_{eN}|^2$


$|V_{eN}|^2$: the “light-heavy” neutrino mixing matrix



<https://arxiv.org/pdf/1802.02965.pdf>

Current Limits and prospects - Z' mass,g1'

G1' :U(1)_{B-L} gauge coupling constant



$M_N [\text{GeV}]$	$M_{Z'} [\text{TeV}]$	$g1'$
100	7	1
200	7	1

By Arindam.D