


A new  
heavy-ion  
collision  
experiment  
at CERN: NA60+


Letter of Intent: [CERN-SPSC-2022-036 ; SPSC-I-259](#)  
(also [arXiv:2212.14452](#))

Website: <https://na60plus.ca.infn.it>





E. Scomparin (INFN Torino, Italy)  
G. Usai (Università and INFN Cagliari, Italy)


# A new heavy-ion experiment at CERN



**ALICE**: general purpose HI detector at LHC: study high  $T$ , zero  $\mu_B$  region (+ **ATLAS**, **CMS**, LHCb)



**NA60+**: high- $\mu_B$  studies of hard and electromagnetic probes of the Quark-Gluon Plasma at SPS energies



**NA61/SHINE**: (only) hadron detector at SPS: study intermediate  $T$ , finite  $\mu_B$  region


Complete access to QGP-related observables in a wide range of  $T$  and  $\mu_B$

# The NA60+ physics program

Several **new and unique measurements** in the region  $6 < \sqrt{s_{NN}} < 17$  GeV ( $20 < E_{lab} < 160$  AGeV)


## 1 Caloric curve of QGP

Measurement of temperature of thermal dimuons vs  $\sqrt{s_{NN}}$




## 2 Chiral symmetry restoration

$\rho$ - $a_1$  mixing in the dimuon channel




## 3 Charmonium melting in the QGP

Charmonium suppression vs  $\sqrt{s_{NN}}$  (dimuon decay channel)



## 4 QGP transport coefficients and charm hadronization

Hadronic decays of open HF mesons/baryons




# The NA60+ physics program


Several **new and unique measurements** in the region  $6 < \sqrt{s_{NN}} < 17$  GeV ( $20 < E_{lab} < 160$  AGeV)

## 1 Caloric curve of QGP

Measurements only at top SPS energy and at very low energy



HADES, Nature Phys. 15(2019) 1040  
NA60, EPJC 61(2009) 711




- Dilepton  $T_{slope}$  measurements  $\rightarrow$  (average) **temperature of the early stage of the system**
- SPS energy  $\rightarrow$  accurate information on the region close to the deconfinement transition temperature  $\rightarrow$  possible signal of a **1<sup>st</sup> order phase transition**


# The NA60+ physics program

Several **new and unique measurements** in the region  $6 < \sqrt{s_{NN}} < 17$  GeV ( $20 < E_{lab} < 160$  AGeV)

- Mixing of vector (V) and axial-vector (A) correlators  
→ **dilepton enhancement**  
for  $m_{\mu\mu} \sim 1-1.4$  GeV/ $c^2$



R. Rapp and H. van Hees, PLB753 (2016) 586



Chiral symmetry restoration

2

C. Jung et al.,  
PRD 95 (2017) 036020

- SPS vs LHC: low-energy measurement expected to be more sensitive to chiral restoration effects  
→ (Exponential) thermal dimuon yield from QGP becomes smaller  
→ Contribution from open charm becomes relatively negligible


No measurements available

# The NA60+ physics program

Several **new and unique measurements** in the region  $6 < \sqrt{s_{NN}} < 17$  GeV ( $20 < E_{lab} < 160$  AGeV)

No measurements below top SPS energy

3 Charmonium melting in the QGP



## J/ψ

- 30% suppression for central Pb-Pb events at top SPS energy  
→ Compatible with suppression of more weakly bound  $\chi_c$  and  $\psi(2S)$  states decaying to J/ψ

## ψ(2S)

- Strong(er) suppression already in peripheral Pb-Pb collisions
- Energy scan towards low SPS energy  
→ Detect **suppression threshold** and correlate with T via thermal dimuons

NA50, PLB 477 (2000) 28  
NA50, EPJC49 (2007) 559


# The NA60+ physics program

Several **new and unique measurements** in the region  $6 < \sqrt{s_{NN}} < 17$  GeV ( $20 < E_{lab} < 160$  AGeV)

□ Charm production in **proton-nucleus**

→ Sensitive to **nPDFs**

→  $Q^2 \sim 10\text{--}40$  GeV<sup>2</sup> and  $0.1 < x_{Bj} < 0.3$  ( $p_T < 3$  GeV/c)  
(from anti-shadowing to EMC region)



No measurements at SPS energy


□ D-meson  $p_T$ -dep. suppression and azimuthal anisotropy

→ Time spent in QGP and hadronic phase varies with  $\sqrt{s_{NN}}$ : constrain the charm diffusion coefficient

→ **Do charm quarks thermalize** in a short-lived QGP?

□  $D_s^+, \Lambda_c$

→ Hadronization studies (quark recombination)




**QGP transport coefficients and charm hadronization**

4

ALICE, PLB 827 (2022) 136986

# The NA60+ detector



Inspired by the **former NA60** detector (2002-2004)

Measurement of **(di)muon** production and hadronic decays of **strange** and **charm** hadrons

SPS **energy scan**: vary z-position of the muon spectrometer and thickness of hadron absorber




# The NA60+ vertex region



**MEP48 dipole** magnet  
Field 1.5 T over a 400mm gap

Stored at **CERN**,  
needs refurbishment



# The NA60+ vertex telescope R&D

Sensor based on 25 mm long units, replicated several times through stitching  
→ up to 15cm length for NA60+




R&D in progress  
Common development  
**ALICE ↔ NA60+**  
**(same timeline!)**

State-of-the-art imaging technology  
TowerJazz 65 nm


Sensor thickness:  
few tens of microns of silicon  
→ material budget **<0.1% X<sub>0</sub>**

Spatial resolution **≤ 5 μm**  
Cooling studies (NA60+ geometry)  
→ **airflow+water**

- ❑ Engineering run for a fully functional prototype
- ❑ Possibility of a second run if optimizations needed





12 inch Si wafer




# The NA60+ muon spectrometer


6 tracking stations, with a modular structure




12x2 modules



36x2 modules



84x2 modules




- Modest rates (FLUKA) already in the upstream stations, thanks to the thick absorber (235 cm BeO +C)
- For a  $10^6 \text{ s}^{-1}$  beam  
→ charged particle rate  $\sim 2 \text{ kHz/cm}^2$


Can be matched by **GEM** or **MWPC** detectors

# The NA60+ muon spectrometer R&D


- First prototype of a **MWPC module built and tested** at Weizmann institute  
→ to be tested on a hadron beam at CERN in spring 2023



Prototype




Cosmic testbench





Weizmann (MWPC) and SBU (GEM) facilities have the technical possibility of **managing the full production** of the detector modules for the NA60+ muon spectrometer

- Ongoing discussions on the final set-up of the spectrometer, various possible solutions, as
  - GEM technology for upstream stations (MS0-MS1)
  - MWPC technology for downstream stations (MS2-MS5)



# The NA60+ toroid



## Warm magnet

Eight sectors,  
12 turns per coil

Conductor has a square  
copper section with a  
circular cooling  
channel in the centre




Prototype (1:5 scale)




- Measurements of resistance, inductance, cooling performance and magnetic field were carried out
- B measurement  
→ agreement with simulations by 3%

**Support and participation of CERN**  
in the design of the final toroid  
is very important



Operating Current [kA]	16.6
Amp-turns [kA]	199
Combined inductance [mH]	9.5
Resistivity Al 1100 @RT [ $\mu\Omega\cdot\text{cm}$ ]	2.67
Length Conductor [m]	800
Total resistance [m $\Omega$ ]	10.4
Dissipated power [MW]	2.8

# Uniqueness of NA60+ program






- The NA60+ physics program needs a large integrated luminosity  
→ Measurement of **rare QGP probes**
- Such a luminosity can be obtained with **Pb-Pb interaction rates  $>10^5$  Hz**, reachable with a  $\sim 10^6$  s<sup>-1</sup> beam intensity in a fixed-target environment
- In the SPS energy range, there **are no other existing/foreseen facilities/experiments** that can approach this kind of performance

- **Complementarity** with experiments accessing
  - different (hadronic) observables in the same energy range (STAR BES, NICA, NA61)
  - similar observables in a lower energy range (CBM at FAIR)

# NA60+: where

- ❑ Thorough studies carried out in 2020/2021 thanks to **PBC support**, with the decisive help of the **CERN-BE-EA group**  
→ integration feasible in the PPE138 area on the H8 beam



Need rail installation (muon spectrometer shifting) and a possible floor excavation due to the current vertical position of the beam line

NA60+, NIM A1047 (2023) 167887



# NA60+: beam studies R&D




- ❑ A **high-intensity Pb beam** ( $\sim 10^6/s$ ) is needed, from 20-30 A GeV to 160 A GeV
- ❑ Beam optics studies carried out to provide sub-mm beam all over the energy range

Goal

Parameter in zone 138	160 GeV/c	30 GeV/c
$\sigma_x$ (mm)	0.19	0.33
$\sigma_y$ (mm)	0.19	0.36
Transmission from T4 (%)	32.43	23.5



N.B.: Vertex spectrometer central hole,  $\varnothing \sim 0.6 \text{ cm}$



A first **test beam in PPE138** was carried out in November 2022, using a telescope of pixel sensors for a precise measurement

Result already promising, further tests needed  
 → Lower beam energy  
 → Higher beam intensity (now  $\sim 10^4 \text{ s}^{-1}$ )

**Pb beam request** submitted for fall 2023

NA60+, NIM A1047 (2023) 167887



# NA60+: beam requests


- Our plan is to run each **~ 1month/year with Pb ions** at a different energy, using a **~10<sup>6</sup> s<sup>-1</sup> beam**
  - Start at top energy, to have a calibration point for observables already studied at that energy
  - At 20 A GeV two months of data taking can be necessary to fulfil the physics program
  - The order of the beam energies is tentative and could be adjusted following the results

	Year 1	Year 2	Year 3	Year 4-5	Year 6	Year 7
Beam energy (A GeV)	160	40	120	20 (30)	80	60
Momentum per charge (GeV/c/Z)	406	101	304	50.7 (76.1)	203	152
Pb ions on target	$\sim 10^{12}$ per energy ( $\sim 30$ days)					
protons on target	$5 - 6 \cdot 10^{13}$ per energy ( $\sim 22$ days)					


- Corresponding periods with **proton beams** at the same energy are also needed
  - Reference for Pb-Pb results
  - Specific studies with p-A collisions
- Integrated luminosity per N-N collision similar for p-A and Pb-Pb
- Beam intensity  $\sim 8 \times 10^8$ /spill, 3000 spills/day (preliminary estimate)

Using a high-intensity beam in the EHN1 surface zone poses non-negligible **radioprotection issues**  
→ Thorough studies carried out by the **CERN-HSE** group

## Top view




Prompt ambient dose, residual ambient dose, air activation and accidental beam loss scenarios were studied



A **massive shielding** around the absorber region, where the beam will be dumped, has been designed

# Physics performance: dimuons





Track matching

Vertex tracking  
(dipole field)


Muon identification in the  
spectrometer (toroidal field)

Track matching: measure  
muon kinematics  
before multiple scattering and  
energy loss



# Physics performance: thermal radiation

1



- Thermal radiation yield
  - Dominated by  $\rho$  contribution at low mass
  - Accessible up to  $M=2.5-3 \text{ GeV}/c^2$
- Drell-Yan contribution
  - to be also estimated via p-A measurements
- Open charm
  - Negligible dimuon source

~1-3% uncertainty on the evaluation of  $T_{\text{slope}}$

Energy (GeV)	Thermal pairs	$T_{\text{slope}}$
6.3	$3.52 \cdot 10^6$	$166 \pm 4.7 \pm 1$
8.8	$3.56 \cdot 10^6$	$169 \pm 4.4 \pm 1$
17.3	$9.70 \cdot 10^6$	$182 \pm 1.8 \pm 1$

(0-5% central Pb-Pb collisions)


2 months →  
1 month →

□ **Elliptic flow** measurement also feasible

Accurate mapping of the region where  $T_{pc}$  is reached  
 → Strong **sensitivity** to possible flattening due to 1<sup>st</sup> order transition

# Physics performance: chiral symmetry



- Detect **modification of continuum** in  $1 < m_{\mu\mu} < 1.4$  GeV, related to chiral symmetry restoration



- Comparison of spectra ( $\sqrt{s_{NN}} = 8.8$  GeV), based on the assumption of no chiral mixing, with expectation of full chiral mixing
- Statistical and systematic uncertainty provide a very good sensitivity to an **increase of the yield due to chiral mixing of  $\sim 20\text{--}30\%$**

# Physics performance: charmonium

3



- 7.5mm Pb target and 1 month data taking  
→  $L_{\text{int}} = 24 \text{ nb}^{-1}$

Can aim at


- $\sim \mathcal{O}(10^4)$   $J/\psi$  at 50 GeV
- $\sim \mathcal{O}(10^5)$   $J/\psi$  at 158 GeV

- Detection of **onset of anomalous suppression** effects down to low SPS energy
- **p-A data taking mandatory** (few weeks/year), to calibrate CNM effects
- $\psi(2S)$  also within reach, down to  $E = 100\text{-}120$  A GeV


- NA60+ is also ideally placed to look for signals of **intrinsic charm** in p-A collisions, which are pushed much closer to midrapidity wrt collider energies

# Physics performance: open charm


- Combine tracks in the vertex spectrometer only, apply topological cuts
- $10^{11}$  minimum bias Pb-Pb collisions:  $>3 \cdot 10^6$  reconstructed  $D^0$  in central Pb-Pb at  $\sqrt{s_{NN}}=17.3$  GeV
  - 2-3 orders of magnitude larger than forthcoming **NA61 results**
  - **$D^0$**  accessible also at lower collision energies with statistical precision at the percent level
  - Measurement of  **$D_s$  and  $\Lambda_c$  yield feasible** with statistical precision of few percent



$D^0 \rightarrow K\pi$



$D_s^+ \rightarrow \Phi\pi \rightarrow KK\pi$



$\Lambda_c^+ \rightarrow pK\pi$

Similar technique allows measurements of **hyperons and hypernuclei**

# Collaboration institutes

## Appendix: NA60+ Collaboration

C. Ahdida<sup>1</sup>, G. Alocco<sup>2,3</sup>, F. Antinori<sup>4</sup>, M. Arba<sup>3</sup>, M. Aresti<sup>2,3</sup>, R. Arnaldi<sup>5</sup>, A. Baratto Roldan<sup>1</sup>, S. Beolè<sup>6,5</sup>, A. Beraudo<sup>5</sup>, J. Bernhard<sup>1</sup>, L. Bianchi<sup>6,5</sup>, M. Borysova<sup>7,8</sup>, S. Bressler<sup>7</sup>, S. Bufalino<sup>9,5</sup>, E. Casula<sup>2,3</sup>, C. Cicalò<sup>3</sup>, S. Coli<sup>5</sup>, P. Cortese<sup>10,5</sup>, A. Dainese<sup>4</sup>, H. Danielsson<sup>1</sup>, A. De Falco<sup>2,3</sup>, K. Dehmelt<sup>11</sup>, A. Drees<sup>11</sup>, A. Ferretti<sup>6,5</sup>, F. Fionda<sup>2,3</sup>, M. Gagliardi<sup>6,5</sup>, A. Gerbershagen<sup>12</sup>, F. Geurts<sup>13</sup>, V. Greco<sup>14,15</sup>, W. Li<sup>13</sup>, M.P. Lombardo<sup>16</sup>, D. Marras<sup>3</sup>, M. Masera<sup>6,5</sup>, A. Masoni<sup>3</sup>, L. Micheletti<sup>1</sup>, L. Mirasola<sup>2,3</sup>, F. Mazzaschi<sup>1,6</sup>, M. Mentink<sup>1</sup>, P. Mereu<sup>5</sup>, A. Milov<sup>7</sup>, A. Mulliri<sup>2,3</sup>, L. Musa<sup>1</sup>, C. Oppedisano<sup>5</sup>, B. Paul<sup>2,3</sup>, M. Pennisi<sup>6,5</sup>, S. Plumari<sup>14</sup>, F. Prino<sup>5</sup>, M. Puccio<sup>1</sup>, C. Puggioni<sup>3</sup>, R. Rapp<sup>17</sup>, I. Ravinovich<sup>7</sup>, A. Rossi<sup>4</sup>, V. Sarritzu<sup>2,3</sup>, B. Schmidt<sup>1</sup>, E. Scomparin<sup>5</sup>, S. Siddhanta<sup>3</sup>, R. Shahoyan<sup>1</sup>, M. Tuveri<sup>3</sup>, A. Uras<sup>18</sup>, G. Usai<sup>2,3</sup>, H. Vincke<sup>1</sup>, I. Vorobyev<sup>1</sup>

1 .European Organization for Nuclear Research (CERN), Geneva, Switzerland

2 .Dipartimento di Fisica dell'Università di Cagliari, Cagliari, Italy

3 .INFN, Sezione di Cagliari, Cagliari, Italy

4 .INFN, Sezione di Padova, Padova, Italy

5 .INFN, Sezione di Torino, Turin, Italy

6 .Dipartimento di Fisica dell'Università di Torino, Turin, Italy

7 .Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot, Israel

8 .Kyiv Institute for Nuclear Research (KINR), Natl. Acad. of Sci. of Ukraine (NASU)

9 .Dipartimento DISAT del Politecnico di Torino, Turin, Italy

10 .Dipartimento di Scienze e Innovazione Tecnologica dell'Università del Piemonte Orientale, Alessandria, Italy

11 .Department of Physics and Astronomy, Stony Brook University, SUNY, Stony Brook, New York, USA

12 .Department of Radiation Oncology, University of Groningen, Groningen, The Netherlands

13 .Department of Physics and Astronomy, Rice University, Houston, Texas, USA

14 .Dipartimento di Fisica e Astronomia dell'Università di Catania, Catania, Italy

15 .INFN, Laboratori Nazionali del Sud, Catania, Italy

16 .INFN, Laboratori Nazionali di Frascati, Frascati, Italy

17 .Cyclotron Institute and Department of Physics and Astronomy, Texas A&M University, College Station, Texas, USA

18 .Institut de Physique des 2 Infinis de Lyon, Université de Lyon, CNRS/IN2P3, Lyon, France

- The LoI was signed by 62 physicists/engineers/technicians representing institutions in
  - **Italy** (Cagliari, Padova, Torino)
  - **Israel** (Weizmann)
  - **USA** (StonyBrook, Rice)
  - **France** (Lyon)
  - and **CERN**
- Support also from prominent members of the QGP theory community
- **Funding for the R&D phase** since 2020 allowed us to complete the LoI preparation
- Contacts ongoing to **strengthen the Collaboration** on specific items and reach critical manpower level
- **Already available** know-how and facilities for Si detector developments and gas detector construction  
→ may just need relatively small adjustments



# Cost estimates

- ❑ Final definition of the set-up details still in progress
- ❑ Estimate of costs related to data acquisition, storage and computing is still in progress
- ❑ Current evaluation subject to oscillation in the cost of raw materials, electronic, etc.
- ❑ Assume 1 Euro ~ 1 CHF ~ 1 US\$

## Toroid

Estimated cost (MCHF)	
Copper Conductor	0.6
Manufacturing of coils	1.7
Power converter (confirmation ~1/8)	0.8
Mechanical structure	0.4
Cooling system	0.3
<b>TOTAL</b>	<b>3.8</b>

**PRELIMINARY**

Sub-system	Estimated cost (MCHF)
Vertex spectrometer	2.5 – 3.1
Muon spectrometer	2.7 – 4.0
Toroidal magnet	3.8
RP monitors, Shielding	1.5
<b>Total</b>	<b>10.5 – 12.4</b>

**Table 17:** Estimated costs of the various NA60+ subsystems.

	kCHF
Engineering runs	600-1200
Wafer post-processing	300
FPC and wire bonding	200
Mechanical support	200
Cables, patch panels	300
Readout and power distribution	900
<b>TOTAL</b>	<b>2500-3100</b>

## MAPS

## Muons

**All MWPCs**

	kCHF
Detectors	500
FEE	1000
HV system	150
Mechanical support	750
Gas system	300
<b>TOTAL</b>	<b>2700</b>

**MSO/1 GEMS**


	GEM: kCHF
Detectors	530
Readout electronics	790
HV system	20
Mechanical support	50
Gas system	50
<b>TOTAL</b>	<b>1,440</b>

# Timeline

- ❑ Project followed by PBC since 2016
- ❑ EoI in 2019
- ❑ LoI in 2022

Our current plan is to have the experiment on the floor by the end of LS3 → **2029**

- ❑ Possible roadmap
  - ❑ Technical proposal: **2024-2025**
  - ❑ Construction and installation: **2026-2028**



# To do list towards technical proposal

## Muon spectrometer

- Beam tests on prototypes
- Finalize set-up and resolution (strip size) for MWPC and GEM detectors
- Define/design read-out electronics

## Toroidal magnet

- Design of the full-scale magnet, based on expertise gained with prototype

## Vertex spectrometer

- Continue R&D on MAPS development. N.B.: **same timeline as ALICE** for the final detector!
- Test first prototype of stitched MAPS detector
- Finalize test set-up with NA60+ geometry (dummy sensors), perform mechanics and cooling tests

## Simulation, reconstruction, DAQ

- From fast simulation/reconstruction to final framework
- Define DAQ framework for the estimated trigger rate

Based on this to do list, we estimate a 1.5-2.5 yrs timeline for the submission of TP

# Summary and final considerations


- ❑ A new heavy-ion experiment at CERN can **address several important questions** that cannot be answered with other facilities/experiments
- ❑ We have submitted a **LoI** for a new experiment that couples state-of-the-art (MAPS) and well known (MWPC, GEM) detection techniques
- ❑ The proponents have a **solid leadership in QGP studies** with hard and e.m. probes, that dates back to the first round of SPS experiments and continues until today at RHIC and LHC
- ❑ Strong **expertise in silicon and gas detector** construction and characterization (ALICE, ATLAS, PHENIX,...)
- ❑ Once R&D studies and detector set-up are finalized, and the Collaboration is strengthened, it is our intention to **submit a proposal** to SPSC (2024-2025)
- ❑ In our current timeline first data taking could occur in **2029**, after LS3

# Backup

# Trigger and DAQ

# Data acquisition, processing, computing (1)

- ❑ **Data rate** dominated by the vertex telescope, for the assumed  $10^6$  ions/s Pb beam intensity,
  - ~ **3.3 GB/s** data rate
  - ~ 3.3 PB of data collected per year
- ❑  **$\delta$ -ray** production from non-interacting Pb ions (85% of the incident beam) significantly contribute to the data rate
- ❑ Consider to acquire data **triggered** by a fast scintillator close to the interaction region
  - increase purity at the price of discarding peripheral Pb-Pb events



selection, %	trigger rate, kHz	purity, %	hits readout per incoming ion	hits readout per trigger	readout rate, GB/s
50	100	80	300	2960	0.94
80	365	35	675	1541	2.1
100	1000	16	1030	1030	3.3

↑  
Centrality selected

# Data acquisition, processing, computing (2)

## ❑ **Offline data reconstruction**

❑ → Use a modified version of the Cellular Automaton track finder developed for the ALICE ITS

❑ **Data decoding and cluster-finding** require ~240 (~450) CPU seconds for 50% (80%) efficiency triggering scenarios, for  $10^6$  incoming ions ← preliminary!



❑ Corresponding **track finding time** ~ 4200 CPU seconds (assume Intel i7-8700K @ 3.7 GHz processor)

❑ Data collected per heavy-ion run can be **fully processed in 2–3 months** by a farm of ~ 100 modern multicore processors or equivalent GRID jobs






# Strangeness and hypernuclei

# Strangeness measurements: hyperons



- ❑ Hyperon decays simulated with EVtGen, decay products propagated in the VT using the fast simulation of NA60+
- ❑ Background from hadron production → **NA49 results**

- ❑ Channels studied



and charge conjugated




- ❑ **Topological selections** applied
- ❑ **BDT employed to enhance the significance of the signal**

- ❑ Among the variables:

- ❑ Product of the impact parameter of decay tracks,
- ❑ Distance of closest approach between the decay tracks
- ❑ Decay length and the cosine of the pointing angle

- ❑ Also  $\phi \rightarrow KK$  and  $K_S \rightarrow \pi\pi$  were studied



# Physics performance: strangeness and hypernuclei



□ **Topological selections with BDT** employed to **enhance the significance** of the signal

- Among the variables:
  - Product of the impact parameter of decay tracks
  - Distance of closest approach between the decay tracks
  - Decay length and the cosine of the pointing angle

□ Also  $\phi \rightarrow KK$  and  $K_S \rightarrow \pi\pi$  have been studied




Low energy HI collisions  
 → **high baryon density** favours the production of hypernuclear clusters

Separation of heavily ionising particles from ordinary hadrons  
 → **size of the clusters** associated with the track

# Muon spectrometer

# The NA60+ muon spectrometer

Low-energy set-up



□ (At least) two configurations of the muon spectrometer are foreseen

□ **Low-energy set-up**

→ Thinner absorber


→ Smaller distance from target

□ **High-energy set-up**

→ Thicker absorber


→ Larger distance from target

High-energy set-up




Keep maximum acceptance around  $y \sim y_{CM}$



# MWPC prototype tests




- ❑ Wire pitch: 3 mm
- ❑ Distance wire to cathode: 3 mm
- ❑ 1 mm strip pitch
- ❑ 2 cathodes with strips running in two different directions  
→ Small angle stereo readout
- ❑ Readout electronics cards with VMM3a ASIC (128 ch each)



Detector tomography



Trigger and MWPC signals



# Vertex spectrometer

# Ongoing R&D on vertex spectrometer



## □ Detector

- Characterization of small-scale structures
- Submission of first large area MAPS with the stitching technique (MOSS)
- Development of test system for large area MAPS

## □ Mechanics

- Positioning and gluing tests of (dummy) sensors on carbon foam/fiber supports with optical bench



- Cooling calculations  
→ Mix air flow + water flow





# Toroid


# The NA60+ toroid



## Warm magnet


Eight sectors,  
12 turns per coil

Conductor has a square  
copper section with a  
circular cooling  
channel in the centre



Complex arrangement of the coils  
close to the beam axis to reduce the  
'dead zone' at forward  $y$

Ongoing discussions on strategy for  
**reducing the dissipated power**  
( $< 2$  month/yr, pulsed operation,...)



Operating Current [kA]	16.6
Amp-turns [kA]	199
Combined inductance [mH]	9.5
Resistivity Al 1100 @RT [ $\mu\Omega\cdot\text{cm}$ ]	2.67
Length Conductor [m]	800
Total resistance [m $\Omega$ ]	10.4
Dissipated power [MW]	2.8


# The NA60+ toroid R&D



- A **prototype (1:5 scale)** was built and tested in 2020-2021 by the **CERN-EP-DT group**, to check calculations and investigate mechanical solutions, in view of the final object

→ **works correctly and as expected**

- Measurements of resistance, inductance, cooling performance and magnetic field were carried out




- B measurement  
→ agreement with simulations by 3%

**Support and participation of CERN** in the design of the final toroid is very important

# NA60+ vs others

# NA60+ vs NA60



Some important improvements:

## Physics program extended to lower energy

→ Fundamental to explore rare probes in the high- $\mu_B$  region

## Larger angular acceptance

→ cope with lab rapidity shift when varying energy down to low SPS energy

## Access new observables (open charm etc.)


NA60: (di)muon trigger  $\sim 5$  kHz

NA60+: MB trigger ( $>100$  kHz)

## State-of-the art detectors

Pixel size: from  $50 \times 425 \mu\text{m}^2$  (NA60) to  $30 \times 30 \mu\text{m}^2$  (NA60+), thinner sensors (from 2% to 0.1%  $X_0$ )

→ Improved resolution and signal over background  
from 21 to 8 MeV at the  $\omega$  mass  
from 70 to 30 MeV at the  $J/\psi$  mass



# Uniqueness of NA60+ program

## NA60+ vs NA61

### NA61

Measurement of hadron production properties for


- Neutrino beams
- Cosmic ray experiments
- Strong interaction

### NA60+

Measurement of rare probes in HI collisions

- Dileptons
- Quarkonium
- Open heavy flavour(\*)
- Strangeness and hypernuclei

(\*) Also part of the NA61 program, but with 2-3 orders of magnitude smaller statistics



□ **Complementarity** with experiments accessing


- different (hadronic) observables in the same energy range (STAR BES, NICA, NA61)
- similar observables in a lower energy range (CBM at FAIR)

# Open charm NA60+ vs NA61

NA60+

NA61

Year	Beam	#days	#events	\$(D^0 + \bar{D}^0)\$	\$(D^+ + D^-)\$
2022	Pb at 150A GeV/c	42	250M	38k	23k
2023	Pb at 150A GeV/c	42	250M	38k	23k
2024	Pb at 40A GeV/c	42	250M	3.6k	2.1k






N.B.: different assumptions for open charm cross section

# Integration, radioprotection, beam



# NA60+: where


- ❑ Thorough studies carried out in 2020/2021 thanks to **PBC support**, with the decisive help of the **CERN-BE-EA group**  
→ integration feasible in the PPE138 area on the H8 beam



Need rail installation (muon spectrometer shifting) and a possible floor excavation due to the current vertical position of the beam line

**High-energy**  
setup

Using a high-intensity beam in the EHN1 surface zone poses non-negligible **radioprotection issues**  
→ Thorough studies carried out by the **CERN-HSE** group



Prompt ambient dose, residual ambient dose, air activation and accidental beam loss scenarios were studied



A **massive shielding** around the absorber region, where the beam will be dumped, has been designed

# First test beam in the H8 experiment location





**Focused optics**

Max beam intensity  
 $\sim 2 \cdot 10^5$  /spill

**Microcollimator**



150  
 AGeV



# Collaboration institute-wise

# Collaboration members

**CERN:** C. Ahdida<sup>HSE-RP</sup>, A. Baratto Roldan<sup>BE-EA-LE</sup>, J. Bernhard<sup>BE-EA-LE</sup>, H. Danielsson<sup>EP-DT-EF</sup>, A. Gerbershagen\*, M. Mentink<sup>TE-MPE-PE</sup>, L. Musa<sup>EP-AIO</sup>, M. Puccio<sup>EP-AIP-PAP</sup>, B. Schmidt<sup>EP-DT</sup>, R. Shahoyan<sup>EP-AIP-SDS</sup>, H. Vincke<sup>HSE-RP</sup>, I. Vorobyev<sup>EP-AIP-PAP</sup> (\*)now at Groningen

**Cagliari Univ. and INFN:** G. Alocco, M. Arba, M. Aresti, E. Casula, C. Cicalo, A. De Falco, F. Fionda, D. Marras, A. Masoni, L. Mirasola, A. Mulliri, B. Paul, C. Puggioni, V. Sarritzu, S. Siddhanta, M. Tuveri, G. Usai

**Padova INFN:** F. Antinori, A. Dainese, A. Rossi

**Torino Univ. and INFN:** R. Arnaldi, S. Beole, L. Bianchi, S. Bufalino, S. Coli, P. Cortese, A. Ferretti, M. Gagliardi, M. Maserà, L. Micheletti, F. Mazzaschi, P. Mereu, C. Oppedisano, M. Pennisi, F. Prino, E. Scomparin

**Weizmann Inst.:** M. Borysova, S. Bressler, A. Milov, I. Ravinovich

**Stony Brook Univ.:** A. Drees, K. Dehmelt




**Rice Univ.:** F. Geurts, W. Li

**IN2P3 Lyon:** A. Uras

**Theorists:** A. Beraudo (Torino), V. Greco (Catania), M.P. Lombardo (Firenze), S. Plumari (Catania), R. Rapp (Texas A&M)

# Charmonia

# Physics performance: charmonium




- 7.5mm Pb target and 1 month data taking  
 $\rightarrow L_{\text{int}} = 24 \text{ nb}^{-1}$   
 Can aim at
  - $\sim \mathcal{O}(10^4)$   $J/\psi$  at 50 GeV
  - $\sim \mathcal{O}(10^5)$   $J/\psi$  at 158 GeV

- Allows detection of **onset of anomalous suppression** effects down to low SPS energy
- **p-A data taking mandatory** (few weeks/year), to calibrate CNM effects
- $\psi(2S)$  also within reach, down to  $E = 100\text{-}120 \text{ A GeV}$

- NA60+ is also ideally placed to look for signals of **intrinsic charm** in p-A collisions, which are pushed much closer to midrapidity wrt collider energies

# J/ψ in Pb-Pb collisions at (various) SPS energies



**Quarkonium production  
not studied  
below top SPS energies!**





Perform an energy scan in  
 **$E_{\text{lab}} = 20 - 158 \text{ GeV}$**

- Decreasing  $\sqrt{s}$ :
  - **Onset of  $\chi_c$  and  $\psi(2S)$  melting**  
→ to be correlated to T measurement via thermal dimuons
  - **Stronger CNM effects**  
→ to be accounted for with pA data taking at the same  $\sqrt{s}$

- With  $\sim 10^{12}$  incident Pb on a 7.5mm Pb target (1 month of data taking) →  $L_{\text{int}} \sim 24 \text{ nb}^{-1}$  NA60+ can aim at
  - **$\sim O(10^4)$  J/ψ at 50 GeV**
  - **$\sim O(10^5)$  J/ψ at 158 GeV**
- N.B.: a factor 3 overall suppression (CNM + QGP) is assumed in these estimates



# NA60+, $R_{AA}$ estimate




- Based on
  - $10^{12}$  incident Pb
  - pA reference:  
 $5 \cdot 10^{13}$  incident p
- Assume only CNM effects for  $N_{\text{part}} < 50$  and 20% extra suppression in Pb-Pb for  $N_{\text{part}} > 50$

→ Precise evaluation of anomalous suppression within reach even at low energy

# Low- $\sqrt{s}$ J/ $\psi$ : studying intrinsic charm

- Intrinsic charm component of the hadron wavefunction  $|uudc\bar{c}\rangle$
- Leads to **enhanced charm production** in the forward region
- Hints from several experiments, but **no conclusive results**
- At colliders, forward  $x_F$  pushed to very high rapidity, difficult to measure  
→ fixed-target configurations more appropriate




Assumed intrinsic charm content varied between 0.1% and 1%

R. Vogt, PRC 103, 035204 (2021)  
R. Vogt, arXiv:2207.04347


# Low- $\sqrt{s}$ J/ $\psi$ : studying intrinsic charm

p-Pb collisions

- EPPS16 shadowing
- $\sigma_{\text{abs}} = 9, 10, 11$  mb at  $E_{\text{lab}} = 120, 80, 40$  GeV
- $P_{\text{ic}}^0$  varied between 0.1 and 1%



(w/o intrinsic charm)



□  $R_{\text{pPb}}$  shape is dominated by intrinsic charm, already with  $P_{\text{ic}}=0.1\%$

# Charmonia: high vs low $\sqrt{s}$

## Collider (LHC)

Hot matter effects: regeneration counterbalances (overcomes) suppression

### Initial state effects:




shadowing  
 $x \sim 10^{-5}$  ( $y \sim 3$ ),  
 $x \sim 10^{-3}$  ( $y = 0$ ),  
 $x \sim 10^{-2}$  ( $y \sim -3$ )

### (Final state) CNM effects:

negligible, extremely short crossing time

$$\tau = L/(\beta_z \gamma) \sim 7 \cdot 10^{-5} \text{ fm/c } (y \sim 3)$$

$$\tau = L/(\beta_z \gamma) \sim 4 \cdot 10^{-2} \text{ fm/c } (y \sim -3)$$



## Fixed target (SPS)


Hot matter effects: suppression effects (if existing) dominate

Initial state effects: moderate anti-shadowing  
 $x \sim 10^{-1}$  ( $y = 0$ )

(Final state) CNM effects: break-up in nuclear matter can be sizeable

$$\tau = L/(\beta_z \gamma) \sim 0.5 \text{ fm/c } (y = 0)$$

# J/ $\psi$ suppression: Pb-Pb at top SPS energy




- Contrary to open charm, accurate studies were performed at  $\sqrt{s}=17.3$  GeV (NA50, NA60)
- J/ $\psi$  yields normalized to Drell-Yan reference
- QGP-induced suppression evaluated with respect to a CNM reference obtained with systematic p-A studies
- **$\sim 30\text{-}40\%$  anomalous suppression effect** possibly due to disappearance of feed-down from  $\chi_c$  and  $\psi(2S)$

# CNM effects are (very) large

- ❑ Shadowing effects are moderate
- ❑ Dominated by nuclear absorption  
→ ~30% effect in p-Pb at  $\sqrt{s_{NN}} = 17$  GeV


## ❑ Strong $\sqrt{s}$ -dependence

→ CNM may become the dominant effect at low energy



Lourenco, Vogt, Woehri, JHEP 0902:014,2009

NA60, PLB 706 (2012) 263



$L$ : thickness of nuclear matter crossed by the cc pair  
(evaluated with Glauber model)




# Prospects for $\psi(2S)$ measurements at low $\sqrt{s}$

Good charmonium resolution ( $\sim 30$  MeV for the  $J/\psi$ ) will help  $\psi(2S)$  measurements

Expectations based on

- 30 days PbPb,  $I_{\text{beam}} = 1\text{e}7$  ions/spill
- 15 days pA,  $I_{\text{beam}} = 8\text{e}8$  p/spill


(assuming stronger suppression for  $\psi(2S)$  than  $J/\psi$ )




□  $\psi(2S)/\psi$  measurement looks feasible down to  $E_{\text{lab}} = 120$  GeV

□ Lower  $E_{\text{lab}}$  would require larger beam intensities/longer running times

# Existing open charm results at SPS energy



- Match track(s) in a muon spectrometer to tracks in a vertex spectrometer
- Excellent resolution on the muon kinematics
- Separate prompt (DY+thermal) from nonprompt sources (open charm)




- Analysis of open charm contribution (semileptonic decays of charm hadron pairs) leads, for In-In collisions at  $\sqrt{s_{NN}} = 17.3$  GeV, to  $\sigma_{cc} = 9.5 \pm 1.3(\text{stat.}) \pm 1.4(\text{syst.}) \mu\text{b}$  assuming kinematic distribution as in PYTHIA6
- Compatible with corresponding p-A measurements by NA50 and supporting the hypothesis of  $N_{\text{coll}}$  scaling

No other results available below top SPS energy



# $\chi_c$ measurements

- $\sim 25\%$  of the  $J/\psi$  comes from the  $\chi_c$  decay  
→  $\alpha(\chi_c)$  important to understand the  $J/\psi$  suppression



- $\chi_c$  not measured at SPS (no AA data)
- Available results at HERA-B, pA@ 920 GeV  
(large  $\chi_c$  sample:  $\sim 15000 \chi_c$   $-0.35 < x_F^{J/\psi} < 0.15$ )
- HERA-B observed no significant difference between  $\alpha(\chi_c)$  and  $\alpha(J/\psi)$   
→ similar “global” CNM effects on both resonances in the covered kinematical range (average value  $\Delta\alpha = 0.05 \pm 0.04$ ), but more accurate results are needed
- Non-trivial measurement, needs detection of low-momentum photon ( $< 1$  GeV)  
→ conversion or calorimetry




HERA-B, Phys.Rev.D79:012001,2009

# Thermal dileptons and chiral symmetry

# Elliptic flow of thermal dileptons

Vujanovic et al, Phys. Rev. C 98, 014902 (2018)


- ❑ **No measurements at present**
- ❑ Predictions at RHIC energies
  - ❑ LMR dominated by hadron gas: almost linear increase of  $v_2$  vs mass
  - ❑ IMR dominated by QGP: small  $v_2$



- ❑ No prediction at SPS energies
- ❑ Two possible scenarios:  $v_2=0$ 
  - ❑ **Measurement with uncertainty between 0.003 and 0.008**
- ❑  $v_2 = v_2^{\text{RHIC}}$ 
  - increase of  $v_2$  versus mass (HG) and a drop in the IMR (QGP)

# Fireball lifetime

- Thermal “excess” radiation in the mass region  $0.3 < M < 0.7 \text{ GeV}/c^2$ 
  - sensitive to all emission stages
  - tracks the **total fireball lifetime** within an accuracy of  $\sim 10\%$
- NA60 measurement, In-In at  $\sqrt{s_{NN}}=17.3 \text{ GeV}$  :  $\tau_{FB} = 8 \pm 1 \text{ fm}/c$






Black points → NA60+ projections  
Excellent accuracy

Soft mixed phase in a **first-order transition**  
→ pressure gradients in the system are small and thus stall the fireball expansion  
→ increased lifetime in the collision-energy regime where the mixed phase forms

# Open charm

# Open charm at low $\sqrt{s}$ in pA: nuclear PDFs

- Sensitivity to **nuclear PDFs in p-A** collisions
  - Probe EMC and anti-shadowing for  $\sqrt{s_{NN}} \sim 10\text{-}20$  GeV
  - Perform measurements with various nuclear targets to access the A-dependence of nPDF
- NA60+ offers a unique opportunity to investigate the **large  $x_{Bj}$  region** (study ratio to pA/pBe)
  - $0.1 < x_{Bj} < 0.3$  at  $Q^2 \sim 10\text{-}40$  GeV<sup>2</sup>




Lourenco, Wohri,  
Phys.Rept.433 (2006) 127


Eskola et al. , EPJ C77 (2017) 13

# Open charm in Pb-Pb: $R_{AA}$ and $v_2$

- Insight into **QGP transport properties**
  - Charm diffusion coefficient larger in the hadronic phase than in the QGP around  $T_c$
  - Hadronic phase represents a large part of the collision evolution at SPS energies
    - Sensitivity to hadronic interactions
    - Test models which predict strongest in-medium interactions in the vicinity of the quark-hadron transition
  - Measurement also important for precision estimates of diffusion coefficients at the LHC
  
- Study **charm thermalization at low  $\sqrt{s}$** 
  - Current measurements of HF-decay electron  $v_2$  at  $\sqrt{s_{NN}}=39$  and 62 GeV/c from RHIC
    - Smaller  $v_2$  than at  $\sqrt{s}=200$  GeV
    - Not conclusive on  $v_2 > 0$



Prino, Rapp, JPG43 (2016) 093002





STAR, PRC 95 (2017) 034907

# Open charm hadrochemistry

- Reconstruct different charm hadron species to get insight into **hadronization mechanism**

- Strange/non-strange** meson ratio ( $D_s/D$ ):


- $D_s/D$  enhancement expected in A-A collisions due to hadronisation via **recombination** in the strangeness rich QGP




STAR, PRL 127 (2021) 092301  
ALICE, PLB827 (2022) 136986

- Baryon/meson** ratios ( $\Lambda_c/D$ ):


- Expected to be enhanced in A-A in case of hadronisation via coalescence
- Interesting also in p-A since  $\Lambda_c/D^0$  in pp (p-Pb) at LHC is higher than in  $e^+e^-$



ALICE, PRL127 (2021) 202301



ALI-PUB-488630



ALI-PREL-321702






# Total charm cross section

- Total charm cross section in A-A collisions
  - Measured so far by NA60 in In-In collisions from intermediate-mass dimuons with 20% precision  
NA60, EPJ C59 (2009) 607
  - Upper limit from NA49 measurements of  $D^0$  mesons  
NA49, PRC73 (2006) 034910
- Precise measurement requires to reconstruct all meson and baryon ground states ( $D^0$ ,  $D^+$ ,  $D_s^+$  and  $\Lambda_c^+$  and their antiparticles)
- Charm cross section **ideal reference for charmonia**

# D-meson performance studies

## Fast simulations for central Pb-Pb collisions:


- D-meson signal simulation:  $p_T$  and  $y$  distributions from POWHEG-BOX+PYTHIA
- Combinatorial background:  $dN/dp_T$  and  $dN/dy$  of  $p$ ,  $K$  and  $\pi$  from NA49
- Parametrized simulation of VT detector resolution + track reconstruction with Kalman filter
- Reconstruct D-meson decay vertex from decay tracks
- Geometrical selections based on displaced decay vertex topology
  - For  $D^0$  in central Pb-Pb:
    - initial S/B  $\sim 10^{-7}$
    - $\rightarrow$  after selections S/B  $\sim 0.5$



# Towards a precise measurement of open charm at SPS energy

A measurement of **hadronic decays** is required

	Mass MeV	$c\tau$ ( $\mu\text{m}$ )	Decay	BR
$D^0$	1865	123	$K^-\pi^+$	3.95%
$D^+$	1869	312	$K^-\pi^+\pi^+$	9.38%
$D_s^+$	1968	147	$\phi\pi^+$	2.24%
$\Lambda_c^+$	2285	60	$pK^-\pi^+$	6.28%
			$pK_s^0$	1.59%
			$\Lambda\pi^+$	1.30%



# Next future

# Status & next future

- ❑ First prototype of a **MWPC module built and tested** at Weizmann institute
  - to be tested on a hadron beam at CERN in spring 2023
- ❑ R&D on **stitched MAPS** ongoing in the frame of a collaboration between ALICE and NA60+
- ❑ Toroidal **magnet prototype** built and tested
  - Mechanical and magnetic parameters under control
- ❑ Studies for LoI carried out with
  - ❑ **Fast simulation and reconstruction** tool
  - ❑ **FLUKA** calculations for background rates
- ❑ Finalize set-up
  - location of MWPC and GEM detectors
  - Define resolution for each station
- ❑ Build test mechanics with dummy sensors to investigate various aspects (cooling, alignment,...), continue R&D
- ❑ Extrapolate to full-scale magnet and start engineering design
- ❑ Define final sim/reco/analysis framework and investigate/develop related tools
  - GEANT4, ACTS, Aliroot framework,...