





International Workshop on Hadron Structure and Spectroscopy 2023

Transverse spin asymmetries in COMPASS Drell-Yan data

Małgorzata Niemiec
University of Warsaw, Poland
28 VI 2023



Collaboration:

24 institutes from **13** countries

Fixed target experiment

CERN SPS North-Area
(M2 beam-line)

Drell-Yan with 190 GeV π^- beam
and p^\uparrow NH₃, Al, W targets








DVCS and hard exclusive processes with
160 GeV μ^\pm beam and liquid H₂ target.

SIDIS with 160 GeV (200 GeV) μ^+ beam
and longitudinally/transversely - polarised
proton (NH₃) or deuteron (⁶LiD) target

Hadron spectroscopy with hadron beams
and nuclear targets.

Transverse Momentum Dependent PDFs

Nucleon spin structure described 8 twist-2 TMD PDFs.

		Nucleon Polarisation		
		U	L	T
Quark Polarisation	U	 $f_1^q(x, k_T^2)$ Number Density		 $f_{1T}^{q\perp}(x, k_T^2)$ Siverson
	L		 $g_1^q(x, k_T^2)$ Helicity	 $g_{1T}^{q\perp}(x, k_T^2)$ Worm-Gear T
	T	 $h_{1T}^{q\perp}(x, k_T^2)$ Boer-Mulders	 $h_{1L}^{q\perp}(x, k_T^2)$ Worm-Gear L	 $h_{1T}^{q\perp}(x, k_T^2)$ Transversity $h_{1T}^{q\perp}(x, k_T^2)$ Pretzelosity



This talk

$$f_{1T}^{q\perp}(x, k_T^2)$$

Siverson

$$h_{1T}^q(x, k_T^2)$$


Transversity

$$h_{1T}^{q\perp}(x, k_T^2)$$

Pretzelosity


Single polarised Drell-Yan process

Drell-Yan process in LO




$$\frac{d\sigma^{LO}}{d\Omega d^4q} \propto \left\{ 1 + D_{\sin^2\theta} \cos(2\varphi_{CS}) A_U^{\cos(2\varphi_{CS})} + |S_T| \left[\begin{array}{l} \sin(\varphi_S) A_T^{\sin(\varphi_S)} \\ D_{\sin^2\theta} \left(\begin{array}{l} \sin(2\varphi_{CS} + \varphi_S) A_T^{\sin(2\varphi_{CS} + \varphi_S)} \\ \sin(2\varphi_{CS} - \varphi_S) A_T^{\sin(2\varphi_{CS} - \varphi_S)} \end{array} \right) \end{array} \right] \right\}$$

$$A_{DY} \propto PDF_\pi \otimes PDF_N$$



Target rest frame



Collins-Soper frame

$$A_U^{\cos(2\varphi_{CS})} \propto h_{1,\pi}^{\perp q} \otimes h_{1,\pi}^{\perp q}$$

$$A_T^{\sin(\varphi_S)} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$$


$$A_T^{\sin(2\varphi_{CS} + \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1T,p}^{\perp q}$$

$$A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$$

Single polarised Drell-Yan process

The convolution of the TMDs runs over intrinsic transverse momenta

$$\mathcal{C} [w(\mathbf{k}_{T,\pi}, \mathbf{k}_{T,N}, \mathbf{q}_T) f_\pi f_N] = \frac{1}{N_c} \sum_q e_q^2 \int d^2\mathbf{k}_{T,\pi} d^2\mathbf{k}_{T,N} \delta^{(2)}(\mathbf{q}_T - \mathbf{k}_{T,\pi} - \mathbf{k}_{T,N}) \\ \times w(\mathbf{k}_{T,\pi}, \mathbf{k}_{T,N}, \mathbf{q}_T) \left[f_\pi^q(x_\pi, k_{T,\pi}^2) f_N^q(x_N, k_{T,N}^2) + f_\pi^q(x_\pi, k_{T,\pi}^2) f_N^q(x_N, k_{T,N}^2) \right]$$



Sivers function
and number density

$$A_T^{\sin(\varphi_S)} = \frac{-\mathcal{C} \left[\frac{\mathbf{q}_T \cdot \mathbf{k}_{T,N}}{q_T \cdot M_N} f_{1,\pi}^q f_{1T,N}^{q\perp} \right]}{\mathcal{C} \left[f_{1,N}^q f_{1,\pi}^q \right]}$$

Pretzelosity and
Boer-Mulders function

$$A_T^{\sin(2\varphi_{CS} + \varphi_S)} = \frac{-\mathcal{C} \left[\frac{2\mathbf{q}_T \cdot \mathbf{k}_{T,N} \left[2(\mathbf{q}_T \cdot \mathbf{k}_{T,\pi}) (\mathbf{q}_T \cdot \mathbf{k}_{T,N}) - q_T^2 (\mathbf{k}_{T,\pi} \cdot \mathbf{k}_{T,N}) \right] - q_T^2 k_{T,N}^2 (\mathbf{q}_T \cdot \mathbf{k}_{T,\pi})}{2q_T^3 M_\pi M_N^2} h_{1,\pi}^{q\perp} h_{1T,N}^{q\perp} \right]}{\mathcal{C} \left[f_{1,N}^q f_{1,\pi}^q \right]}$$

Transversity and
Boer-Mulders function

$$A_T^{\sin(2\varphi_{CS} - \varphi_S)} = \frac{\mathcal{C} \left[\frac{\mathbf{q}_T \cdot \mathbf{k}_{T,\pi}}{q_T \cdot M_\pi} h_{1,\pi}^{q\perp} h_{1,N}^q \right]}{\mathcal{C} \left[f_{1,N}^q f_{1,\pi}^q \right]}$$

Semi-inclusive Deep Inelastic Scattering

$$\frac{d\sigma^{LO}}{dx dy dz d^2p_T d\phi_h d\phi_\psi} \propto \left\{ \begin{array}{l} 1 + \cos(2\phi_h) \varepsilon A_{UU}^{\cos(2\phi_h)} \\ \sin(\phi_h + \phi_S) A_T^{\sin(\phi_h + \phi_S)} \\ \sin(\phi_h - \phi_S) A_T^{\sin(\phi_h - \phi_S)} \\ \sin(3\phi_h - \phi_S) A_T^{\sin(3\phi_h - \phi_S)} \end{array} \right\}$$


$$A_{SIDIS} \propto PDF_p \otimes FF$$

Drell-Yan process


$$\frac{d\sigma^{LO}}{d\Omega d^4q} \propto \left\{ \begin{array}{l} 1 + D_{\sin^2 \theta} \cos(2\varphi_{CS}) A_U^{\cos(2\varphi_{CS})} \\ \sin(\varphi_S) A_T^{\sin(\varphi_S)} \\ D_{\sin^2 \theta} \left(\begin{array}{l} \sin(2\varphi_{CS} + \varphi_S) A_T^{\sin(2\varphi_{CS} + \varphi_S)} \\ \sin(2\varphi_{CS} - \varphi_S) A_T^{\sin(2\varphi_{CS} - \varphi_S)} \end{array} \right) \end{array} \right\}$$

$$A_{DY} \propto PDF_\pi \otimes PDF_N$$




$A_{UU}^{\cos(2\phi_h)} \propto h_{1T,p}^{\perp q} \otimes H_{1q}^{\perp h}$	\longleftrightarrow Boer-Mulders	\longrightarrow	$A_U^{\cos(2\varphi_{CS})} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^{\perp q}$
$A_T^{(\phi_h - \phi_S)} \propto f_{1T,p}^{\perp q} \otimes D_{1q}^h$	\longleftrightarrow Sivers	\longrightarrow	$A_T^{\sin(\varphi_S)} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$
$A_T^{(3\phi_h - \phi_S)} \propto h_{1T,p}^{\perp q} \otimes H_{1q}^{\perp h}$	\longleftrightarrow Pretzelosity	\longrightarrow	$A_T^{\sin(2\varphi_{CS} + \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1T,p}^{\perp q}$
$A_T^{(\phi_h + \phi_S)} \propto h_{1,p}^q \otimes H_{1q}^{\perp h}$	\longleftrightarrow Transversity	\longrightarrow	$A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$



**Universality in TMD-QCD
parton model approach**



COMPASS DY setup






Drell-Yan setup

- 190 GeV π^- beam
- COMPASS spectrometer
- transversely polarised NH_3 target p^\uparrow
- Al, W targets
- hadron absorber

Drell-Yan measurement at COMPASS

- “Low mass” $\longleftrightarrow 1 < M_{\mu\mu}/(\text{GeV}/c^2) < 2$
 - large background and contamination
- “Intermediade mass” $\longleftrightarrow 2 < M_{\mu\mu}/(\text{GeV}/c^2) < 2.5$
 - high DY cross section
- “Charmonia mass” $\longleftrightarrow 2.5 < M_{\mu\mu}/(\text{GeV}/c^2) < 4.3$
 - strong J/ψ signal
 - good signal/background
- “High mass” $\longleftrightarrow 4.3 < M_{\mu\mu}/(\text{GeV}/c^2) < 8.5$
 - valence quark region
 - low background



“High mass” $\longleftrightarrow 4.3 < M_{\mu\mu}/(\text{GeV}/c^2) < 8.5$

- valence quark region
- low background





$$\langle x_\pi \rangle = 0.50$$

$$\langle x_N \rangle = 0.17$$

$$\langle q_T \rangle = 1.17 \text{ GeV}/c$$

$$\langle M_{\mu\mu} \rangle = 5.3 \text{ GeV}/c^2$$


Statistics in 2015 ~ 35 000 events
Statistics in 2018 ~ 37 000 events
Total ~ 71 000 events



Sivers: SIDIS and Drell-Yan TSA results


$$A_T^{\sin(\varphi_S)} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$$

Sivers TSA in Drell-Yan, 2015+2018 DATA



Sivers TSA in SIDIS


$$A_T^{(\phi_h - \phi_S)} \propto f_{1T,p}^{\perp q} \otimes D_{1q}^h$$



Pretzelosity: Drell-Yan and SIDIS TSA results

$$A_T^{\sin(2\varphi_{CS}+\varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1T,p}^{\perp q}$$


Pretzelosity TSA in Drell-Yan, 2015+2018 DATA



COMPASS, PLB 770 (2017) 130

Pretzelosity TSA in SIDIS


$$A_T^{(3\phi_h-\phi_S)} \propto h_{1T,p}^{\perp q} \otimes H_{1q}^{\perp h}$$



Transversity: Drell-Yan and SIDIS TSA results

$$A_T^{\sin(2\varphi_{CS}-\varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$$


Transversity TSA in Drell-Yan, 2015+2018 DATA






COMPASS, PLB 770 (2017) 130

Collins TSA in SIDIS

$$A_T^{(\phi_h + \phi_S)} \propto h_{1,p}^q \otimes H_{1q}^{\perp h}$$



Drell-Yan TSA results



Weighted TSAs in Drell-Yan: Introduction

Possible alternative: **weighting with powers of the transverse momentum**

Example: integration over $d^2\mathbf{q}_T$ with weight $= q_T/M_N$

$$A_T^{\sin\phi_S \frac{q_T}{M_N}} = \frac{2}{|\mathbf{S}_T|} \frac{\int d\phi_S d q_T^2 d\phi \sin\phi_S \frac{q_T}{M_N} d\sigma_{DY}}{\int d\phi_S d q_T^2 d\phi d\sigma_{DY}} = \frac{\int d^2\mathbf{q}_T \frac{q_T}{M_N} F_T^{\sin\phi_S}}{\int d^2\mathbf{q}_T F_U^1}$$

TMD framework

The n-th moment of TMD PDF

$$f^{(n)}(x) = \int d^2\mathbf{k}_T \left(\frac{k_T^2}{2M_h^2} \right)^n f(x, k_T^2)$$

Weighted **Sivers** asymmetry

$$A_T^{\sin\phi_S \frac{q_T}{M_N}} = -2 \frac{\sum_q e_q^2 [f_{1,\pi}^{\bar{q}}(x_\pi) f_{1T,N}^{q(1)\perp}(x_N) + (q \leftrightarrow \bar{q})]}{\sum_q e_q^2 [f_{1,\pi}^{\bar{q}}(x_\pi) f_{1,N}^q(x_N) + (q \leftrightarrow \bar{q})]}$$

Weighted asymmetry induced by proton **pretzelosity** and pion **Boer-Mulders** function


$$A_T^{\sin(2\varphi+\varphi_S) \frac{q_T^3}{2M_\pi M_N^2}} = -6 \frac{\sum_q e_q^2 [h_{1,\pi}^{\bar{q}(1)\perp}(x_\pi) h_{1T,N}^{q(2)\perp}(x_N) + (q \leftrightarrow \bar{q})]}{\sum_q e_q^2 [f_{1,\pi}^{\bar{q}}(x_\pi) f_{1,N}^q(x_N) + (q \leftrightarrow \bar{q})]}$$

Weighted asymmetry induced by proton **transversity** and pion **Boer-Mulders** function

$$A_T^{\sin(2\varphi-\varphi_S) \frac{q_T}{M_\pi}} = -2 \frac{\sum_q e_q^2 [h_{1,\pi}^{\bar{q}(1)\perp}(x_\pi) h_{1,N}^q(x_N) + (q \leftrightarrow \bar{q})]}{\sum_q e_q^2 [f_{1,\pi}^{\bar{q}}(x_\pi) f_{1,N}^q(x_N) + (q \leftrightarrow \bar{q})]}$$



Sivers: Drell-Yan and SIDIS weighted TSA results

Sivers
weighted TSA
in Drell-Yan,
2015 + 2018
DATA




Sivers weighted TSA in SIDIS

[COMPASS, Nucl.Phys. B940 (2019) 34]



Transversity weighted TSA in Drell-Yan, 2015+2018 DATA



Combined conventional Drell-Yan TSA

Sivers found to be positive, $\sim 1\sigma$ away from zero

Transversity found to be negative, $\sim 2\sigma$ away from zero

Pretzelosity found to be small and compatible with zero

Transverse momentum weighted asymmetries

A way to overcome the convolution over intrinsic k_T

A direct access to the k_T^2 moments of TMD PDFs

Combined Drell-Yan weighted TSA

Result consistent with conventional TSAs

Prospects

Pretzelosity weighted TSA study ongoing

Extraction of first k_T^2 -moment of Sivers

A close-up view of an ornate astronomical clock face. The clock is circular with a blue background and gold-colored markings. It features concentric rings of Roman numerals and zodiac signs. A central black sphere is mounted on a metal structure, with a golden sunburst ornament above it. The clock is set against a dark, textured background.

Thank you for attention!